

FUNGI-
FOR ECONOMY
DO UNDER NO. 7001
COMMENTS



BEST MALASPRAY 8

ACTIVE INGREDIENTS:	BY WT.
Malathion* - - - - -	80.0%
Aromatic Petroleum Solvent - - -	6.9%
INERT INGREDIENTS - - - - -	13.1%
TOTAL 100.0%	

*0,0-dimethyl dithiophosphate of diethyl mercaptosuccinate.

U.S.D.A. Registration No. 7001-7

CONTAINS 8 LBS. MALATHION PER GALLON

DIRECTIONS FOR USE

NOTE: The rates of water necessary to add to the tank will vary with the application method. The normal requirements for the following application methods are as follows: AIR APPLICATION — 5 to 20 gals. per acre. GROUND APPLICATION — 15 to 200 gals. per acre. ORCHARD APPLICATION — 70 to 600 gals. per acre. (Citrus may require up to 1,500 gals.) (70 to 100 gals. per acre rates are considered concentrate rates and require special concentrate equipment.)

- BEANS:** Lygus Bug, Aphid. 1 to 1-3/4 Pts./Acre. Do not apply within 1 day of harvest.
- BROCCOLI, BRUSSELS SPROUTS, CABBAGE, CAULIFLOWER:** Aphid, Cabbage Looper, Imported Cabbage Worm. 1 to 2-1/2 Pts./Acre. Do not apply within 3 days of harvest on Broccoli, or within 7 days on Cabbage, Brussels Sprouts or Cauliflower.
- LETTUCE, SPINACH:** Aphid, Leafhopper. 1 to 2 Pts./Acre. Do not apply within 14 days of harvest on Leaf Lettuce, or within 7 days of harvest on Head Lettuce or Spinach.
- ONIONS (Including Green Onions):** Onion Thrip. 1 to 2 Pts./Acre. Do not apply within 3 days of harvest on Green Onions.
- PEPPERS:** Aphid. 3/4 to 1-1/2 Pts./Acre. Do not apply within 3 days of harvest.
- COTTON:** Aphid, Leafhopper, Lygus Bug, Leaf Perforator, Leafworm. 1 to 4 Pts./Acre. Do not use treated seed as food or feed.
- CELERY:** Aphid, Leafhopper, Lygus Bug. 1 to 1-1/2 Pts./Acre. Do not apply within 7 days of harvest.
- SAFFLOWER:** Aphid, Lygus Bug, Thrips. 1-1/4 Pts./Acre. Do not apply within 3 days of harvest.
- WALNUTS:** Walnut Husk Fly. 2 Pts./100 gals. water. No limitations.
- EGGPLANT:** Aphid, Lace Bug, Spider Mite. 1-1/2 to 3-1/2 Pts./Acre. Do not apply within 3 days of harvest.
- PEAS:** Pea Aphid, Lygus Bug, Leafhopper, Thrips, Pea Weevil. 1 to 2-1/2 Pts./Acre. Do not apply within 3 days of harvest.
- CORN:** Aphid, Corn Rootworm Adults, Sap Beetle, Thrips, Young Grasshopper. 1 Pt./Acre. Do not apply within 5 days of harvest for forage use. CAUTION: Injury may occur in the whorls and silk stages.
- HOPS:** Aphids. 3/4 to 1-1/4 Pts./Acre. Do not apply within 7 days of harvest.

- CUCUMBERS, SQUASH, PUMPKINS, MELONS:** Aphids, Spider Mite, Pickleworm, Leafhopper, Cucumber Beetle, Darkling Ground Beetle, Squash Vine Borer, Thrips. 1-3/4 Pts./Acre. Do not apply on Pumpkins within 3 days of harvest. Do not apply to Cucumbers, Squash or Melons within 1 day of harvest. Do not apply unless plants are dry.
- POTATOES:** Potato Aphid, Blister Beetle, False Chinch Bug, Leafhopper, Mealy Bug. 1 to 3 Pts./Acre. No time limitations.
- TOMATOES:** Aphid, Spider Mite. 1 to 2 Pts./Acre. Do not apply within 1 day of harvest.
- ALFALFA, CLOVER:** Aphid, Leafhopper, Spider Mite, Alfalfa Weevil Larvae, Armyworm, Clover Leaf Weevil, Grasshopper, Lygus Bug, Vetch Bruchid, Spittle Bug. 1 to 1-1/2 Pts./Acre. No time limitations. Do not apply to crops in bloom.
- APPLES:** Red-banded Leafroller, Woolly Apple Aphid, Forbes Scale, Bud Moth, Spider Mite, Mealy Bug. 1-1/2 Pts./Acre per 100 gals. water. Do not apply within 3 days of harvest. May cause injury to McIntosh and Cortland varieties if applied within 4 weeks of harvest.
- CHERRIES:** Black Cherry Aphid, Leafroller, Fruit Fly. 1 Pt./100 gals. water. Do not apply within 3 days of harvest. May injure foliage on some varieties.
- PEARS:** Red-banded Leafroller, Pear Psylla, Aphid, Woolly Aphid, Mealy Bug, Spider Mite, Leafroller. 1 to 1-1/2 Pts./Acre per 100 gals. water. Make one or more applications as needed. Do not apply within 1 day of harvest.
- PEACHES:** Spider Mite, Aphid, Lygus Bug, Bud Moth. 1-1/2 Pts./100 gals. water. Full coverage spray. Do not apply within 7 days of harvest.
- PLUMS, PRUNES:** Mealy Plum Aphid, Bud Moth, Leafhopper. 3/4 Pt./100 gals. water. Make one or more applications as needed. Do not apply within 3 days of harvest.
- GRAPES:** Grape Leafhopper (California and southwest), Spider Mite, Mealy Bug. 1 Pt./100 gals. water. Do not apply within 3 days of harvest. Injury may occur to Ribier, Italia, Cardinal or Almiria varieties after clusters appear.
- STRAWBERRIES:** Aphid, Thrips, Spider Mite, Leafhopper, Leafroller, Strawberry Root Weevil. 1 to 2 Pts./Acre. Do not apply within 3 days of harvest.
- CANE BERRIES (Blackberries, Boysenberries, Loganberries, Raspberries):** Spider Mite, Thrips, Leafhopper. 1 Pt./100 gals. water. Do not apply within 1 day of harvest.
- CITRUS (Lemons, Oranges, Grapefruit, Limes, Tangerines):** Aphid, Thrips, Red Scale, Yellow Scale, Purple Scale, Black Scale (single brooded), Soft Scale, Citricola Scale. 3/4 to 1 Pt./100 gals. water. Do not apply when trees are in bloom. Do not apply within 7 days of harvest.
- BEETS (Sugar & Table):** Aphid, Thrips, Leafminer, Leafhopper, Grasshopper. 1 to 2-1/2 Pts./Acre. Do not apply within 7 days of harvest if tops are to be used as food or feed.
- ALFALFA SEED, CLOVER SEED:** Aphid, Alfalfa Weevil, Lygus Bug. 1-1/2 to 2 Pts./Acre. Do not apply within 7 days of harvest. Do not apply during full bloom.
- ORNAMENTALS:** Aphid, Spider Mite, White Flies, Thrips, Mealy Bug, Leafhopper, Tent Caterpillar, Lace Bug, Oyster Shell Scale, Tarnished Plant Bug, Leafminer. 1 to 2 Pts./100 gals. water. Do not use on Boston, Maidenhair and Pteris ferns, and some varieties of Grassula.

CA
HARMFU
AVOID PROL
SP
AVOID PROL
CONT

KEEP OUT OF
Wash Tho
AVOID CONTAMINATION

Do not allow this product
crops as residues may cor
unfit for sale, feed or hu
food or dairy animals, stre

THIS PRODUCT

THE BEST F

Division of Occiden
P. O. BOX 198 • LATHROP,

CONDITIONS OF SALE: 1. Sell
ingredient specified and is read
No one, other than an officer
warranty, guarantee or direction
time, place, rate of application
Seller's control, Seller's liability
product is limited to rep aceme

ACCEPTED WITH COMMENTS

2 LOTS

BEST MALASPRAY 8 EC

BY WT.

- - 80.0%

- - 6.9%

- - 13.1%

TOTAL 100.0%

7001-7

PER GALLON

FOR USE

Apply with the application method. The normal rate is: AIR APPLICATION — 5 to 20 gals. per acre. ORCHARD APPLICATION — 70 to 600 gals. per acre. Concentrate rates are considered concentrate rates.

Apply within 1 day of harvest.

Aphid, Cabbage Looper, Imported Cabbage Moth — Broccoli, or within 7 days on other crops.

Do not apply within 14 days of harvest on Leafhopper.

Do not apply within 3 days of harvest on other crops.

Days of harvest.

Form. 1 to 4 Pts./Acre. Do not use treated seed.

Do not apply within 7 days of harvest.

Do not apply within 3 days of harvest.

Days.

per acre. Do not apply within 3 days of harvest.

1 to 2-1/2 Pts./Acre. Do not apply within 3 days of harvest.

Grasshopper. 1 Pt./Acre. Do not apply within 7 days of harvest.

Days of harvest.

CUCUMBERS, SQUASH, PUMPKINS, MELONS: Aphids, Spider Mite, Pickleworm, Leafhopper, Cucumber Beetle, Darkling Ground Beetle, Squash Vine Borer, Thrips. 1-3/4 Pts./Acre. Do not apply on Pumpkins within 3 days of harvest. Do not apply to Cucumbers, Squash or Melons within 1 day of harvest. Do not apply unless plants are dry.

POTATOES: Potato Aphid, Blister Beetle, False Chinch Bug, Leafhopper, Mealy Bug. 1 to 3 Pts./Acre. No time limitations.

TOMATOES: Aphid, Spider Mite. 1 to 2 Pts./Acre. Do not apply within 1 day of harvest.

ALFALFA, CLOVER: Aphid, Leafhopper, Spider Mite, Alfalfa Weevil Larvae, Armyworm, Clover Leaf Weevil, Grasshopper, Lygus Bug, Vetch Bruchid, Spittle Bug. 1 to 1-1/2 Pts./Acre. No time limitations. Do not apply to crops in bloom.

APPLES: Red-banded Leafroller, Woolly Apple Aphid, Forbes Scale, Bud Moth, Spider Mite, Mealy Bug. 1-1/2 Pts./Acre per 100 gals. water. Do not apply within 3 days of harvest. May cause injury to McIntosh and Cortland varieties if applied within 4 weeks of harvest.

CHERRIES: Black Cherry Aphid, Leafroller, Fruit Fly. 1 Pt./100 gals. water. Do not apply within 3 days of harvest. May injure foliage on some varieties.

PEARS: Red-banded Leafroller, Pear Psylla, Aphid, Woolly Aphid, Mealy Bug, Spider Mite, Leafroller. 1 to 1-1/2 Pts./Acre per 100 gals. water. Make one or more applications as needed. Do not apply within 1 day of harvest.

PEACHES: Spider Mite, Aphid, Lygus Bug, Bud Moth. 1-1/2 Pts./100 gals. water. Full coverage spray. Do not apply within 7 days of harvest.

PLUMS, PRUNES: Mealy Plum Aphid, Bud Moth, Leafhopper. 3/4 Pt./100 gals. water. Make one or more applications as needed. Do not apply within 3 days of harvest.

GRAPES: Grape Leafhopper (California and southwest), Spider Mite, Mealy Bug. 1 Pt./100 gals. water. Do not apply within 3 days of harvest. Injury may occur to Ribier, Italia, Cardinal or Almiria varieties after clusters appear.

STRAWBERRIES: Aphid, Thrips, Spider Mite, Leafhopper, Leafroller, Strawberry Root Weevil. 1 to 2 Pts./Acre. Do not apply within 3 days of harvest.

CANE BERRIES (Blackberries, Boysenberries, Loganberries, Raspberries): Spider Mite, Thrips, Leafhopper. 1 Pt./100 gals. water. Do not apply within 1 day of harvest.

CITRUS (Lemons, Oranges, Grapefruit, Limes, Tangerines): Aphid, Thrips, Red Scale, Yellow Scale, Purple Scale, Black Scale (single brooded), Soft Scale, Citricola Scale. 3/4 to 1 Pt./100 gals. water. Do not apply when trees are in bloom. Do not apply within 7 days of harvest.

BEETS (Sugar ^{beet} & Table ^{beets}): Aphid, Thrips, Leafminer, Leafhopper, Grasshopper. 1 to 2-1/2 Pts./Acre. Do not apply within 7 days of harvest if tops are to be used as food or feed.

ALFALFA SEED, CLOVER SEED: Aphid, Alfalfa Weevil, Lygus Bug. 1-1/2 to 2 Pts./Acre. Do not apply within 7 days of harvest. Do not apply during full bloom.

ORNAMENTALS: Aphid, Spider Mite, White Flies, Thrips, Mealy Bug, Leafhopper, Tent Caterpillar, Lace Bug, Oyster Shell Scale, Tarnished Plant Bug, Leafminer. 1 to 2 Pts./100 gals. water. Do not use on Boston Maidenhair and Pteris ferns, and some varieties of Crassula.

CAUTION

HARMFUL IF SWALLOWED!

AVOID PROLONGED BREATHING OF SPRAY OR MIST

AVOID PROLONGED OR REPEATED CONTACT WITH SKIN

KEEP OUT OF REACH OF CHILDREN

Wash Thoroughly After Using

AVOID CONTAMINATION OF FEED AND FOODSTUFFS

Do not allow this product to drift onto adjoining food or feed crops as residues may contaminate these crops making them unfit for sale, feed or human consumption. Keep clear of food or dairy animals, streams, fish ponds and wildlife areas.

THIS PRODUCT IS POISONOUS TO FISH

26:10/66

THE BEST FERTILIZERS CO.

Division of Occidental Petroleum Corporation

P. O. BOX 198 • LATHROP, CALIFORNIA 95330 • (209) 858-2511

CONDITIONS OF SALE: 1. Seller warrants that this product consists of the ingredients specified and is reasonably fit for use as directed on the label. No one, other than an officer of Seller, is authorized to make any other warranty, guarantee or direction concerning this product. 2. Because the time, place, rate of application and other conditions of use are beyond Seller's control, Seller's liability from the handling, storage and use of this product is limited to replacement of product or refund of purchase price.