



BEST BBC 12

FOR CONTROL OF PLANT PARASITIC NEMATODES

ACTIVE INGREDIENTS:	BY WT.
1, 2 - Dibromo - 3 - Chloropropane - - -	81.7%
Other Halogenated C ₃ Compounds - - -	4.3%
INERT INGREDIENTS - - - - -	14.0%
	TOTAL 100.0%

U.S.D.A. Reg. No. 7001-10
**BEST BBC 12 CONTAINS 12 LBS.
 BBC SOIL FUMIGANT PER GALLON**

RECOMMENDED APPLICATION RATES

- VEGETABLES** (Broccoli, Brussels Sprouts, Cabbage, Cauliflower, Carrots, Eggplant, Endive, Lettuce, Okra, Parsnips, Radishes and Snap Beans):
 PREPLANT, AT PLANTING, POSTPLANT. Apply postplant as soon as plants become firmly established. 1 1/2 to 2 gals. per acre.
- CUCURBITS** (Cucumbers, Melons, Summer Squash):
 PREPLANT, AT PLANTING. 1-1/2 to 2 gals. per acre.
- TOMATOES:**
 PREPLANT, DIRECT SEEDED. 1 to 2 gals. per acre.
 AT PLANTING, DIRECT SEEDED. 1/2 to 3/4 gal. per acre.
 Preplanting transplants, wait 2 weeks before planting. 1 gal. per acre.
- COTTON:**
 PREPLANT, AT PLANTING, POSTPLANT. 1 to 2 gals. per acre.
- PINEAPPLES:**
 PREPLANT, AT PLANTING, POSTPLANT. Do not apply within 9 months of harvest. 3 to 11 gals. per acre.
- BERRIES** (Blackberries, Boysenberries, Dewberries, Loganberries, Raspberries):
 PREPLANT, AT PLANTING, POSTPLANT. Do not apply within 48 days of harvest. 2 gals. per acre. Strawberries, do not apply within 300 days of harvest. 2 gals. per acre.
- CITRUS FRUIT:**
 PREPLANT, AT PLANTING, POSTPLANT. Western states only. Do not apply more than once every three years. 3 to 7 gals. per acre.
- DECIDUOUS FRUITS** (Apricots, Grapes, Nectarines, Peaches):
 PREPLANT, AT PLANTING, POSTPLANT. 3 to 7 gals. per acre.
 FIGS: 3 to 5 gals. per acre. Do not apply within 1 year of harvest.
- NUTS:** English Walnuts:
 PREPLANT, AT PLANTING, POSTPLANT. 3 to 7 gals. per acre. Do not apply within 100 days of harvest.
- PEANUTS:**
 PREPLANT, AT PLANTING. 1 1/2 to 2 gals. per acre. Do not feed, sell, or introduce into commerce hay or hulls taken from treated soil. Forage crops grown on treated soil should not be used as feed for dairy animals or animals being finished for slaughter until two years after row treatments or three years after overall treatments.
- SYMPHYLANS:**
 To control SYMPHYLANS use only overall applications at the dosages indicated. Apply during late summer or early fall when the soil is warm.

WARNING! KEEP OUT OF REACH OF CHILDREN

HARMFUL LIQUID AND VAPOR — COMBUSTIBLE MIXTURE — MAY BE ABSORBED THROUGH SKIN CAUSES SKIN IRRITATION AND BLISTERS ON PROLONGED CONTACT

DO NOT BREATHE VAPOR — KEEP CONTAINER CLOSED — DO NOT TAKE INTERNALLY

WASH THOROUGHLY AFTER USING — DO NOT GET IN EYES, ON SKIN OR ON CLOTHING

Use Only in a Well-Ventilated Area. Keep Away from Heat and Open Flame, Feed and Foodstuffs.

In case of contact, immediately remove contaminated shoes and clothing and wash skin with soap and water. Flush eyes with plenty of water for at least 15 minutes and get medical attention. Do not wear shoes or clothing until absolutely free of all chemical odor.

If illness results from inhalation, remove to fresh air and call a doctor.

If swallowed, call a doctor. Induce vomiting by giving an emetic such as 2 tablespoons of table salt in a glass of warm water.

To Protect Fish and Wildlife, Do Not Contaminate Streams, Lakes, or Ponds with This Material
 Do Not Feed Plant Remains to Dairy or Meat Animals

GENERAL DIRECTIONS

Apply when soil temperatures at a depth of 6 to 8 inches are between 40° and 80° F. The soil should be in good tilth and contain adequate moisture for good seed germination. Extremely wet or dry soils should not be treated. Application on commercial acreages should be made with equipment designed for even distribution at specified depth.

PREPLANT TREATMENT: OVERALL: Apply evenly in rows 12 inches apart at a depth of 6 to 8 inches taking care to seal chisel channels with a drag or press wheel. **ROW:** Apply in the rows to be planted at a depth of 6 to 8 inches, preferably with two shanks spaced 12 inches apart, placing the fumigant 6 inches to each side of the seed row. Care should be taken to seal chisel channels with a press wheel.

DRENCH TREATMENT: For either preplant or postplant applications in irrigation water feed evenly in the water flow in order to make sure that all irrigated areas receive equal treatment. This product is also recommended for use at the rate of 30 ppm of the active ingredient in sufficient water to give penetration to the desired depth. One gallon in 48,000 gallons of water is 30 ppm.

MECHANICAL MIXING IS REQUIRED — RECOMMENDED FOR USE ON MINERAL SOILS ONLY

FOR PRE-PLANT APPLICATIONS: A 7- to 14-day exposure period on shallow rooted crops and a 4- to 6-weeks exposure period on deeprooted trees and shrubs should be allowed.

RECOMMENDED APPLICATION RATES ARE FOR OVERALL TREATMENT
 Row application should be in proportion to overall applications

DO NOT EXCEED OVERALL PER ACRE RECOMMENDATION IN A SINGLE YEAR

MIXING DIRECTIONS

This product can be used "as is" or further diluted with mineral spirits. Dilute sufficiently so that the application equipment will deliver the proper amount of Soil Fumigant per acre.

Avoid Spillage When Mixing. When Container is Empty Immediately Wash Thoroughly with Water and Detergent and Destroy. Never Re-Use Container.

1-67

OCCIDENTAL CHEMICAL COMPANY

DIVISION OF OCCIDENTAL PETROLEUM CORPORATION
 LATHROP, CALIFORNIA

CONDITIONS OF SALE: 1. Seller warrants that this product consists of the ingredients specified and is reasonably fit for use as directed on the label. No one, other than an officer of Seller, is authorized to make any other warranty, quote fee, or direction concerning this product. 2. Because of the time, place, rate of application and other conditions of use, Seller's control of responsibility from the handling, storage and use of this product is limited to replacement of product or refund of purchase price.