

ACCEPTED

July-20-1967

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND ROENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
UNDER NO. 7001-46

And Get Results With

BEST

Comment

BEST MAG-CHLOR* DEFOLIANT

ACTIVE INGREDIENT:	BY WT.
Sodium Chlorate -----	19.3%
INERT INGREDIENTS: -----	80.7%
TOTAL	100.0%

* Contains 2 lbs. Magnesium Chloride as a fire retardant

U.S.D.A. Registration No. 7001-46

DIRECTIONS FOR USE

COTTON DEFOLIATION: Use 2½ gallons per acre in 10 gallons of water by air. 20 to 30 gallons of water by ground sprayer. Make first application 2 to 3 weeks before anticipated picking date. Use no more than 2 applications. Apply not later than 7 days before harvest. Keep animals off treated area. Do not feed treated gin trash to livestock.

NOTE: It is recommended that a spreader-activator be used with this product in order to aid in obtaining uniform coverage. Consult the label of the product selected for manufacturers instructions.

THE BEST FERTILIZERS CO.

DIVISION OF OCCIDENTAL PETROLEUM CORPORATION
P. O. BOX 198 • LATHROP, CALIFORNIA 95330 • (209) 858-2511

CONDITIONS OF SALE: 1. Seller warrants that this product consists of the ingredients specified and is reasonably fit for use as directed on the label. No one, other than an officer of Seller, is authorized to make any other warranty, guarantee or direction concerning this product. 2. Because the time, place, rate of application and other conditions of use are beyond Seller's control, Seller's liability from the handling, storage and use of this product is limited to replacement of product or refund of purchase price.

WARNING

Harmful if swallowed. Avoid skin contact. Wash clothing and spray equipment after handling. Avoid contamination of feed and foodstuffs. Keep containers tightly closed.

KEEP OUT OF REACH OF CHILDREN
AND ANIMALS

Do not allow drift of this material onto other crops or pasturelands. Do not apply during high mid-day temperatures or extremely cold or windy conditions. A second application may be necessary to complete defoliation on rank growth or if plants to be defoliated are not at proper stage for complete defoliation to occur.

Avoid Spillage When Mixing. When Container Is Empty Immediately Wash Thoroughly and Destroy.
NEVER RE-USE!

96:12/66

APPLICATION SUGGESTIONS: For best results in cotton defoliation the following conditions should be present:

Temperature High
Plants Mature - Growth stopped
Complete Coverage
Large Droplet Size
Low Nitrogen Condition at Time of Defoliation
Soil Moisture Moderate (Neither Very High or Very Low)
Uniform Plant Population
Insects and Weeds Controlled
Uniform Land (No Saline or Alkaline Areas)

ACCEPTED WITH COMMENTS