

BEST CHLORDANE 8 EC

ACTIVE INGREDIENTS:	BY WT.
Chlordane*	72%
Aromatic Petroleum Solvents	18%
INERT INGREDIENTS	10%
TOTAL 100%	

*Equivalent to 43.2% Octachloro-4,7-methanotetrahydrochlordane and 28.8% related compounds.

CONTAINS 8 LBS. OF CHLORDANE PER GALLON

U.S.D.A. 7001-57

DIRECTIONS FOR USE

NOTE: The rates of water necessary to add to the tank will vary with the application method. The normal application rates for the following application methods are as follows: AIR APPLICATION — 5 to 20 gals. per acre. GROUND APPLICATION — 15 to 200 gals. per acre.

Beans, Broccoli, Kale, Brussels Sprouts, Celery, Cabbage, Cauliflower, Collards, Radishes, Strawberries, Cucumbers, Peppers:

LEAFMINERS, THRIPS, SOWBUGS, SQUASH BUGS, TOMATO HORNWORMS, ARMYWORMS, CLIMBING CUTWORMS, LYGUS BUGS, STINK BUGS, CABBAGE LOOPERS, FLEA BEETLES, DIAMOND-BACK MOTHS, COLORADO POTATO BEETLES: 1 Quart per acre. Do not apply to Green or Snap Beans after pods begin to form. Do not apply to Celery after plants are half grown or start to bunch. Do not apply to Broccoli, Brussels Sprouts, Cabbage, Cauliflower, Kale, Collards, Cucumbers, Peppers, Radishes or Strawberries after edible parts start to form. May be injurious to some varieties of Cucumbers.

Tomatoes:


LEAFMINERS, THRIPS, SOWBUGS, SQUASH BUGS, TOMATO HORNWORMS, ARMYWORMS, CLIMBING CUTWORMS, LYGUS BUGS, STINK BUGS, CABBAGE LOOPERS, FLEA BEETLES, DIAMOND-BACK MOTHS, COLORADO POTATO BEETLES: 2½ Quarts per acre. Do not apply to Tomatoes after edible parts start to form.

Carrots:

LEAFMINERS, THRIPS, SOWBUGS, SQUASH BUGS, TOMATO HORNWORMS, ARMYWORMS, CLIMBING CUTWORMS, LYGUS BUGS, STINK BUGS, CABBAGE LOOPERS, FLEA BEETLES, DIAMOND-BACK MOTHS, COLORADO POTATO BEETLES: 1 Quart per acre. No time limitation.

Potatoes:

LEAFMINERS, THRIPS, SOWBUGS, SQUASH BUGS, TOMATO HORNWORMS, ARMYWORMS, CLIMBING CUTWORMS, LYGUS BUGS, STINK BUGS, CABBAGE LOOPERS, FLEA BEETLES, DIAMOND-BACK MOTHS, COLORADO POTATO BEETLE: 1 Pint per acre. No time limitation. For control of WIREWORMS and CUTWORMS: 5 Quarts per acre. Preplant soil application or at time of planting.



Cotton:

COTTON FLEA HOPPER, CUTWORMS, FALL ARMYWORM, FLEA BEETLES: 1 Quart per acre. Do not apply after bolls open. Do not graze treated fields by meat or dairy animals.

Lettuce:

CORN EARWORM, FLEA BEETLE, LEAFMINERS: 1 Quart per acre. Do not apply after edible parts start to form.

Lettuce:

WIREWORMS, WHITE GRUBS, CUTWORMS, DIABROTICA LARVAE: 1¼ Gallons per acre. Work into upper few inches of soil as a preplant treatment.

Melons:

WIREWORMS, CUTWORMS: 1¼ Gallons per acre. Preplant soil application, work into upper few inches of soil.

WARNING! KEEP OUT OF REACH OF CHILDREN

HARMFUL IF SWALLOWED

Contact with Skin Can Cause Toxic Symptoms. In Case of Contact with Skin Wash with Soap and Water

AVOID BREATHING SPRAY MIST

AVOID CONTAMINATION OF FEED AND FOODSTUFFS

Do not allow this product to drift onto adjoining food or feed crops as residues may contaminate these crops making them unfit for sale, feed or human consumption. Keep clear of food or dairy animals, streams, fish ponds and wildlife areas.

Avoid spillage while mixing and filling spray equipment. When container is empty, immediately wash thoroughly with water and detergent and destroy. Discard in a safe place; never re-use container.

2:11/68

OCCIDENTAL CHEMICAL COMPANY

DIVISION OF OCCIDENTAL PETROLEUM CORPORATION

LATHROP, CALIFORNIA

CONDITIONS OF SALE: 1. Seller warrants that this product consists of the ingredients necessarily fit for use as directed on the label. No one, other than an official representative, shall make any other warranty, guarantee or direction concerning this product. The rate of application and other conditions of use are beyond Seller's control. The handling, storage and use of this product is limited to replacement.