

BEST BASIC COPPER "53"

(Basic Copper Sulphate)

7001-66

JUL 11 1966



DIRECTIONS FOR USE

NOTE: The rates of water necessary to add to the tank will vary with the application method. The normal requirements for the following application methods are as follows: AIR APPLICATION — 5 to 20 gals. per acre. GROUND APPLICATION — 15 to 200 gals. per acre. ORCHARD APPLICATION — 70 to 600 gals. per acre. (Citrus may require up to 1,500 gals.) 170 to 100 gals. per acre rates are considered concentrate rates and require special concentrate equipment.)

ALMONDS: Brown Rot, Shot-hole Fungus. 4 to 5 lbs. plus 1/2 to 1 gal. Best Dormant Flowable Oil per 100 gals. water. For Brown Rot Control apply in popcorn stage. Apply in petal fall or jacket stage for shot-hole fungus.

APRICOTS: Brown Rot Fungus, Shot-hole Fungus. 4 to 5 lbs. plus 1/2 to 1 gal. Best Dormant Flowable Oil per 100 gals. water. Apply in red bud to popcorn stage. Brown Rot may also require a jacket spray, especially during wet years.

CHERRIES: Brown Rot, Blossom Rot. 4 to 5 lbs. plus 1/2 to 1 gal. Best Dormant Flowable Oil per 100 gals. water. Apply in popcorn stage.

PEACHES, NECTARINES: Peach Blight (shot-hole), Peach Leaf Curl. 4 to 5 lbs. plus 1/2 to 1 gal. Best Dormant Flowable Oil per 100 gals. water from Nov. to Dec. 15. Brown Rot, Peach Blight (shot-hole) - apply in pink bud stage using 4 to 5 lbs. plus 1/2 to 1 gal. Best Dormant Flowable Oil.

PRUNES, PLUMS: Blossom Rot, Brown Rot, Shot-hole. 4 to 5 lbs. plus 1/2 to 3/4 gal. Best Dormant Flowable Oil. Apply in early popcorn stage.

CUCUMBERS: Downy Mildew, Anthracnose, Angular Leaf Spot. 10 lbs. per acre. Begin application before disease appears and repeat at 7 to 10 day intervals.

ONIONS: Downy Mildew. 10 lbs. per acre. Begin application before disease appears and repeat at 7 to 10 day intervals.

ACTIVE INGREDIENT:	BY WT.
Copper expressed as Metallic (in Basic Copper Sulfate) - - - - -	53%
INERT INGREDIENTS - - - - -	47%
	TOTAL 100%

MIXING DIRECTIONS

With tank 1/4 full of water and with agitator running, add the amount of Copper required. When tank is two-thirds full, add the oil spreader and fill tank. Keep agitator running during filling and spraying operations.

BEST BASIC COPPER "53" is compatible with spray oils and can be included with higher percentages of Dormant Spray Oils.

CAUTION

KEEP OUT OF REACH OF CHILDREN

HARMFUL IF TAKEN INTERNALLY

AVOID INHALING DUST OR SPRAY MIST

DO NOT APPLY ON TENDER FOLIAGE OF PEACHES, ALMONDS, APRICOTS, AND JAPANESE PLUMS

170:1/68

OCCIDENTAL CHEMICAL COMPANY

OCCIDENTAL CHEMICAL CORPORATION
LATHROP, CALIFORNIA


NOTICE: Occidental Chemical Company warrants that this product is a pure, chemically synthesized product and is not a by-product of any other process. The name "Occidental" is a registered trademark of Occidental Chemical Company. Occidental Chemical Company is authorized to make any other warranty or guarantee for this product. This product is sold as a concentrate and the rate of application and other conditions of use are beyond the liability of Occidental Chemical Company. The handling, storage and use of this product is limited to replacement of product or refund of purchase price.

BEST BASIC COPPER "53"

(Basic Copper Sulphate)

7001-66

Jul 11 1968
UNDER THE FEDERAL RESISTANCE AND REPRESSION ACT FOR ECONOMIC POWERS OF THE UNITED STATES TO REVENUE COMMISSION



DIRECTIONS FOR USE

NOTE: The rates of water necessary to add to the tank will vary with the application method. The normal requirements for the following application methods are as follows: AIR APPLICATION — 5 to 20 gals. per acre. GROUND APPLICATION — 15 to 200 gals. per acre. ORCHARD APPLICATION — 70 to 600 gals. per acre. (Citrus may require up to 100 gals. 170 to 100 gals. per acre rates are considered concentrate rates and require special concentrate equipment.)

ALMONDS: Brown Rot, Shot-hole Fungus. 4 to 5 lbs. plus 1/2 to 1 gal. Best Dormant Flowable Oil per 100 gals. water. For Brown Rot Control apply in popcorn stage. Apply in petal fall or jacket stage for shot-hole fungus.

APRICOTS: Brown Rot Fungus, Shot-hole Fungus. 4 to 5 lbs. plus 1/2 to 1 gal. Best Dormant Flowable Oil per 100 gals. water. Apply in red bud to popcorn stage. Brown Rot may also require a jacket spray, especially during wet years.

CHERRIES: Brown Rot, Blossom Rot. 4 to 5 lbs. plus 1/2 to 1 gal. Best Dormant Flowable Oil per 100 gals. water. Apply in popcorn stage.

PEACHES, NECTARINES: Peach Blight (shot-hole), Peach Leaf Curl. 4 to 5 lbs. plus 1/2 to 1 gal. Best Dormant Flowable Oil per 100 gals. water from Nov. to Dec. 15. Brown Rot, Peach Blight (shot-hole) - apply in pink bud stage using 4 to 5 lbs. plus 1/2 to 1 gal. Best Dormant Flowable Oil.

PRUNES, PLUMS: Blossom Rot, Brown Rot, Shot-hole. 4 to 5 lbs. plus 1/2 to 3/4 gal. Best Dormant Flowable Oil. Apply in early popcorn stage.

CUCUMBERS: Downy Mildew, Anthracnose, Angular Leaf Spot. 10 lbs. per acre. Begin application before disease appears and repeat at 7 to 10 day intervals.

ONIONS: Downy Mildew. 10 lbs. per acre. Begin application before disease appears and repeat at 7 to 10 day intervals.

17 :1,68

ACTIVE INGREDIENT:

BY WT.

Copper expressed as Metallic (in Basic Copper Sulfate) - - - - - 53%

INERT INGREDIENTS - - - - - 47%

TOTAL 100%

MIXING DIRECTIONS

With tank 1/4 full of water and with agitator running, add the amount of Copper required. When tank is two-thirds full, add the oil spreader and fill tank. Keep agitator running during filling and spraying operations.

BEST BASIC COPPER "53" is compatible with spray oils and can be included with higher percentages of Dormant Spray Oils.

CAUTION

KEEP OUT OF REACH OF CHILDREN

HARMFUL IF TAKEN INTERNALLY

AVOID INHALING DUST OR SPRAY MIST

DO NOT APPLY ON TENDER FOLIAGE OF PEACHES, ALMONDS, APRICOTS AND JAPANESE PLUMS