

You Get Results With

**BEST**<sup>®</sup>

ACCEPTED

6-12-72

# MALATHION\* 5 DUST

Read This Label Carefully

## CAUTION

**KEEP OUT OF REACH OF CHILDREN**

**Harmful If Swallowed**

HARMFUL IF SWALLOWED!

AVOID BREATHING OF DUST

AVOID CONTACT WITH SKIN

WASH THOROUGHLY AFTER USING

AVOID CONTAMINATION OF FEED AND FOODSTUFFS

Do not allow this product to drift onto adjoining food or feed crops as residues may contaminate these crops making them unfit for sale, feed or human consumption. Keep clear of streams, fish ponds and wildlife areas

Highly toxic to fish. Do not contaminate any body of water, by direct application, cleaning of equipment or disposal of wastes and containers. Do not apply this product to any crop not specifically indicated on this label

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service

**DESTROY BAG IMMEDIATELY WHEN EMPTY**

**BURY IN SAFE PLACE, OR BURN; STAY OUT OF SMOKE**

**POULTRY** *Chicken Red Mite, Lice, Northern Fowl Mite* Make a thorough application to birds. For dust box treatment per 100 birds use 10 to 12 lbs. per 6 cu. ft. of box. Poultry house control apply 2 to 5 lbs. per 100 sq. ft. on litter, floor and walls

(28 4 72)

## DIRECTIONS FOR USE - Continued

**PEACHES** *Aphid* Apply 50 to 60 lbs. per acre. Do not apply within 7 days of harvest

**PEAS** *Aphid, Pea Weevil* Apply 20 to 25 lbs. per acre. Do not apply within 3 days of harvest or 7 days of harvest if forage is to be fed to livestock

**POTATOES** *Aphid, Leafhopper, False Chinch Bug, Mealy Bug* Apply 30 to 60 lbs. per acre. No time limitations

**SOYBEANS** *Mexican Bean Beetle* Apply 30 to 40 lbs. per acre. No time limitation

**STRAWBERRIES** *Strawberry Leafroller, Spittle Bug, Aphid, Lygus Bug, Leafhopper, Spider Mite, Strawberry Root Weevil, Field Crickets* Apply 20 to 40 lbs. per acre. Do not apply within 3 days of harvest

**TOMATOES** *Aphid, Spider Mite, Tomato Russet Mite* Apply 30 to 40 lbs. per acre. Do not apply within 1 day of harvest

**WALNUTS** *Walnut Husk Fly, Aphid* Apply 50 to 60 lbs. per acre. No time limitations

## ACTIVE INGREDIENTS:

Malathion\* ..... 5%

## INERT INGREDIENTS:

TOTAL ..... 100%

\* O,O-Dimethyl dithiophosphate of diethyl mercaptosuccinate

E. P. A. Reg. No. 7001-82-AA

## DIRECTIONS FOR USE

**ALFALFA, CLOVER** *Spotted Alfalfa Aphid, Pea Aphid, Alfalfa Weevil Larvae, Lygus Bug, Leafhopper, Spittle Bug, Vetch Bruchid, Armyworm, Spider Mites* Apply 30 to 50 lbs. per acre. Do not apply within 7 days of harvest. Do not apply during full bloom

**ASPARAGUS** *Asparagus Beetle, Thrips* Apply 20 to 25 lbs. per acre. Do not apply within 1 day of harvest

**BEANS** *Aphids, Lygus Bug, Grasshoppers, Leafhoppers, Thrips, Mexican Bean Beetle* Apply 20 to 35 lbs. per acre. Do not apply within 7 days of harvest

**BEETS** (Sugar) *Aphids, Leafhopper, Spider Mites* Apply 35 to 40 lbs. per acre. Do not apply within 3 days of harvest or within 7 days of harvest if tops are to be fed to livestock

**BEETS** (Table) *Aphids, Leafhopper, Spider Mites* Apply 35 to 50 lbs. per acre. Do not apply within 7 days of harvest

**BLACKBERRIES, BOYSENBERRIES, LOGANBERRIES, RASPBERRIES** *Aphids, Spider Mites, Thrips, Leafhoppers, Rose Scale* Apply 35 to 50 lbs. per acre. Do not apply within 1 day of harvest

**BROCCOLI, BRUSSELS SPROUTS, CABBAGE, CAULIFLOWER** *Aphid, Leafhopper, Lygus Bug, Flea Beetle, Imported Cabbage Worm, Cabbage Looper* Apply 35 to 50 lbs. per acre. Do not apply within 7 days of harvest

on Brussels Sprouts, Cabbage and Cauliflower. Do not apply within 3 days of harvest on Broccoli

**CARROTS** *Leafhopper, Aphid* Apply 30 to 40 lbs. per acre. Do not apply within 7 days of harvest

**CELERY** *Aphid, Leafhopper, Spider Mite* Apply 25 to 30 lbs. per acre. Do not apply within 7 days of harvest

**CHERRIES** *Black Cherry Aphid, Leafroller* Apply 35 to 40 lbs. per acre. Do not apply within 3 days of harvest. May injure foliage on some varieties

**CORN** *Sap Beetle, Corn Earworm, Thrips* Apply 20 to 30 lbs. per acre. Do not apply within 5 days of harvest

**COTTON** *Lygus Bug, Spider Mite, Thrips, Aphid, Fleahopper, Armyworm, Cotton Leaf Perforator* Apply 35 to 65 lbs. per acre. No time limitations

**CUCUMBERS, MELONS, SQUASH, PUMPKINS** *Aphid, Spider Mite, Pickleworm, Cucumber Beetle, Leafhoppers, Squash Vine Borers* Apply 30 to 40 lbs. per acre. Do not apply within 1 day on Cucumbers, Melons, and Squash. Do not apply within 3 days of harvest on Pumpkins

**EGGPLANT, PEPPERS** *Leafhopper, Spider Mite, Aphid, Pepper Maggot* Apply 30 lbs. per acre. Do not apply within 3 days of harvest

**GRAPES** *Leafhopper, Thrips, Spider Mite* Apply 30 to 55 lbs. per acre. Do not apply within 3 days of harvest

**LETTUCE, SPINACH** *Lygus Bug, Aphid, Leafhopper, Cabbage Looper, Spider Mite* Apply 35 to 40 lbs. per acre. Do not apply within 7 days of harvest on head lettuce and spinach. Do not apply within 14 days of harvest on leaf lettuce

**ONIONS** *Thrips, Onion Maggot* Apply 30 to 40 lbs. per acre. Do not apply within 3 days of harvest on Green Onions

**OCCIDENTAL CHEMICAL COMPANY**

P. O. BOX 198 LATHROP, CALIFORNIA 95330 PHONE 425-8811

ACCEPTED WITH RESERVATIONS