


PHOSDRIN[®] 4 EC

ORGANIC PHOSPHATE COMPOUND MIXTURES, LIQUID N.O.S.



ACTIVE INGREDIENTS:

2-Carbomethoxy-1-Methylvinyl dimethyl phosphate, Alpha isomer-----	28.3%
Related Compounds-----	18.9%
Xylene-----	47.6%

INERT INGREDIENTS

5.2%
TOTAL 100.0%

E.P.A. Reg. No. 7001-100-AA

Contains 4 pounds of Phosdrin[®] insecticide per gal

DIRECTIONS FOR DILUTION

DILUTE APPLICATION

Ground Application Apply specified rate in 20 to 60 gallons of water per acre
Orchard Application Apply specified rate in 100 to 800 gallons of water per acre

CONCENTRATE APPLICATION

Ground Application Apply specified rate in not less than 5 gallons of water per acre
Orchard Application Apply specified rate in 20 to 100 gallons of water per acre
Special concentrate equipment is necessary for these uses

AIR APPLICATION

Apply specified rate in 5 to 20 gallons of water per acre. Orchard rates should not be applied in less than 10 gallons of water per acre

DIRECTIONS FOR USE

ALFALEA CLOVER Spotted Alfalfa Aphid, Alfalfa Caterpillar, Leafhopper, Grasshopper, Cutworm (climbing), Lygus Bug, Red Spider Mite, Alfalfa Weevil Larvae 1/2 to 1 pt per acre. Do not apply within 1 day of harvest

ARTICHOKES Aphid, Plume Moth 2 pts per acre. Do not apply within 2 days of harvest

BEANS PEAS TOMATOES Aphid, Red Spider Mite, Lygus Bug, Grasshopper, Leafhopper, Cabbage Looper 1/2 to 1 pt per acre. Do not apply within 1 day of harvest

BROCCOLI BRUSSELS SPROUTS CABBAGE CAULIFLOWER CELERY Aphid, Cabbage Looper, Adult Dipterous Leafminer, Imported Cabbage Worm, Cutworm (climbing), Leafhopper, Lygus Bug, False Chinch Bug, Grasshopper, Red Spider Mite, Salt Marsh Caterpillar, Corn Earworm 1 pt to 1 qt per acre. Do not apply within 3 days of harvest on broccoli brussels sprouts, cabbage, 5 days on celery

LETTUCE SPINACH Aphid, Leafhopper, Adult Dipterous Leafminer, Cabbage Looper, Lygus Bug, Corn Earworm, Imported Cabbage Worm, Cutworm (climbing), Grasshopper, Salt Marsh Caterpillar 1 qt per acre. Do not apply within 4 days of harvest on lettuce, 7 days on spinach

ONIONS (including green onions) Thrips, Cutworm (climbing), Aphid 1/2 to 1 pt per acre. Do not apply within 1 day of harvest

BY WT.

DIRECTIONS FOR USE — Continued

CUCUMBERS CANTALOUPE HONEYDEW MUSKMELONS SQUASH (Summer), **WATERMELONS** Aphid, 1/4 to 1/2 pt per acre Cabbage Looper, Leafhopper, Dipterous Leafminer (Adults), Lygus Bug, Red Spider Mite, False Chinch Bug, Cutworm (Climbing), Grasshopper, Salt Marsh Caterpillar 1/2 to 1 pt per acre. Do not apply within 1 day of harvest

TURNIPS BEETS (including tops), **CARROTS EGGPLANT PEPPERS** Aphid 1/4 to 1/2 pt per acre Leafhopper, Dipterous Leafminer (adults), Grasshopper, Lygus Bug, False Chinch Bug, Imported Cabbage Worm, Corn Earworm, Armyworm, Red Spider Mite, Salt Marsh Caterpillar 1/2 to 1 pt per acre. Do not apply within 3 days of harvest on Beets or Turnips, or within 1 day of harvest on Carrots, Eggplant and Peppers

CORN Aphid, Red Spider Mite, Corn Earworm, Cutworm (climbing) 1/2 to 1 pt per acre. Do not apply within 1 day of harvest

SORGHUM (forage & grain) Aphids 1/4 to 1/2 pt per acre Corn Earworm, Webworm, Fall Armyworm 1/2 to 1 pt per acre. Do not apply within 3 days of harvest

POTATOES Aphid, Red Spider Mites, Lygus Bugs, Leaf Hopper, Cabbage Looper, Apply 1/2 pint per acre. Do not apply within 4 days of harvest


APPLES CHERRIES PEACHES PEARS PLUMS Aphid, Red Spider Mite, Lygus Bug, Grasshopper, Red-banded Leafroller, 1 to 2 qts per acre. Do not apply to Cherries within 2 days of harvest. Do not apply to Apples, Peaches, Pears and Plums within 1 day of harvest

GRAPES Aphid 1/2 to 1 pt per acre Leaf Folder, Leafhopper, Red Spider Mite, Lygus Bug, Redbanded Leafroller 1 to 2 pts per acre. Do not apply within 2 days of harvest

CITRUS (Oranges, Lemons, Grapefruit) Fruit Tree Leafroller, Orange Tortrix, Omnivorous Leafroller 2 pts per acre Western Tussock Moth Larvae, Citrus Cutworm, Variegated Cutworm, Pink Scavenger Caterpillar 4 pints per acre Aphids 1 to 2 pts per acre. Allow at least 7 days between applications. Do not apply within 1 day of harvest

STRAWBERRIES Aphid, Red Spider Mite, Thrips, Strawberry Leafroller, Lygus Bug, Grasshopper, Salt Marsh Caterpillar 1/2 to 2 pt per acre. Do not apply within 1 day of harvest

WALNUTS Aphid, 2 to 3 pts per acre Red Spider Mite, Orange Tortrix, Omnivorous Looper, Fruit Tree Leafroller, Western Tussock Moth 3 to 5 pts per acre. Do not apply within 1 day of harvest



Antidotes: CALL

NOTE TO PHYSICIAN: Atropine is an antidote for 2-PAM. 2-PAM is also antidotal and may be used.

INTERNAL: If swallowed give a tablespoon of water. Have victim lie down and keep eyes open. Have victim breathe fresh air.
EYES: Flush immediately with water for 15 minutes.
SKIN: In case of contact wash immediately with soap and water.

DANGER KEI

POISONOUS OR ABSORBED (Rapidly) DO NOT

Do not get in the eyes, on skin or on clothing. In case of contact wash immediately with water. For more information contact the U.S. Bureau of Mines for PHOSDRIN products. Do not apply in areas where there may be danger of drift and volatile residues have dissipated. Do not eat or drink, or smoke, or use tobacco, or eat or drink, or use tobacco, or eat or drink, or use tobacco, or eat or drink, or use tobacco. Wash all contaminated clothing before reuse. Do not allow this product to drift onto adjacent property. Do not make them unfit for sale, feed or human consumption. This product is toxic to fish, birds, and other animals. Do not apply where runoff is likely to occur. Do not apply where it will contaminate water by cleaning of equipment. Do not allow this product to drift onto adjacent property. Because of the danger of killing bees, do not apply in areas where bees are present. Do not apply in areas where bees are present. Do not apply in areas where bees are present. Avoid spillage while mixing and filling. Do not use for any further purpose. Discard immediately. Do not use, pour, spill, or store near heat or fire.

OCCIDENTAL CHEMICAL COMPANY
A DIVISION OF OCCIDENTAL PETROLEUM CORPORATION
P. O. Box 198 • LATHROP, CALIFORNIA 95330 • Phone (209) 858-2511


CONDITIONS OF SALE: 1. Seller warrants that the product is as represented and is reasonably fit for use as directed and authorized to make any other warranty, guarantee, or refund of purchase price.

PHOSDRIN[®] 4 EC

ORGANIC PHOSPHATE COMPOUND MIXTURES, LIQUID N.O.S.

BY WT.

Alpha isomer	28.3%
	18.9%
	47.6%
	5.2%
TOTAL	100.0%



DIRECTIONS FOR USE — Continued

CUCUMBERS, CANTALOUPE, HONEYDEW, MUSKMELONS, SQUASH (Summer), **WATERMELONS** Aphid. $\frac{1}{4}$ to $\frac{1}{2}$ pt per acre **Cabbage Looper, Leafhopper, Dipterous Leafminer (Adults), Lygus Bug, Red Spider Mite, False Chinch Bug, Cutworm (Climbing), Grasshopper, Salt Marsh Caterpillar** $\frac{1}{2}$ to 1 pt per acre Do not apply within 1 day of harvest

TURNIPS, BEETS (including tops), **CARROTS, EGGPLANT, PEPPERS** Aphid $\frac{1}{4}$ to $\frac{1}{2}$ pt per acre **Leafhopper, Dipterous Leafminer (adults), Grasshopper, Lygus Bug, False Chinch Bug, Imported Cabbage Worm, Corn Earworm, Armyworm, Red Spider Mite, Salt Marsh Caterpillar** $\frac{1}{2}$ to 1 pt per acre Do not apply within 3 days of harvest on Beets or Turnips, or within 1 day of harvest on Carrots, Eggplant, and Peppers

CORN Aphid, Red Spider Mite, Corn Earworm, Cutworm (climbing) $\frac{1}{2}$ to 1 pt per acre Do not apply within 1 day of harvest

SORGHUM (forage & grain) Aphids $\frac{1}{4}$ to $\frac{1}{2}$ pt per acre **Corn Earworm, Webworm, Fall Armyworm** $\frac{1}{2}$ to 1 pt per acre Do not apply within 3 days of harvest

POTATOES Aphid, Red Spider Mites, Lygus Bugs, Leaf Hopper, Cabbage Looper, Apply $\frac{1}{2}$ pint per acre Do not apply within 4 days of harvest

APPLES, CHERRIES, PEACHES, PEARS, PLUMS Aphid, Red Spider Mite, Lygus Bug, Grasshopper, Red-banded Leafroller. 1 to 2 qts per acre Do not apply to Cherries within 2 days of harvest Do not apply to Apples, Peaches, Pears and Plums within 1 day of harvest

GRAPES Aphid $\frac{1}{2}$ to 1 pt per acre **Leaf Folder, Leafhopper, Red Spider Mite, Lygus Bug, Redbanded Leafroller** 1 to 2 pts per acre Do not apply within 2 days of harvest

CITRUS (Oranges, Lemons, Grapefruit) **Fruit Tree Leafroller, Orange Tortrix, Omnivorous Leafroller** 2 pts per acre **Western Tussock Moth Larvae, Citrus Cutworm, Variegated Cutworm, Pink Scavenger Caterpillar** 4 pints per acre **Aphids** 1 to 2 pts per acre Allow at least 7 days between applications Do not apply within 1 day of harvest

STRAWBERRIES Aphid, Red Spider Mite, Thrips, Strawberry Leafroller, Lygus Bug, Grasshopper, Salt Marsh Caterpillar $\frac{1}{2}$ to 2 pt per acre Do not apply within 1 day of harvest

WALNUTS Aphid, 2 to 3 pts per acre **Red Spider Mite, Orange Tortrix, Omnivorous Looper, Fruit Tree Leafroller, Western Tussock Moth** 3 to 5 pts per acre Do not apply within 1 day of harvest



Antidotes: CALL A PHYSICIAN IMMEDIATELY!

NOTE TO PHYSICIAN: Atropine is antidotal.

2-PAM is also antidotal and may be administered in conjunction with atropine.

INTERNAL: If swallowed give a tablespoonful of salt in a glass of warm water. Repeat until vomit fluid is clear. Have victim lie down and keep quiet.

EYES: Flush immediately with water for at least 15 minutes.

SKIN: In case of contact wash immediately with soap and water.

DANGER KEEP OUT OF REACH OF CHILDREN

**POISONOUS IF SWALLOWED, INHALED
OR ABSORBED THROUGH SKIN OR EYES**

(Rapidly Absorbed Through Skin)

DO NOT BREATHE VAPORS!

Do not get in the eyes, on skin or on clothing. Wear natural rubber gloves, protective clothing and goggles. In case of contact wash immediately with soap and water. Wear a mask or respirator of a type passed by the U.S. Bureau of Mines for PHOSDRIN protection. Keep all unprotected persons out of operating areas or vicinity where there may be danger of drift. Vacated areas should not be re-entered until drifting insecticide and volatile residues have dissipated. Wash hands, arms and face thoroughly with soap and water before eating or smoking. Wash all contaminated clothing with soap and hot water before reuse.

Do not allow this product to drift onto adjoining food or feed crops as residues may contaminate these crops making them unfit for sale, feed or human consumption. Keep clear of food or dairy animals or animals being finished for slaughter.

This product is toxic to fish, birds, and other wildlife. Keep out of any body of water. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

Because of the danger of killing bees, do not apply to legumes while in bloom. If infestation requires immediate attention to save the crop, spray in early morning or late evening while bees are inactive.

Avoid spillage while mixing and filling spray equipment. When container is empty, immediately wash thoroughly with water and detergent and destroy by punching drum full of holes or crushing so that it cannot be used for any further purpose. Discard in a safe place, never reuse container.

Do not use, pour, spill, or store near heat or open flame.

(39 3 72)

OCCIDENTAL CHEMICAL COMPANY

A DIVISION OF OCCIDENTAL PETROLEUM CORPORATION

P. O. Box 198 • LATHROP, CALIFORNIA 95330 • Phone (209) 858-2511

CONDITIONS OF SALE: 1. Seller warrants that this product consists of the ingredients specified and is reasonably fit for use as directed on the label. No one, other than an officer of Seller, is authorized to make any other warranty, guarantee or direction concerning this product. 2. Because the time, place, rate of application and other conditions of use are beyond Seller's control, Seller's liability from the handling, storage and use of this product is limited to replacement of product or refund of purchase price.