

(67)

ACCEPTED



3-19-73

FUNGICIDE AND RODENTICIDE ACT.
FOR E.C. WHICH IS REGISTERED UNDER NO. 7001-104

You Get Results With
BEST

PHOSPHAMIDON 8 EC

READ THIS LABEL CAREFULLY

	ACTIVE INGREDIENTS	BY WT.		
	*Phosphamidon	75.5%		78%
	Related Compounds	2.5%		
POISON	INERT INGREDIENTS	TOTAL	100%	

Contains 8 lbs Phosphamidon Insecticide per gallon
* 2-Chloro 2-diethylcarbamoyl 1-methylvinyl dimethyl phosphate U.S. Pat. No. 2,908,605
Phosphamidon 8 EC is a liquid systemic insecticide for the control of both sucking and chewing insects. It may be applied as a diluted spray or as a concentrate spray.

DANGER! KEEP OUT OF REACH OF CHILDREN
SEE WARNING AND ANTIDOTE STATEMENTS
ON BACK PANEL

(230:4/72) rjw

NET CONTENTS 1 GALLON
MANUFACTURED FOR:

OCCIDENTAL CHEMICAL COMPANY

A DIVISION OF OCCIDENTAL PETROLEUM CORPORATION

P.O. BOX 198 LATHROP, CALIFORNIA 95330 PHONE (209) 858-2511

2001-104



P O I S O N



3-19-73

7001-104

ANTIDOTE: If swallowed Give a tablespoonful of salt in a glass of warm water and repeat until vomit fluid is clear. Have victim lie down and keep quiet. **CALL A PHYSICIAN IMMEDIATELY!** **Note to Physician:** Atropine is Antidote. 2-PAM is also antidotal and may be administered in conjunction with atropine. If on Skin Remove contaminated clothing, immediately wash skin with soap and water.

DANGER: POISONOUS IF SWALLOWED, INHALED OR ABSORBED THROUGH SKIN - RAPIDLY ABSORBED THROUGH SKIN!

DO NOT INHALE DO NOT GET ON SKIN DO NOT TAKE INTERNALLY

Wear a mask or respirator of a type passed by the U.S. Bureau of Mines for phosphamidon protection. Do not get eyes, on skin or on clothing. Wear natural rubber gloves, protective clothing and goggles. In case of contact, wash immediately with soap and water. Wash hands, arms and face thoroughly with soap and water before eating or drinking. Wash all contaminated clothing with soap and hot water before reuse. Keep all unprotected persons out of operating areas or vicinity where there may be danger of drift. Vacated areas should not be re-entered until drifting insecticide and volatile residues have dissipated. Not for use or storage in or around the home. Do not store or transport near food or food. **DO NOT USE POUR SPILL OR STORE NEAR HEAT OR OPEN FLAME.** Do not use on Maple Trees, Peaches, Plums or Sweet Cherries as injury may occur, avoid drift to these varieties. Rinse equipment and containers and dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed of by puncturing holes in them and burying with wastes. Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.

This product is toxic to wildlife. Birds feeding in treated areas may be killed. Do not apply where birds are nesting. Do not contaminate water by cleaning of equipment or disposal of wastes.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective dusts are not obtained from your Agricultural Extension Service.

When container is empty immediately wash thoroughly and destroy. Never reuse. Store in a cool, dry place.

E.P.A. Reg. No. 7001-104-AA

DIRECTIONS FOR USE

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH CAUTIONS, WARNINGS AND DIRECTIONS AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

KEEP PRODUCT IN ORIGINAL CONTAINER. DO NOT PUT CONCENTRATE OR DILUTE INTO FOOD OR DRINK CONTAINERS.

SEE DIRECTIONS FOR AMOUNT OF PRODUCT IN THE FOLLOWING GALLONS WATER PER ACRE UNLESS OTHER WISE NOTED.

Ground Application - Vegetable & Field Crops - 30 to 250 gals. **Deciduous Fruit Crops** - 60 to 800 gals. **Citrus** - 400 to 3000 gals. On young fruit trees use minimum of 35 gals. water per acre. Where Recommended for

Air Application - Field Crops - 2 to 10 gals. **Orchard Crops** - 10 to 20 gals.

APPLES: For control of Damage from *Green and Rosy Apply Aphids, European Red Mite, San Jose Scale, Fruit Tree Leaf Roller. Apply 10 to 20 gals. per acre. Super 94 Spray Oil in Western States per 100 gals. water. For control of *Apple Scab* apply 5 to 10 gals. per acre. For control of *Apple Scab* apply 5 to 10 gals. per acre. Pre-blossom and Post-blossom for control of *Apple Scab*, *First Broad Leafing Moth, Leathoppers, Green and Rosy Apply Aphids, Leaf Miners.* Apply 10 to 20 gals. per acre. Do not use Phosphamidon EC on apples within 30 days of harvest.*

BROCCOLI, CAULIFLOWER, APRICOTS, IMPORTED CABBAGEWORM: Apply 10 to 20 gals. per acre in sufficient water to cover one acre. Apply 10 to 20 gals. per acre for control of *Imported Cabbageworm.* Do not apply within 3 days of harvest.

CANTALOUPE, WATERMELON, APHIDS, LEATHOPPER, LEATHOPPER: Apply 10 to 20 gals. per acre in sufficient water to cover. Do not apply within 3 days of harvest.

COTTON: For control of *Leaf Miner, Aphids, Thrips, Fleahoppers (except Texas & Oklahoma), Spider Mites.* Apply 2 to 3 gals. per acre with ground equipment. Spray directly on young plants in sufficient water to cover. Make final application at the second true leaf stage. Mid and Late Season. *Lygus, Flea Beetles, Leathoppers.*

DANGER: POISONOUS IF SWALLOWED, INHALED OR ABSORBED THROUGH SKIN - RAPIDLY ABSORBED THROUGH SKIN!

DO NOT INHALE DO NOT GET ON SKIN DO NOT TAKE INTERNALLY

Wear a mask or respirator of a type passed by the U.S. Bureau of Mines for phosphorimidon protection. Do not get eyes on skin or on clothing. Wear natural rubber gloves, protective clothing and goggles. In case of contact, wash immediately with soap and water. Wash hands, arms and face thoroughly with soap and water before eating or drinking. Wash all contaminated clothing with soap and hot water before reuse. Keep all unprotected persons out of operation areas or vicinity where there may be danger of drift. Vacated areas should not be re-entered until drifting insecticide and volatile residues have dissipated. Not for use or storage in or around the home. Do not store or transport near food or food. **DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME.** Do not use on Maple, Elm, Peach, Plum or Sweet Cherries as injury may occur, avoid drift to these varieties. Rinse equipment and containers and dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed of by puncturing them and burying with wastes. Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.

This product is toxic to wildlife. Birds feeding in treated areas may be killed. Do not apply where runoff is likely. Do not contaminate water by cleaning of equipment or disposal of wastes.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information is obtained from your Agricultural Extension Service.

When container is empty immediately wash thoroughly and destroy. Never reuse. Store in a cool, dry place.

E.P.A. Reg. No. 7001-104-AA

DIRECTIONS FOR USE

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH CAUTIONS, WARNINGS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

KEEP PESTICIDE IN ORIGINAL CONTAINER. DO NOT PUT CONCENTRATE OR DILUTE INTO FOOD OR DRINK CONTAINERS.

USE DESIGNATED AMOUNT OF PRODUCT IN THE FOLLOWING GALLONS WATER PER ACRE UNLESS OTHERWISE NOTED.

Ground Application - Vegetable & Field Crops - 30 to 250 gals. **Deciduous Fruit Crops** - 60 to 800 gals. **Citrus** - 400 to 3,000 gals. On young Fruit Plantings - Use minimum of 15 gals. water per acre. Where Recommended for

Air Application - Field Crops - 3 to 10 gals. **Orchard Crops** - 10 to 20 gals.

APPLES - Early, Delayed Dormant Onset - *Green and Rosy Apply Aphids, European Red Mite, San Jose Scale, Fruit Tree Leaf Roller*. Apply 1/2 pt. plus 2 gals. Super 94 (Spray Oil) or 1/2 gal. Super 94 (Spray Oil) in Western States per 100 gals. water in full coverage sprays. Do not combine of sprays with Sulfur, Captain or Phaltan. Pre-blossom and Post-blossom. First brood cover sprays. *First Brood Codling Moth, Leafhoppers, Green and Rosy Apply Aphids, Leaf Miners*.

Apply 1/2 pt. plus 200 gals. water. Do not use Phosphorimidon 8 EC on apples within 30 days of harvest.

BROCCOLI - CABBAGEFLOWER - *Aphids, Imported Cabbageworm*. Apply 1/2 to 1 pt. in sufficient water to cover one acre. Apply every 10 to 14 days in a preventative program for Aphid control. Do not apply within 3 days of harvest.

CANTALOUPE - WATERMELON - *Aphids, Leafminer, Leafhopper*. Apply 1/2 to 1 pt. per acre in sufficient water to cover. Do not apply within 1 day of harvest.

CITRUS - Early Season - 2 through 10 gal. type. *Aphids, Thrips, Fleahoppers (except Texas & Oklahoma), Spider Mites*. Apply 1/2 to 1 gal. per acre. Do not use in combination with other insecticides. Spray directly on young plants in sufficient water to cover. Make first application at the seedling stage. Mid and Late Season - *Lygus, Flea Beetles, Leafhoppers*.

Apply 1/2 to 1 gal. per acre in sufficient water to cover by ground or air. Do not apply within 2 weeks of harvest.

CUCUMBERS - *Aphids, Leaf Miner, Leafhopper*. Apply 1/2 to 1 pt. per 100 gals. water per acre. Do not apply within 3 days of harvest.

CUCUMBERS - *Aphids, Leaf Miner, Leafhopper*. Apply 1/2 to 1 pt. per 100 gals. water per acre. Do not apply within 3 days of harvest.

CUCUMBERS - *Aphids, Leaf Miner, Leafhopper*. Apply 1/2 to 1 pt. per 100 gals. water per acre. Do not apply within 3 days of harvest.

CUCUMBERS - *Aphids, Leaf Miner, Leafhopper*. Apply 1/2 to 1 pt. per 100 gals. water per acre. Do not apply within 3 days of harvest.

PEPPERS - *Aphids, Leaf Miner, White Fly*. Apply 1/2 to 1 pt. per acre in sufficient water to cover. Do not apply within 3 days of harvest.

POTATOES - *Aphids, Colorado Potato Beetles, Leaf Miners, White Fly, Flea Beetles, Stinkbugs*. Apply 1/2 to 1 pt. per acre in sufficient water to cover. Do not apply within 3 days of harvest.

POTATOES - *Aphids, Colorado Potato Beetles, Leaf Miners, White Fly, Flea Beetles, Stinkbugs*. Apply 1/2 to 1 pt. per acre in sufficient water to cover. Do not apply within 3 days of harvest.

POTATOES - *Aphids, Colorado Potato Beetles, Leaf Miners, White Fly, Flea Beetles, Stinkbugs*. Apply 1/2 to 1 pt. per acre in sufficient water to cover. Do not apply within 3 days of harvest.

POTATOES - *Aphids, Colorado Potato Beetles, Leaf Miners, White Fly, Flea Beetles, Stinkbugs*. Apply 1/2 to 1 pt. per acre in sufficient water to cover. Do not apply within 3 days of harvest.

POTATOES - *Aphids, Colorado Potato Beetles, Leaf Miners, White Fly, Flea Beetles, Stinkbugs*. Apply 1/2 to 1 pt. per acre in sufficient water to cover. Do not apply within 3 days of harvest.

POTATOES - *Aphids, Colorado Potato Beetles, Leaf Miners, White Fly, Flea Beetles, Stinkbugs*. Apply 1/2 to 1 pt. per acre in sufficient water to cover. Do not apply within 3 days of harvest.



DIRECTIONS FOR USE

ALFALFA CLOVER Aphid, armyworm, grasshoppers, clover leaf weevil, leafhopper, spider mites - 1 1/4 to 2 pts Rates of 1 1/4 to 1 1/2 pts No time limitation Rates over 1 1/2 pts to 2 pts do not apply within 7 days of harvest
ALMONDS Aphid, peach twig borer, red spider mites - 2 1/2 to 8 pts No time limitation
APPLES, PEARS Aphid, mealybugs, mites, pear psylla - 2 to 15 pts Codling moth, fruit tree leaf roller, plum curculio, red-banded leaf roller, San Jose scale - 4 to 15 pts Dormant and delayed dormant sprays - aphid, mites, red-banded leaf roller - 3 to 15 pts plus 6 to 8 gals Super 94 (spray oil) For concentrate sprays and delayed dormant use the lower rate of Super 94 (spray oil) Caution May cause injury to McIntosh and Cortland varieties Do not apply during bloom period and within 3 days of harvest on apples Do not apply within 1 day of harvest on pears
APRICOTS Aphid, codling moth, orange tortrix, soft brown scale, terrapin scale - 4 to 10 pts Do not apply within 7 days of harvest Full coverage spray
ASPARAGUS Asparagus beetle, thrips - 1 1/4 pt Do not apply within 1 day of harvest
AVOCADOS Green house thrips, Latania scale, omniverous looper, soft brown scale, orange tortrix - 4 to 9 pts Do not apply within 7 days of harvest
BARLEY OATS RYE WHEAT Aphid, cereal leaf beetle, grasshoppers, greenbugs - 1 1/4 pt Do not apply within 7 days of harvest
BEANS Aphid, cucumber beetle, Japanese beetle, leafhopper, lygus bug, Mexican bean beetle, mites - 1 1/4 to 1 1/2 pt Do not apply within 1 day of harvest
BEETS, DANDELIONS Aphid - 1 to 2 1/2 pts Do not apply within 7 days of harvest
BLACKBERRIES BOYSENBERRIES DEWBERRIES LOGANBERRIES RASPBERRIES Japanese beetle, leafhoppers, mites, thrips - 1 to 4 pts Aphid, rose scale - 2 to 4 pts Do not apply within 1 day of harvest
BLUEBERRIES Cherry fruit worm, cranberry fruit worm, Japanese beetle, plum curculio, sharp nose leaf hopper - 1 1/2 to 2 1/2 pts Do not apply within 1 day of harvest
BROCCOLI BRUSSELS SPROUTS MUSTARD GREENS TURNIPS CABBAGE CAULIFLOWER COLLARDS RALE Aphid, cabbage looper, diamond back moth, imported cabbage worm - 1 1/2 to 2 1/2 pt Do not apply within 7 days of harvest except on broccoli within 3 days of harvest
CARROTS Aphid - 1 to 2 pts Leafhopper - 1 1/2 to 2 pts Do not apply within 7 days of harvest
CELERY Aphid, spider mites - 1 to 1 1/2 pts Do not apply within 7 days of harvest
CHERRIES Black cherry aphid, bud moth, cherry fruit fly, torbes scale, fruit tree leaf roller, San Jose scale - 2 1/2 to 8 pts Lesser peach tree borer - 4 to 8 pts Do not apply within 3 days of harvest May injure foliage of some varieties
CHESTNUTS Mites - 2 to 5 pts No time limitation
CITRUS (Grapefruit lemons limes oranges tangerines tangelos kumquats) Aphid, black scale, California red scale, citricola scale, Florida red and purple scale, soft scale, thrips, yellow scale - 7 to 25 pts Mediterranean fruit fly - 1 to 8 pts Do not apply within 7 days of harvest Do not apply during full bloom
CORN Aphid, cereal leaf beetle, corn earworm - apply to silks as soon as they appear Make 3 to 4 applications at 2-3 day intervals as infestations warrant reapply immediately after rain corn rootworm - adult grasshoppers, sap beetle, thrips - 1 pt Do not apply within 5 days of harvest For grain or forage use
COTTON Aphid, boll weevil, cotton leaf perforator, cotton leafworm, leafhoppers, lygus bug, mites, thrips, white flies, fall armyworm, fleahopper, garden webworm, grasshopper - 1 1/2 to 4 pts per acre No time limitation
CRANBERRIES Black headed fireworm, cranberry fruitworm, leafhoppers, spittlebug nymphs - 1 to 2 pts Do not apply within 3 days of harvest
CUCUMBERS MELONS PUMPKINS SQUASH Aphid, pickleworm, spider mites, thrips - 1 to 1 1/2 pts Cucumber beetle - 1 1/4 pt Do not apply within 3 days of harvest on pumpkins 1 day of harvest on cucumbers, melons and squash Do not apply unless plants are dry
CURRANTS GOOSEBERRIES Mites Japanese beetle - 1 to 2 pts Currant aphid, imported currantworm - 2 pts Do not apply within 3 days of harvest on Gooseberries 1 day of harvest on currants
DATES Nitidulid beetle - 2 pts Do not apply within 7 days of harvest
EGGPLANT Aphid, spider mite - 1 1/4 to 3 1/2 pts Lace bug - 2 to 3 1/2 pts Do not apply within 3 days of harvest

CYTHION* 8 AQUAMUL

Active Ingredients:

Malathion (O,O-Dimethyl dithiophosphate of diethyl mercaptosuccinate) 81.8

Inert Ingredients:

18.2

TOTAL 100.0%

Contains 8 lbs Malathion per gallon

* Reg Trademark of American Cyanamid Company

EPA Reg. No. 7001-114-AA

DIRECTION FOR USE

Do not use this product for any uses other than those specified on this label All rates are for a dilute spray All rates are per acre.

DILUTION RATE

Ground Application Apply specified rate in 20 to 60 gallons of water per acre Orchard Application Apply specified rate in 100 to 800 gallons of water per acre

CONCENTRATE APPLICATION

Ground Application Apply specified rate in not less than 5 gallons of water per acre Orchard Application Apply specified rate in 20 to 100 gallons of water per acre Special concentrate equipment is necessary for these uses Air Application Apply specified rate in 5 to 20 gallons of water per acre Orchard rates should not be applied in less than 10 gallons of water per acre

CAUTION

KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed Avoid breathing of spray mist Avoid contact with skin wash thoroughly after using Avoid contamination of feed and foodstuff Highly toxic to fish Do not contaminate any body of water by direct application, cleaning of equipment or disposal of wastes and containers Highly toxic to bees exposed to direct treatment or residues on crops Protective information may be obtained from your cooperative Agricultural Extension Service When container is empty immediately wash thoroughly with water and detergent and destroy in a safe place Never reuse container


WARNING - FLAMMABLE Do not use, pour, spill, or store near heat or open flame OCCIDENTAL CHEMICAL COMPANY

A DIVISION OF OCCIDENTAL PETROLEUM CORPORATION P O Box 198 • LATHROP, CALIFORNIA 95330 • Phone (209) 858-2811

CONDITIONS OF SALE: Seller warrants that this product consists of the ingredients specified on this label and that it is to be used as directed on the label. No other than the use of Seller is authorized to make any other warranty, guarantee or direction concerning this product. Because the time, place, rate of application and other conditions of use are beyond Seller's control, Seller's liability from the handling, storage and use of this product is limited to replacement of product or refund of purchase price.

DIRECTIONS FOR

ENDIVE Aphid, mites - 1 to 2 pts Do not apply within 3 days of harvest
FIGS Dried fruit beetle, vinegar flies - 2 1/2 pts Do not apply within 3 days of harvest
FILBERT Apple mealybug, eye-spotted bud moth, No time limitation
GRAPES Drosophila, Japanese beetle, leafhopper Do not apply within 3 days of harvest Caution May cause injury to fruit
GRASS GRASS HAY Aphid, armyworm, grasshopper
HOPS Aphid - 1 1/2 to 1 3/4 pt Do not apply within 7 days of harvest
HORSE RADISH KOHLRABI SALSIFY, SHALLOT 1 to 2 pts Do not apply within 7 days of harvest
LENTILS Aphid - 1 pt Do not apply within 3 days of harvest
LETTUCE Aphid, leafhopper - 1 1/4 to 2 pts Cabbage root fly - 1 1/2 to 2 1/2 pts Do not apply within 7 days of harvest on head lettuce
MACADAMIA NUTS Green stink bug - 3 to 15 pts
MANGO PASSION FRUIT GUAVA Fruit flies - 1 to 2 pts Do not apply within 2 days of harvest
OKRA Aphid, Japanese beetle - 1 1/2 pt Do not apply within 7 days of harvest
ONIONS GARLIC Thrips - 1 to 2 pts Onion maggot and green onions
PARSLEY PARSNIPS RADISHES RUTABAGAS Do not apply within 7 days of harvest on parsley, radishes, rutabagas
PEACHES NECTARINES Aphid (green, peach, borer), mites (European red, two spotted) - 2 1/2 to 9 pts Do not apply within 7 days of harvest on peach, nectarines, oriental fruit moth, terrapin scale - 5 to 9 pts
PEACHES (dormant and delayed dormant) Aphid - 1 to 2 pts Super 94 (spray oil) Use lower rate of Super 94 (spray oil)
PEANUTS Leafhoppers, thrips - 1 pt No time limitation
PEAS Aphid, pea weevil - 1 to 2 1/2 pt Do not apply within 7 days of harvest
PECANS Aphid, mites, pecan bud moth, pecan leaf miner - 2 1/2 to 3 1/2 pts No time limitation
PEPPERS Aphid - 1 to 2 1/2 pts Maggots - 1 pt
PEPPERMINT, SPEARMINT Adult flea beetle - up to 7 days of harvest
PINEAPPLE Mealybug - 1 pt Do not apply within 7 days of harvest
PLUMS PRUNES Lesser peach tree borer, mealybug - 1 to 10 pts Dormant and delayed dormant aphid - 5 to 10 pts Use lower rate of Super 94 (spray oil) for dormant and delayed dormant aphid - 5 to 10 pts Do not apply within 3 days of harvest Full coverage spray
POTATOES Aphid, false chinch bug, leafhopper, mite, Colorado potato beetle - 1 to 2 pts Do not apply within 7 days of harvest
RICE Rice leaf miner, rice stink bug - 1 pt Do not apply within 7 days of harvest
SAFFLOWER Aphid, grasshopper, lygus bug - 1 to 2 pts Do not apply within 7 days of harvest
SOYBEANS Grasshoppers, Japanese beetle, Mexican bean beetle, aphid, field crickets, lygus bug, leafhopper, strawberry root weevil, thrips, white fly, strawberry root weevil, apply to soil surface before planting
SUGAR BEETS Aphid, mites - 1 to 2 pts Grasshopper - 1 to 2 pts Do not apply within 7 days of harvest
SWEET POTATOES Leafhoppers - 1 to 1 1/2 pts 1 day of harvest
TOMATOES Aphid, spider mite - 1 to 2 pts Drosophila, fruit worms - 3 to 5 pts Do not apply within 7 days of harvest
WALNUTS Aphid, mites, walnut husk fly - 4 to 12



CYTHION* 8 AQUAMUL

Active Ingredients:

Malathion (0,0-Dimethyl dithiophosphate of diethyl mercaptosuccinate) 81.8

Inert Ingredients:

18.2

TOTAL 100.0%

Contains 8 lbs. Malathion per gallon

* Reg. Trademark of American Cyanamid Company

EPA Reg. No. 7001-114-AA

DIRECTION FOR USE

Do not use this product for any uses other than those specified on this label. All rates are for a dilute spray. **All rates are per acre.**

DILUTION RATE

Ground Application - Apply specified rate in 20 to 60 gallons of water per acre

Orchard Application - Apply specified rate in 100 to 800 gallons of water per acre

CONCENTRATE APPLICATION

Ground Application - Apply specified rate in not less than 5 gallons of water per acre

Orchard Application - Apply specified rate in 20 to 100 gallons of water per acre

Special concentrate equipment is necessary for these uses

Air Application - Apply specified rate in 5 to 20 gallons of water per acre. Orchard rates should not be applied in less than 10 gallons of water per acre.

CAUTION

KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed. Avoid breathing of spray mist. Avoid contact with skin. Wash thoroughly after using. Avoid contamination of feed and foodstuff.

Highly toxic to fish. Do not contaminate any body of water by direct application. Cleaning of equipment or disposal of wastes and containers.

Highly toxic to bees exposed to direct treatment. For details on traps. Protective information may be obtained from your cooperative Agricultural Extension Service.

When container is empty, immediately wash thoroughly with water and detergent and destroy. Discard in a safe place. Never reuse container.

WARNING - FLAMMABLE! Do not use, pour, spill, or store near heat or open flame.

OCCIDENTAL CHEMICAL COMPANY

A DIVISION OF OCCIDENTAL PETROLEUM CORPORATION
P.O. Box 198 • LATHROP, CALIFORNIA 95330 • Phone (209) 858 2511

WARRANTY: Occidental Chemical Company warrants that this product consists of the ingredients specified on this label and is fit for use as described on the label. No other party, including Occidental Chemical Company, is authorized to make any other warranty, guarantee or representation concerning this product. Because the time, place, rate of application, and other conditions of use are beyond Occidental Chemical Company's control, Occidental Chemical Company's liability from the handling, storage and use of this product is limited to replacement of product or refund of purchase price.

DIRECTIONS FOR USE (continued)

ENDIVE *Aphid, mites* - 1 to 2 pts. Do not apply within 7 days of harvest

FIGS *Dried fruit beetle, vinegar flies* - 2 1/2 pts. plus 2 gals. unsulfurized molasses as a bait spray. Do not apply within 3 days of harvest

FILBERT *Apple mealybug, eye-spotted bud moth, filbert aphid, scale (crawler stage), tingids* - 2 1/2 to 3 pts. No time limitation

GRAPES *Drosophila, Japanese beetle, leathopper, mealybugs, spider mite, terrapin scale* - 2 to 2 1/4 pts. Do not apply within 3 days of harvest. Caution: May cause injury to foliage on some varieties

GRASS GRASS HAY *Aphid, armyworm, grasshopper, leathopper* - 1 1/4 pt. No time limitation

HOPS *Aphid* - 1/2 to 1 1/4 pt. Do not apply within 7 days of harvest

HORSERADISH KOHLRABI SALSIFY, SHALLOTS, SPINACH, SWISS CHARD WATERCRESS *Aphid,* 1 to 2 pts. Do not apply within 7 days of harvest

LENTILS *Aphid* - 1 pt. Do not apply within 3 days of harvest

LETTUCE *Aphid, leathopper* - 1 1/4 to 2 pts. *Cabbage looper, mites* - 2 pts. Do not apply within 14 days of harvest on leaf lettuce. 7 days of harvest on head lettuce

MACADAMIA NUTS *Green stink bug* - 3 to 15 pts. No time limitation

MANGO, PASSION FRUIT, GUAVA *Fruit flies* - 1/4 pt. plus 1 lb. partially hydrolyzed yeast protein or enzymatic yeast hydrolyzate. Do not apply within 2 days of harvest

OKRA *Aphid, Japanese beetle* - 1 1/2 pt. Do not apply within 1 day of harvest

ONIONS, GARLIC *Thrips* - 1 to 2 pts. *Onion maggot* - 2 pts. Do not apply within 3 days of harvest on garlic and green onions

PARSLEY PARSNIPS, RADISHES, RUTABAGAS *Aphid* - 1 to 2 pts. Do not apply within 21 days of harvest on parsley. 7 days of harvest on parsnips. radishes. 3 days of harvest on rutabagas

PEACHES, NECTARINES *Aphid (green peach, black cherry, black peach, rusty plum), Japanese beetle, mites (European red, two spotted)* - 2 1/2 to 9 pts. *Cottony peach scale, lesser peach tree borer, plum curculio, oriental fruit moth, terrapin scale* - 5 to 9 pts. Do not apply within 7 days of harvest. Full coverage spray

PEACHES (Dormant and delayed dormant) *Aphid, peach twig borer, scale (San Jose)* - 4 to 9 pts. plus 6.8 gals. Super 94 (spray oil). Use lower rate of Super 94 (spray oil) at delayed dormant and concentrate spray

PEANUTS *Leathoppers, thrips* - 1 pt. No time limitation

PEAS *Aphid, pea weevil* - 1 to 2 1/2 pt. Do not apply within 3 days of harvest

PECANS *Aphid, mites, pecan bud moth, pecan leaf casebearer, pecan nut casebearer, pecan phylloxera* - 2 1/2 to 12 1/2 pts. No time limitation

PEPPERS *Aphid* - 1/4 to 1 1/2 pts. *Maggots* - 1 1/2 pts. Do not apply within 3 days of harvest

PEPPERMINT, SPEARMINT *Adult flea beetle, aphid, leathoppers, spider mite* - 1 pt. Do not apply within 7 days of harvest

PINEAPPLE *Mealybug* - 5 pts. Do not apply within 7 days of harvest

PLUMS, PRUNES *Lesser peach tree borer, mealy plum aphid, European red, two spotted mite* - 5 to 10 pts. Dormant and delayed dormant aphid. San Jose scale - 5 to 10 pts. plus 8 gals. Super 94 (spray oil). Use lower rate of Super 94 (spray oil) for concentrate sprays. Dormant. Do not apply within 3 days of harvest. Full coverage spray

POTATOES *Aphid, false chinch bug, leathopper, mealybug* - 1 to 3 pts. No time limitation

RICE *Rice leaf miner, rice stink bug* - 1 1/2 pt. Do not apply within 7 days of harvest

SAFFLOWER *Aphid, grasshopper, lygus bug* - 1 to 1 1/4 pt. Do not apply within 3 days of harvest

SOYBEANS *Grasshoppers, Japanese beetle, Mexican bean beetle* - 2 pts. No time limitation

STRAWBERRIES *Aphid, field crickets, lygus bug, potato leathopper, spider mite, spittlebug, strawberry leafroller, strawberry root weevil, thrips, white flies* - 1 to 2 pts. Do not apply within 3 days of harvest. For strawberry root weevil apply to soil surface before planting and work into top 6 to 8 inches

SUGAR BEETS *Aphid, mites* - 1 to 2 pts. *Grasshoppers* - 2 pts. Do not apply within 3 days of harvest

SWEET POTATOES *Leathoppers* - 1 to 1 1/4 pts. *Morning Glory leafminer* - 1 1/2 pt. Do not apply within 3 days of harvest

TOMATOES *Aphid, spider mite* - 1 to 2 pts. *Drosophila tomato russet mite* - 1 1/2 to 3 1/2 pts. *Armyworm, fruit worms* - 3 1/2 pts. Do not apply within 1 day of harvest at the 1 to 2 pt. rate. Rates over 2 pts. to 3 1/2 pts. do not apply within 5 days of harvest

WALNUTS *Aphid, mites, walnut husk fly* - 4 to 12 1/2 pts. No time limitation

