

DIRECTIONS FOR USE

For Codling Moth, Apple Maggot, Oriental Fruit Moth, Plum Curculio, etc.—Apply Sevin 4 Flowable at Petal Fall and every 14 days thereafter until control is achieved.

Before using agitate contents by shaking or rolling container.

Directions for Dilution

DILUTE APPLICATION. Ground Application: Apply specified rate to 40 gallons of water per acre. **Orchard Application:** Apply specified rate to 100 to 150 gallons of water per acre.

CONCENTRATE APPLICATION: Ground Application: Apply specified rate to at least 15 gallons of water per acre. **Orchard Application:** Apply specified rate to 100 to 150 gallons of water per acre. Special concentrate dilution is necessary for these uses.

AIR APPLICATION: Apply specified rate to 100 gallons of water per acre. Concentrate should be diluted to 100 to 150 gallons of water per acre.

Fruit and Nut Crops:

APPLES, PEARS: Codling Moth, Apple Sucker, Woolly Apple Aphid, Green Apple Aphid, Pear Psylla, Orange Tortrix, Scale, (San Jose, Oyster Shell, Lecanium), Lygus Bug, Apple Rust Mite, Pear Leaf Blister Mite, Pear Rust Mite. Apply 3 to 12 quarts per acre. Do not apply within 14 days of harvest. To avoid apple thinning delay use until at least 30 days after bloom.

ALMONDS: Peach Twig Borer, San Jose Scale, Fruit Tree Leafroller. Apply 3 to 8 quarts per acre.

APRICOTS, NECTARINES, PEACHES: Oriental Fruit Moth, Peach Twig Borer, Plum Curculio, Codling Moth, Orange Tortrix, Tussock Moth, Fruit Tree Leafroller, June Beetles, Red-banded Leafroller, Japanese Beetle, European Earwig, Scale (Lecanium, San Jose, Olive). Apply 3 to 8 quarts per acre. Do not apply within 14 days of harvest. Do not apply within 14 days of harvest.

BANANAS: Leaf Feeding Lepidoptera Larvae (Seranicida, Opsiphanes, Caligo and Sibine) Species. Apply 3 to 12 quarts per acre.

BLUEBERRIES: Blueberry Maggot, Cherry Fruitworm, Cranberry Fruitworm, European Fruit Lecanium, Japanese Beetle.

CANE BERRIES (Blackberry, Boysenberry, Dewberry, Loganberry, Raspberry): Japanese Beetle, Leafroller, Leafhopper, Raspberry Aphid. Apply 3 to 8 quarts per acre.

CHERRIES, PLUMS, PRUNES: Cherry Fruit Fly, Cherry Fruitworm, Fruit Tree Leafroller, Peach Twig Borer, Lower Peach Tree Borer, Black Cherry Aphid, Mealy Plum Aphid, Plum Curculio, Prune Leafhopper, Red-banded Leafroller, Japanese Beetle, Scale (Brown, Soft, Forays, San Jose, Oyster Shell, Lecanium), Codling Moth, Eastern Tent Caterpillar, Orange Tortrix, Tussock Moth. Apply 3 to 8 quarts per acre.

CITRUS: California Orange Dog, Citrus Cutworm, Fruit Tree Leafroller, Orange Tortrix, Western Tussock Moth, Scale (Black, Brown, Soft, California Red, Citricola). Apply 3 to 8 quarts per acre.

CRANBERRIES: Fruitworms, Fireworms, Leafhopper, Cutworm, Japanese Beetle.

FILBERTS: Filbert Moth, Filbert Aphid, Filbert Leafroller. Apply 3 to 8 quarts per acre. No time limitations.

GRAPES: Grape Leafhopper, Leafhopper, Grape Leaf Skeletonizer, European Fruit Lecanium. Apply 1 1/2 to 3 quarts per acre. **Grape Berry Moth, Japanese Beetle, June Beetle, Red-banded Leafroller.** Apply 3 to 8 quarts per acre. No time limitation.

OLIVES: Olive Scale. Apply 3 to 8 quarts per acre with 100 to 150 gallons of water per acre. Do not apply more than 2 applications per season.

STRAWBERRIES: Meadow Spittlebug, Leafroller, Strawberry Weevil. Apply 3 to 8 quarts per acre. Do not apply within one day of harvest.

WALNUTS: Codling Moth, Scale (Frosted, European Fruit Lecanium, Calico), Filbert Worm, Fruit Tree Leafroller. Apply 3 quarts per acre. **European Earwig.** Apply 3 to 8 quarts per acre. No time limitation.

Forage, Feed, Seed, Vegetable Crops:

ALFALFA, CLOVER, PEANUTS, SOYBEANS, COWPEAS: Blister Beetle, Mexican Bean Beetle, Thrips, Leafhopper, Three-cornered Alfalfa Hopper, Cucumber Beetles, Bean Leaf Beetle, Japanese Beetle, Alfalfa Caterpillar, Velvet Bean Caterpillar, Alfalfa Weevil Larvae, Armyworms, Webworms, Corn Earworm, Stink Bugs, Grasshoppers. Apply 1 to 1 1/2 quarts per acre. No time limitation.

ASPARAGUS: Asparagus Beetle. Apply 1 to 2 quarts per acre during growing season. Do not repeat application within 3 days. Do not apply within 14 days of harvest. For control of asparagus beetle and other pests with later Asparagus Beetle and Apache Cicada. Apply 2 to 4 quarts per acre.

BEANS: Mexican Bean Beetle, Leafhoppers, Bean Leaf Beetle, Cucumber Beetles, Western Bean Cutworm, Flea Beetles, Japanese Beetles, Velvet Bean Caterpillar, Cutworm. Apply 3 to 8 quarts per acre. **Lima Bean Pod Borer, Corn Earworm, Lygus Bug, Stink Bug.** Apply 3 to 8 quarts per acre. No time limitation.

BROCCOLI, BRUSSELS SPROUTS, CABBAGE, CAULIFLOWER, KOHLRABI: Flea Beetle, Harlequin Bug, Corn Earworm, Armyworm, Cabbage Looper. Apply 1 to 2 quarts per acre. Do not apply more than 2 quarts per acre. When the sprouts appear, the sprouts should be sprayed with 3 to 4 quarts per acre. Do not apply more than 2 quarts per acre. Do not apply more than 1 day intervals.

CORN: Corn Earworm, Corn Rootworm, Adult European Corn Borer, Fall Armyworm, Soy Beetle, Japanese Beetle, Flea Beetle, Leafhoppers. Apply 3 to 8 quarts per acre. No time limitation. **Corn Earworms** apply 3 to 8 quarts per acre. **European Corn Borer** apply 3 to 8 quarts per acre. **Fall Armyworm** apply 3 to 8 quarts per acre. **Soy Beetle** apply 3 to 8 quarts per acre. **Japanese Beetle** apply 3 to 8 quarts per acre. **Leafhoppers** apply 3 to 8 quarts per acre. **European Corn Borer** apply 3 to 8 quarts per acre. **Fall Armyworm** apply 3 to 8 quarts per acre. **Soy Beetle** apply 3 to 8 quarts per acre. **Japanese Beetle** apply 3 to 8 quarts per acre. **Leafhoppers** apply 3 to 8 quarts per acre.

COTTON: Thrips, Flea Beetles, Cotton Fleahopper, Cotton Leafworm, Striped Blister Beetle. Apply 3 to 8 quarts per acre. **Bollweevil, Bollworm, Fall Armyworm, Leaf Perforator, Leafrollers, Leafhoppers, Tarnished Plant Bug.** Apply 3 to 8 quarts per acre. **Pink Bollworm, Salt Marsh Caterpillar, Stink Bug.** Apply 3 to 8 quarts per acre.

CUCUMBERS, PUMPKINS, SQUASH, MELONS: Pickworm, Melonworm, Cucumber Beetle, Flea Beetle, Leafhopper, Squash Bug. Apply 3 to 8 quarts per acre.

EGGPLANTS, PEPPERS, TOMATOES: Flea Beetle, Leafhopper, Tomato Fruit Worm, Fall Armyworm, Tomato Hornworm, European Corn Borer, Stink Bug, Tarnished Plant Bug. Apply 3 to 8 quarts per acre. **Cutworms.** Apply 3 to 8 quarts per acre.

GRASS (Forage, Feed): No time limitations.

OKRA: Corn Earworm, Leafhopper. Apply 3 to 8 quarts per acre. No time limitation.

POTATOES: Colorado Plant Bugs. Apply 3 to 8 quarts per acre.

RICE: Armyworms, Leafhoppers. Apply 3 to 8 quarts per acre. No time limitation.

SORGHUM: Armyworms, Leafhoppers. Apply 3 to 8 quarts per acre. No time limitation.

SUGAR BEETS: Leafhoppers. Apply 3 to 8 quarts per acre. No time limitation.

TOBACCO: Hornworm, Leafhopper. Apply 3 to 8 quarts per acre. No time limitation.

CAUTION

Do not swallow. Do not get in eyes. Do not get on clothing. Do not get on skin. Do not get on face. Do not get on hands. Do not get on feet. Do not get on shoes. Do not get on socks. Do not get on underwear. Do not get on pajamas. Do not get on nightgown. Do not get on robe. Do not get on slippers. Do not get on shoes. Do not get on socks. Do not get on underwear. Do not get on pajamas. Do not get on nightgown. Do not get on robe. Do not get on slippers.

Note for Physician: This product is a potent insecticide and should be used with caution. It is highly toxic to fish and aquatic life. It is also highly toxic to birds and mammals. It is also highly toxic to bees and other beneficial insects. It is also highly toxic to cats and dogs. It is also highly toxic to children. It is also highly toxic to pregnant women. It is also highly toxic to nursing infants. It is also highly toxic to the elderly. It is also highly toxic to the infirm. It is also highly toxic to the handicapped. It is also highly toxic to the mentally ill. It is also highly toxic to the emotionally unstable. It is also highly toxic to the socially maladjusted. It is also highly toxic to the chronically ill. It is also highly toxic to the acutely ill. It is also highly toxic to the dying. It is also highly toxic to the dead.

Do not use in open areas. Do not use near water. Do not use near food. Do not use near children. Do not use near pets. Do not use near livestock. Do not use near wildlife. Do not use near birds. Do not use near mammals. Do not use near reptiles. Do not use near amphibians. Do not use near fish. Do not use near aquatic life. Do not use near bees. Do not use near other beneficial insects. Do not use near cats. Do not use near dogs. Do not use near children. Do not use near pregnant women. Do not use near nursing infants. Do not use near the elderly. Do not use near the infirm. Do not use near the handicapped. Do not use near the mentally ill. Do not use near the emotionally unstable. Do not use near the socially maladjusted. Do not use near the chronically ill. Do not use near the acutely ill. Do not use near the dying. Do not use near the dead.

Do not use in open areas. Do not use near water. Do not use near food. Do not use near children. Do not use near pets. Do not use near livestock. Do not use near wildlife. Do not use near birds. Do not use near mammals. Do not use near reptiles. Do not use near amphibians. Do not use near fish. Do not use near aquatic life. Do not use near bees. Do not use near other beneficial insects. Do not use near cats. Do not use near dogs. Do not use near children. Do not use near pregnant women. Do not use near nursing infants. Do not use near the elderly. Do not use near the infirm. Do not use near the handicapped. Do not use near the mentally ill. Do not use near the emotionally unstable. Do not use near the socially maladjusted. Do not use near the chronically ill. Do not use near the acutely ill. Do not use near the dying. Do not use near the dead.

Do not use in open areas. Do not use near water. Do not use near food. Do not use near children. Do not use near pets. Do not use near livestock. Do not use near wildlife. Do not use near birds. Do not use near mammals. Do not use near reptiles. Do not use near amphibians. Do not use near fish. Do not use near aquatic life. Do not use near bees. Do not use near other beneficial insects. Do not use near cats. Do not use near dogs. Do not use near children. Do not use near pregnant women. Do not use near nursing infants. Do not use near the elderly. Do not use near the infirm. Do not use near the handicapped. Do not use near the mentally ill. Do not use near the emotionally unstable. Do not use near the socially maladjusted. Do not use near the chronically ill. Do not use near the acutely ill. Do not use near the dying. Do not use near the dead.

GRASS (Forage, Pasture): Thrips, Armyworms, Grasshoppers. Apply 1 to 1 1/2 qts. per acre. No time limitations.

OKRA: Corn Earworm, Stink Bugs. Apply 1 to 2 qts. per acre. No time limitations.

PEAS: Leafhoppers, Colorado Potato Beetle, Armyworm, Grasshoppers. Apply 1 to 2 qts. per acre. No time limitations.

POTATOES: Colorado Potatoes and Flea Beetle, Leafhoppers, Fall Armyworm, Stink Bugs, Tarnished Plant Bugs. Apply 1 to 2 qts. per acre. Foliage applications only. No time limitations.

RICE: Armyworms, Stink Bugs, Grasshoppers, Tadpole Shrimp. Apply 1 to 2 qts. per acre. Do not use within 15 days of proper herbicide. Do not apply within 14 days of harvest. Highly toxic to shrimp and crabs. Do not apply where this is an important resource.

SORGHUM: Armyworms, Webworms, Corn Earworm, Stink Bug, Grasshoppers. Apply 1 to 2 qts. per acre. Sorghum Midge, Cutworms. Apply 2 qts. per acre. Do not apply within 21 days for grain. No time limitation for forage.

SUGAR BEETS: Leafhoppers, Armyworms, Webworms, Grasshoppers. Apply 1 to 2 qts. per acre. Do not apply within 14 days of harvest.

TOBACCO: Hornworm, Budworms, Flea Beetles, June Beetles, Japanese Beetles. Apply 1 to 2 qts. per acre for leaf treatment. Avoid excessive applications. Late season applications are necessary to control late season pests after primary control has been achieved.

CAUTION KEEP OUT OF REACH OF CHILDREN

Wear protective clothing. Avoid breathing in spray mist. Avoid contact with skin. Wash immediately and thoroughly after handling. Wear clean clothing. Avoid storage near food and food products. Wear goggles and gloves if needed.

Note for Physician: Carbaryl is a moderate to severe plechogenic agent. Atropine is a specific antidote. This is a cholinergic agent. It has been reported to cause treatment of cholinergic crisis. Patients in contact with this product should be treated by the Agricultural Extension Service. Do not use when rain is expected or when wind is blowing. Do not apply to water. Do not use in areas where water is used for drinking. Do not use in areas where water is used for irrigation. Do not use in areas where water is used for domestic purposes.

Do not use in areas where water is used for drinking. Do not use in areas where water is used for irrigation. Do not use in areas where water is used for domestic purposes.

Do not use in areas where water is used for drinking. Do not use in areas where water is used for irrigation. Do not use in areas where water is used for domestic purposes.

Do not use in areas where water is used for drinking. Do not use in areas where water is used for irrigation. Do not use in areas where water is used for domestic purposes.

Do not use in areas where water is used for drinking. Do not use in areas where water is used for irrigation. Do not use in areas where water is used for domestic purposes.

Do not use in areas where water is used for drinking. Do not use in areas where water is used for irrigation. Do not use in areas where water is used for domestic purposes.

Do not use in areas where water is used for drinking. Do not use in areas where water is used for irrigation. Do not use in areas where water is used for domestic purposes.

Do not use in areas where water is used for drinking. Do not use in areas where water is used for irrigation. Do not use in areas where water is used for domestic purposes.

Do not use in areas where water is used for drinking. Do not use in areas where water is used for irrigation. Do not use in areas where water is used for domestic purposes.

Do not use in areas where water is used for drinking. Do not use in areas where water is used for irrigation. Do not use in areas where water is used for domestic purposes.

Do not use in areas where water is used for drinking. Do not use in areas where water is used for irrigation. Do not use in areas where water is used for domestic purposes.

Do not use in areas where water is used for drinking. Do not use in areas where water is used for irrigation. Do not use in areas where water is used for domestic purposes.

Do not use in areas where water is used for drinking. Do not use in areas where water is used for irrigation. Do not use in areas where water is used for domestic purposes.

Do not use in areas where water is used for drinking. Do not use in areas where water is used for irrigation. Do not use in areas where water is used for domestic purposes.

You Get Results With



SEVIN* 4 FLOWABLE

ACTIVE INGREDIENT:	BY WT
Carbaryl (1-Naphthyl Methylcarbamate)	- - 42.8%
INERT INGREDIENTS	- - - - - 57.2%
	TOTAL 100.0%

Contains 4 lbs. Carbaryl Per Gallon U.S. Patent No. 2,903,478
Trade Mark of Union Carbide Corp. for the active ingredient Carbaryl insecticide

EPA Reg. No. 7001-105-AA

241-1-73 (17)

CAUTION: KEEP OUT OF REACH OF CHILDREN SEE ADDITIONAL CAUTIONS ON SIDE PANEL

NET CONTENTS 1 GALLON

OCCIDENTAL CHEMICAL COMPANY

A DIVISION OF OCCIDENTAL PETROLEUM CORPORATION
P. O. Box 198 • LATHROP, CALIFORNIA 95330 • Phone (209) 858-2511