

Store in Cool, Dry Place

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH CAUTIONS, WARNINGS AND DIRECTIONS, AND IN CONFORMITY WITH FEDERAL AND STATE REGULATIONS.

KEEP PESTICIDE IN ORIGINAL CONTAINER. DO NOT PUT CONCENTRATE OR DILUTE INTO FOOD OR DRINK CONTAINERS.

USE DESIGNATED AMOUNT OF PRODUCT IN THE FOLLOWING GALLONS WATER PER ACRE UNLESS OTHERWISE NOTED.

GROUND APPLICATION—Vegetable and Field Crops—30 to 250 gals. Cotton—3 to 20 gals. Deciduous Fruit Crops—40 to 800 gals. Citrus—400 to 3,000 gals. (On Young Fruit Plantings—Use a minimum of 15 gals. water per acre).

Where Recommended for **AIR APPLICATION**—Field Crops—3 to 10 gals.

Begin application at first sign of insects. Repeat as necessary. Do not apply within 4 days of harvest unless otherwise noted. Do not use with highly alkaline materials such as lime or bordeaux.

MIXING DIRECTIONS: Add to spray tank when 1/4 to 1/2 full with agitator running and maintain agitation until spraying is completed.

BEANS (Dry or Field), SOYBEANS, LIMA BEANS AND PEAS for Processing: Alfalfa Looper, Aphids, Spider Mites, Lygus, Leaf Miner; **GREEN BEANS:** Aphids, Spider Mites, Loopers, Nitidulids, Leafhoppers—1 pt. per acre in water to cover. 100 gals. per acre is preferable. Large Loopers, Stink Bugs—1 1/2 pts. per acre in water to cover.

BROCCOLI, CABBAGE, CAULIFLOWER, BRUSSELS SPROUTS, KALE, COLLARDS: Imported Cabbageworm, Diamond-back Caterpillars, Aphids—1 pt. per acre. Looper—2 pts. per acre. Do not apply within 1 day of harvest, except do not apply to kale and collards within 4 days of harvest.

CELERY: Aphids, Loopers, Armyworms, Salt Marsh Caterpillars, Leaf Miners—Use ground application only with fine mist spray—1 pt. in 50 to 75 gals. water per acre on plants less than 1/2 grown. On larger plants use 1 1/2 pts. in 100 to 150 gals. water per acre.

COTTON: Lygus—1/2 to 1 pt. per acre by air or ground. Leaf Perforator, Fleahoppers, Armyworms, Bollworms, Loopers and for Suppression of Mites—1 pt. per acre by air or ground. Apply at 7 to 10 day intervals while pests persist. For Heavy Population of Listed Insects—Use 1/2 pt. per acre by air or ground in control programs with other insecticides registered for use on cotton.

CUCUMBERS, SUMMER SQUASH: Cucumber Beetles, Aphid, Spider Mites, Loopers, Thrips, Armyworms—1 pt. per acre in water to cover. Pickleworms, Leaf Miners—1/2 pt. per acre in water to cover. By Air—Use minimum of 10 gals. water per acre. May be used up to harvest.

EGGPLANT, PEPPERS: Aphids, Spider Mites, Blister Beetles, Flea Beetles, Leaf Miners, Whitefly—1 pt. per acre. Do not apply within 1 day of harvest.

WALNUTS: Aphids—1/2 to 1 pt. in 100 gals. water. Do not use over 10 pts. per acre. Do not apply within 10 days of harvest.

GRAPES: Drosophila, Leafhoppers—1/2 to 3/4 pt. per 100 gals. water. (100 to 130 gals. diluted spray per acre depending on site of vines.) Use only 1/2 pt. per 100 gals. water on Muscat varieties. Do not use on Italian varieties as fruit injury may occur.

HOPS: Aphids, Spider Mites, Corn Earworm, Hop Looper, Cutworm—1 pt. per acre.

SPINACH, CHARD, T. GREENS: Aphids, Loopers, Armyworms, Salt Marsh Caterpillars, Leaf Miners—Use ground application only with fine mist spray—1 pt. in 50 to 75 gals. water per acre on plants less than 1/2 grown. On larger plants use 1 1/2 pts. in 100 to 150 gals. water per acre. Do not apply within 1 day of harvest.

MELONS (Cantaloupes, Honeydew, Muskmelons, Watermelons, Pumpkins, Winter Squash): Loopers, Armyworms, Leafhoppers, Spider Mites, Leaf Miners—1 to 2 pts. per acre (200 gals. diluted spray per acre). Do not apply within 1 day of harvest. Do not apply when temperature is over 90° F.

ORANGES, LEMONS, GRAPEFRUIT, TANGERINES: Aphids, Soft Brown Scale Crawlers, Mites, Citrus Cutworms, Leaf Roller (Tortrix)—1 pt. to 100 gals. water. Thrips (Bloom)—1/2 pt. per 100 gals. water. Use up to 7 days of harvest. (Calif. and Ariz.—Do not apply within 1 day of harvest.)

PEACHES (Western States Only): Lygus, Suppression of Mites, Peach Twig Borer, Oriental Fruit Moth (California Only)—1/2 pt. to 100 gals. water. To avoid fruit spotting, do not use within 30 days of harvest. Do not apply when temperature is over 90° F. Do not apply more than 8 pts. per acre.

RICE: Grasshoppers—1/2 to 3/4 pt. per acre. Apply at first sign of insects. Do not make more than 3 applications. Do not apply within 2 days of harvest.

STRAWBERRIES: Leaf Roller, Spider Mites, Aphids, Omnivorous Leaf Tier, Spittlebugs (Northwest)—1 pt. per acre. Do not apply within 1 day of harvest.

SUGAR BEETS: Spider Mites, Leafhoppers—1 pt. per acre by air or ground. By Air—Use in 1 to 5 gals. water per acre. Do not apply within 5 days of harvest.

TOBACCO: Hornworm—1 pt. to 100 gals. water using up to 200 gals. diluted spray per acre. Flea Beetles—1 pt. per acre in water to cover. Apply at first sign of insects. Repeat as necessary.

TOMATOES (Field): Fruit Flies (Drosophila)—1 pt. in at least 40 gals. water per acre with ground sprayer. 7 to 10 gals. by air. Make first application 5 to 7 days before first picking and continue at 5 to 7 day intervals until the day before harvest. (Field and Greenhouse); Tomato Fruit Worm, Hornworm, Leaf Miners, Flea Beetles, Mites—1 pt. in 100 gals. water. Spray plants thoroughly. For Field—Use up to 200 gals. diluted spray per acre. Do not apply when temperature is over 90° F. Do not apply within 1 day of harvest.

GREENHOUSE VAPOR TREATMENT OF ROSES; OTHER ORNAMENTAL PLANTS, TOMATOES, CUCUMBERS: Spider Mites, Adult Whiteflies, Aphids, Leafrollers, Mealybugs—1 fluid ounce per 10,000 cu. ft. when plants are dry. Apply undiluted on heat pipes or in open pans or hot plates. Apply continuously or to intermittent sections of pipes on each side and on 1 or more of the pipes through center of each range depending upon width. Application to cold pipes should be followed by immediately heating pipes to 160° F. In steam heated houses, close vents for at least 1 hour after pipes are hot. With hot water systems, close house for at least 3 hours. Houses may remain closed overnight following treatment. When application is made to hot pipes, wear goggles and approved respirator. Ventilate house before workers re-enter. Wear respirator when entering treated houses to ventilate. Protect hives of bees in cucumber and other houses by covering with plastic or rubber sheet during treatment and until house is ventilated. Do not apply to tomatoes or cucumbers closer than 1 day before harvest. Spider Mites (Resistant and Non-resistant)—Make 3 to 4 treatments at 3 or 4-day intervals. Repeat as necessary. Adult Whiteflies—Make 2 applications at 7-day intervals. Repeat as necessary. Aphids—Apply when aphids appear. Repeat as necessary. **WARNING:** Avoid overtreatment and direct application to plants as injury may result. White Butterfly and Golden Rapture roses, Pink Champagne Chrysanthemums, Green Wandering Jew, Poinsettias and Dutchman's Pipe may be injured by DIBROM vapor.

FRUIT FLIES (Drosophila sp.) in and around Food Processing Plants, Loading Docks, Cell Piles and Refuse Areas, Cider Mills: Use 2 ounces in 2 1/2 gals. water. 5 pints to 100 gals. Apply as a coarse spray to walls, floors, doorways, windows, refuse and cell piles where insects congregate. Apply every 5 to 7 days as long as necessary to maintain control. Do not apply to cultivated or refuse piles to be fed to livestock. Avoid contamination of feeds, foodstuffs and food processing machinery. Do not apply when plants are in operation or when foods are present or exposed. Do not spray surfaces which will come into contact with food. Cover food containers during spraying periods.

ADULT MOSQUITO: Treat shrubbery, foliage around stagnant pools, marshes, and other areas where water is present. Do not apply to any body of water in which fish are present. **CONTROL—RESIDENTIAL AREAS, CROPS, FEED LOTS:** Consult your State Extension Agent for recommendations and directions. Do not apply to any body of water in which fish are present. See crop recommendations for shrimp. Do not apply to tidal creeks. **CRAFT:** East—Apply 0.8 to 4 lbs. per acre. West—Apply 1 to 2 lbs. per acre. Diluted spray per acre. **MIST OR EQUIPMENT:** Rate of travel and other factors. **CORRALS, HOLDING PENS (Beef and Poultry):** Apply 1 gal. diluted spray per acre. **CLOVER MITES, ROACHES, BARBER:** Apply 1/2 to 1 gal. water. Apply to 1/2 to 1 gal. water. **GRASSHOPPERS:** Field Areas (crops) Grasshoppers—1/2 pt. per acre by air or ground. By Air—Use in 1 to 5 gals. water per acre. Repeat as necessary. Animals may be protected near harvest.

DANGER! Concentrate May Cause Damage. Do Not Use in Field or Goggles. Harmful if Swallowed. Avoid Breathing. Contamination of feed, foodstuffs, open flame. Do not use on Nectariferous plants. May be toxic to bees. Diluted spray. Make new dilution. Avoid Spillage When Empty. Immediately Wash. Do not apply in edible or other areas where food is being raised. Wash areas while food is being raised.

ENVIRONMENTAL CAUTIONS: This or other treated areas may be toxic to fish. Do not apply where runoff is likely from areas treated. Do not contaminate this product only as specified on this label. This product is highly toxic to bees. Formulation may be obtained from your distributor by breaking and burying.

CONDITIONS OF SALE: 1. Seller warrants that the product is as described and is reasonably fit for use as intended. 2. Seller is not authorized to make any other warranty, representation, or statement of liability from the handling, storage, or use of the product. 3. Seller's liability is limited to the refund of purchase price.

7001-108

You Results With

BEST

DIBROM 8 EC

READ THIS LABEL CAREFULLY

Naled Insecticide

	BY WT.
ACTIVE INGREDIENTS: *Naled	58%
Xylene	20%
INERT INGREDIENTS	22%
TOTAL 100%	

Registered Trademark of Chevron Chemical Co. U.S. Patent No. 2,971,882
CONTAINS 8 POUNDS OF TECHNICAL NALED PER GALLON, EQUIVALENT TO 7.2 POUNDS NALED

KEEP OUT OF REACH OF CHILDREN

SEE BACK PANEL FOR ADDITIONAL CAUTIONS

NET CONTENTS 1 GALLON

E.P.A. Reg. No.
7001-108-AA

E.P.A. Est. 7001-CA-1

(224 2/75)

MANUFACTURED BY

OCCIDENTAL CHEMICAL COMPANY

P. O. BOX 198 • LATHROP, CALIFORNIA 95330 • PHONE (209) 858-2511



ADULT MOSQUITO. Treat shrubbery and vegetation where mosquitoes may rest. Shrubby and vegetation around stagnant pools, marshy areas, ponds and shore lines may be treated. Application of this product to any body of water is prohibited. **GNAT, STABLEFLY (Dog Flies), AND HOUSEFLY CONTROL** — RESIDENTIAL AREAS, CROPS LISTED ON LABEL, TIDAL MARSHES, SWAMPS, WOODLAND, **FEED LOTS:** Consult your State Fish and Game Agency before applying this product. It is not necessary to avoid farm buildings and dairy barns. Make applications during peak of infestation and repeat as necessary. See crop recommendations for use limitations near harvest. This rate of application will kill shrimp. Do not apply to tidal or marsh waters which are important shrimp producing areas. **AIR-CRAFT:** East — Apply 0.8 to 4 fluid ozs. per acre diluted with water. Apply 2 to 8 quarts of diluted spray per acre. West — Apply 1.6 to 4 fl. ozs. per acre diluted with water. Apply 2 to 8 quarts of diluted spray per acre. **MIST OR COLD FOG:** Use 3 to 5 qts. per 100 gals. water. Calibrate equipment (rate of travel and output) to apply 0.1 to 0.25 lb. technical Naled per acre.

CORRAL, HOLDING PENS (Dairy and Beef Cattle, Hogs, Sheep, Horses): Houseflies, Mosquitoes — **Area Application** — Use 1.6 to 4 fluid ounces per acre. Dilute 1 to 2 1/2 pts. in 10 gals. water. Apply diluted spray per acre. Apply over areas with animals present.

FLOWERS, MARSHES, SARWIGS AND LEAFHOPPERS Outside Dwellings and in Lawns: Use 1 fl. oz. per 100 sq. ft. Apply to turf and to soil surfaces around flowers, shrubs and trees for general control.

GRASS (TOPPER) — Sold Areas (crops listed on label) — 1/2 to 1/4 pt. per acre by air or ground. Young Grasses — 1 pt. per acre by air or ground. **Armyworms (except rice)** — 1 pt. per acre by air or ground. **Spotted Armyworm** — Use 1 to 5 gals. water per acre. Apply when hoppers hatch. Repeat as necessary. Animals may be present during treatment. See crop recommendations for use limitations.

DANGER! Concentrate May Cause Skin Damage. This Formulation is Corrosive & Causes Eye Damage. Do Not Get on Skin, Eyes or Clothing. Use Waterproof Gloves and Face Shield or Goggles When Handling Concentrate.

Hazards: If Swallowed, Avoid Breathing Spray Mist. In case of contact, immediately remove contaminated clothing and flush skin or eyes with plenty of water; for eyes, get medical attention. Avoid contamination of feed, foodstuffs and drinking water. Do not use, pour, spill or store near heat or open flame. Do not use on Nectarines, Ornamental Cherries, Liquidambar or Chrysanthemums as injury may occur. May be toxic to bees, avoid application during periods of bee activity. Do not store diluted spray. Make new dilution for each use. **WASH EQUIPMENT THOROUGHLY AFTER USE.**

Avoid Spillage When Mixing and Filling Spray Equipment. When Container is Empty, Immediately Wash Thoroughly and Destroy. Never Re-Use.

Do not apply in edible products areas of food processing plants, restaurants or other areas where food is commercially prepared or processed. Do not use in serving areas while food is exposed.

ENVIRONMENTAL CAUTIONS: This product is toxic to fish, shrimp, crab, birds and other wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, streams, ponds, tidal marshes and estuaries. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service. Containers should be disposed of by breaking and burying with wastes.

CONDITIONS OF SALE: 1. Seller warrants that this product consists of the ingredients specified and is reasonably fit for use as directed on the label. No one, other than an officer of Seller, is authorized to make any other warranty, guarantee or direction concerning this product. 2. Because the time, place, rate of application and other conditions of use are beyond Seller's control, Seller's liability from the handling, storage and use of this product is limited to replacement of product or refund of purchase price.

Store in Cool, Dry Place

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH CAUTIONS, WARNINGS AND DIRECTIONS, AND IN CONFORMITY WITH FEDERAL AND STATE REGULATIONS.

KEEP PESTICIDE IN ORIGINAL CONTAINER. DO NOT PUT CONCENTRATE OR DILUTE INTO FOOD OR DRINK CONTAINERS.

USE DESIGNATED AMOUNT OF PRODUCT IN THE FOLLOWING GALLONS WATER PER ACRE UNLESS OTHERWISE NOTED:

GROUND APPLICATION — Vegetable and Field Crops — 30 to 250 gals. Cotton — 3 to 20 gals. Deciduous Fruit Crops — 60 to 800 gals. Citrus — 400 to 3,000 gals. (On Young Fruit Plantings — Use a minimum of 15 gals. water per acre.)

Where Recommended for **AIR APPLICATION** — Field Crops — 3 to 10 gals.

Begin application at first sign of insects. Repeat as necessary. Do not apply within 4 days of harvest unless otherwise noted. Do not use with highly alkaline materials such as lime or bordeaux.

MIXING DIRECTIONS: Add to spray tank when $\frac{1}{4}$ to $\frac{1}{2}$ full with agitator running and maintain agitation until spraying is completed.

BEANS (Dry or Field), SOYBEANS, LIMA BEANS AND PEAS for Processing: Alfalfa Looper, Aphids, Spider Mites, Lygus, Leaf Miner; **GREEN BEANS:** Aphids, Spider Mites, Loopers, Nitidulias, Leafhoppers — 1 pt. per acre in water to cover. 100 gals. per acre is preferable. Large Loopers, Stink Bugs — $\frac{1}{2}$ pts. per acre in water to cover.

BROCCOLI, CABBAGE, CAULIFLOWER, BRUSSELS SPROUTS, KALE, COLLARDS: Imported Cabbage-worm, Diamond-Back Caterpillars, Aphids — 1 pt. per acre. Looper — 2 pts. per acre. Do not apply within 1 day of harvest, except do not apply to kale and collards within 4 days of harvest.

CELERY: Aphids, Loopers, Armyworms, Salt Marsh Caterpillars, Leaf Miners — Use ground application only with fine mist spray — 1 pt. in 50 to 75 gals. water per acre on plants less than $\frac{3}{4}$ grown. On larger plants use $\frac{1}{2}$ pts. in 100 to 150 gals. water per acre.

COTTON: Lygus — $\frac{1}{2}$ to 1 pt. per acre by air or ground. Leaf Perforator, Fleahoppers, Armyworms, Bollworms, Loopers and For Suppression of Mites — 1 pt. per acre by air or ground. Apply at 7 to 10 day intervals while pests persist. For Heavy Population of Listed Insects — Use $\frac{1}{2}$ pt. per acre by air or ground in control programs with other insecticides registered for use on cotton.

CUCUMBERS, SUMMER SQUASH: Cucumber Beetles, Aphids, Spider Mites, Loopers, Thrips, Armyworms — 1 pt. per acre in water to cover. Pickleworms, Leaf Miners — $\frac{1}{2}$ pts. per acre in water to cover. By Air — Use minimum of 10 gals. water per acre. May be used up to harvest.

EGGPLANT, PEPPERS: Aphids, Spider Mites, Blister Beetles, Flea Beetles, Leaf Miners, Whitefly — 1 pt. per acre. Do not apply within 1 day of harvest.

WALNUTS: Aphids — $\frac{1}{2}$ to 1 pt. to 100 gals. water. Do not use over 10 pts. per acre. Do not apply within 10 days of harvest.

GRAPES: Drosophila, Leafhoppers — $\frac{1}{2}$ to $\frac{3}{4}$ pt. per 100 gals. water. 100 to 130 gals. diluted spray per acre depending on size of vines. Use only $\frac{1}{2}$ pt. per 100 gals. water on Muscat varieties. Do not use on Italia varieties as fruit injury may occur.

HOPS: Aphids, Spider Mites, Corn Earworm, Hop Looper, Cutworm — 1 pt. per acre

SPINACH, CHARD: Miners — Use on plants less than 10 days old. Apply within 10 days of harvest.

MELONS (Cantaloupes): Armyworms, Leafhoppers — 1 pt. per acre. Do not apply within 10 days of harvest.

ORANGES, LEMONS: Cutworms, Leafhoppers — 1 pt. per acre. Use up to harvest.

PEACHES (Western): (California Only) — 1 pt. per acre. Do not apply within 10 days of harvest.

RICE: Grasshoppers — 3 applications.

STRAWBERRIES: — 1 pt. per acre.

SUGAR BEETS: — 1 pt. per acre in 100 gals. water per acre.

TOBACCO: Hornworms — 1 pt. per acre.

TOMATOES (Field): (For ground sprayer, apply at 5 to 10 day intervals.)

WORMS, HORNWORMS: For Field — Use up to 100 gals. water per acre. Do not apply at 90° F. Do not apply within 10 days of harvest.

GREENHOUSE, CUMBERS: Spider Mites — 1 pt. per 100 cu. ft. when plants are continuously or intermittently heated.

center of each pipe are hot. immediately heat pipes are hot. overnight follow

respirator. Ventilate. Use sheet during treatment. Do not apply at 3 or 4 day intervals. Repeat.

FRUITING: Avoid Dutchman's Pipe and Golden Raisin. Do not apply within 10 days of harvest.

FRUIT FLIES (House Flies): Refuse Areas, Refuse — Use coarse spray to every 5 to 7 days. Do not apply when plants are in fruit.

Do not apply when plants are in fruit.

Do not apply when plants are in fruit.

Do not apply when plants are in fruit.

7001-108

You Results With

BEST®

DIBROM 8 EC

READ THIS LABEL CAREFULLY

Naled Insecticide

ACTIVE INGREDIENTS:	*Naled	- - -	58%
	Xylene	- - -	20%
INERT INGREDIENTS	- - -	- - -	22%
			TOTAL 100%

1,1-dibromo-2,2-dichloroethyl phosphite.
Registered Trademark of Chevron Chemical Co. U.S. Patent No. 2,971,882
TECHNICAL NALED PER GALLON, EQUIVALENT TO 7.2 POUNDS NALED

KEEP OUT OF REACH OF CHILDREN

SEE BACK PANEL FOR ADDITIONAL CAUTIONS

NET CONTENTS 1 GALLON


E.P.A. Reg. No.
7001-108-AA

E.P.A. Est. 7001-CA-1

MANUFACTURED BY

OCCIDENTAL CHEMICAL COMPANY

P. O. BOX 198 • LATHROP, CALIFORNIA 95330 • PHONE (209) 858-2511



Caterpillars, Leaf
water per acre on
per acre. Do not

(wash): Loopers,
als. diluted spray
over 90° F.

rs, Mites, Citrus
nt. per 100 gals.
of harvest.)

ental Fruit Moth
within 30 days of
nts. per acre.

make more than

ugs (Northwest)

— Use in 1 to 5

y per acre. Flea
as necessary.

er per acre with
picking and con-
): Tomato Fruit
plants thoroughly.
perature is over

TOMATOES, CU-
ounce per 10,000
plates). Apply
he pipes through
followed by im-
ast 1 hour after
ay remain closed
s and approved
g treated houses
plastic or rubber
cucumbers closer
to 4 treatments
ations at 7-day
cessary. **WARN-**
White Butterfly
Poinsettias and

s, Cull Piles and
s.). Apply as a
ngregate. Apply
it or refuse piles
achinery. Do not
y surfaces which

ADULT MOSQUITO, Treat shrubbery and vegetation where mosquitoes may rest. Shrubby and veg-
tation around stagnant pools, marshy areas, ponds and shore lines may be treated. Application of this
product to any body of water is prohibited. **GNAT, STABLEFLY (Dog Flies), AND HOUSEFLY CON-**
TROL -- RESIDENTIAL AREAS, CROPS LISTED ON LABEL, TIDAL MARSHES, SWAMPS, WOODLAND,
FEED LOTS: Consult your State Fish and Game Agency before applying this product. It is not necessary
to avoid farm buildings and dairy barns. Make applications during peak of infestation and repeat as
necessary. See crop recommendations for use limitations near harvest. This rate of application will kill
shrimp. Do not apply to tidal or marsh waters which are important shrimp producing areas. **AIR-**
CRAFT: East — Apply 0.8 to 4 fluid ozs. per acre diluted with water. Apply 2 to 8 quarts of diluted
spray per acre. **West** — Apply 1.6 to 4 fl. ozs. per acre diluted with water. Apply 2 to 8 quarts of
diluted spray per acre. **MIST OR COLD FOG:** Use 3 to 5 qts. per 100 gals. water. Calibrate
equipment (rate of travel and output) to apply 0.1 to 0.25 lb. technical Naled per acre.

CORRALS, HOLDING PENS (Dairy and Beef Cattle, Hogs, Sheep, Horses): Houseflies, Mosquitoes
— **Airplane Application** — Use 1.6 to 4 fluid ounces per acre. Dilute 1 to 2 1/2 pts. in 10 gals. water.
Apply 1 gal. diluted spray per acre. Apply over areas with animals present.

CLOVER MITES, ROACHES, EARWIGS AND LEAFHOPPERS Outside Dwellings and In Lawns: Use 1 fl.
oz. in 3 gals. water. Apply to turf and to soil surfaces around flowers, shrubs and trees for general
pest cleanup.

GRASSHOPPERS: Field Areas (crops listed on label) — 1/2 to 3/4 pt. per acre by air or ground. **Young**
Grasshoppers — 1/2 pt. per acre by air or ground. **Armyworms (except rice)** — 1 pt. per acre by air or
ground. **By Air** — Use in 1 to 5 gals. water per acre. Apply when hoppers hatch is completed. Repeat
as necessary. Animals may be present during treatment. See crop recommendations for use limita-
tions near harvest.

DANGER! Concentrate May Cause Skin Damage. This Formulation Is Corrosive & Causes Eye
Damage. Do Not Get on Skin, Eyes or Clothing. Use Waterproof Gloves and Face
Shield or Goggles When Handling Concentrate.

Harmful If Swallowed. Avoid Breathing Spray Mist. In case of contact, immediately remove contam-
inated clothing and flush skin or eyes with plenty of water; for eyes, get medical attention. Avoid
contamination of feed, foodstuffs and drinking water. Do not use, pour, spill or store near heat or
open flame. Do not use on Nectarines, Ornamental Cherries, Liquidamber or Chrysanthemums as injury
may occur. May be toxic to bees, avoid application during periods of bee activity. Do not store
diluted spray. Make new dilution for each use. **WASH EQUIPMENT THOROUGHLY AFTER USE.**

**Avoid Spillage When Mixing and Filling Spray Equipment. When Container
Is Empty, Immediately Wash Thoroughly and Destroy. Never Re-Use**

Do not apply in edible products areas of food processing plants, restaurants or
other areas where food is commercially prepared or processed. Do not use in ser-
ving areas while food is exposed.

ENVIRONMENTAL CAUTIONS: This product is toxic to fish, shrimp, crab, birds and other wildlife. Birds
feeding on treated areas may be killed. Keep out of lakes, streams, ponds, tidal marshes and estuar-
ies. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift
from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply
this product only as specified on this label.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective in-
formation may be obtained from your Cooperative Agricultural Extension Service. Containers should be
disposed of by breaking and burying with wastes.

CONDITIONS OF SALE: 1. Seller warrants that this product consists of the ingredients specified
and is reasonably fit for use as directed on the label. No one, other than an officer of Seller, is
authorized to make any other warranty, guarantee or direction concerning this product. 2. Because
the time, place, rate of application and other conditions of use are beyond Seller's control, Seller's
liability from the handling, storage and use of this product is limited to replacement of product or
refund of purchase price.