

METHOXYCHLOR 2 EC

You Get Results With

BEST®

ACTIVE INGREDIENTS:

Methoxychlor Technical -----
Petroleum Solvent -----

INERT INGREDIENTS -----

TOTAL

BY WT.

24.9%
71.8%
3.3%
100.0%

* 24.9% Methoxychlor Technical contains 21.9% 2,2-bis-(p-methoxyphenyl)-propane and 3.0% other isomers and reduction products.

E.P.A. Reg. No. 7001-109-AA

Directions for Dilution

DILUTE APPLICATION: Ground Application: Apply specified rate in 20 to 60 gallons of water per acre. Orchard Application: Apply specified rate in 100 to 800 gallons of water per acre.
CONCENTRATE APPLICATION: Ground Application: Apply specified rate in not less than 5 gallons of water per acre. Orchard Application: Apply specified rate in 20 to 100 gallons of water per acre. Special concentrate equipment is necessary for these uses.
AIR APPLICATION: Apply specified rate in 5 to 20 gallons of water per acre. Orchard rates should not be applied in less than 10 gallons of water per acre.

DIRECTIONS FOR USE

Alfalfa, Clover, Cowpea: Alfalfa Caterpillars, Cowpea Curculio, Flea Beetle, Apply 3 to 5 qts. per acre. Alfalfa Weevil (Larvae), Apply 4 to 6 qts. per acre. Armyworm, Apply 6 qts. per acre. Do not apply within 7 days of harvest or grazing.
Apples, Pears: Apple Maggot, Codling Moth, Pear Slug, Tent Caterpillar, Apply 6 to 24 lbs. per acre. Do not apply within 7 days of harvest. For full coverage application.
Asparagus: Asparagus Beetle, Blister Beetle, Apply 4 to 7 qts. per acre. Do not apply within 7 days of harvest.
Beans, Blackeye Peas, Cabbage: Bean Leaf Beetle, Cucumber Beetle, Apply 4 to 10 qts. per acre. Do not apply within 3 days of harvest.
Beets, Broccoli, Brussel Sprouts, Carrots, Collard, Kale, Spinach, Turnips: Cucumber Beetle, Fall Armyworm, Flea Beetle, Apply 4 to 10 qts. per acre. Do not apply within 14 days of harvest.
Blackberries, Blueberries, Boysenberries, Cranberries, Loganberries, Raspberries: Craneberry Fruitworm, Leafhoppers, San Jose Scale (Crawlers), Apply 4 to 10 qts. per acre. Do not apply within 14 days of harvest.
Cantaloupe, Cauliflower, Cucumbers, Kohlrabi, Melons, Peas, Pumpkins, Radishes, Rutabaga, Squash (summer & winter): Blister Beetle, Fall Armyworm, Flea Beetle, Squash Vine Borer, Apply 4 to 10 qts. per acre. Do not apply within 7 days of harvest.
Cocoa, Eggplant, Peppers, Tomatoes: Fall Armyworm, Cucumber Beetle, Apply 4 qts. per acre. Do not apply within 7 days of harvest.
Cherries, Cherries, Fruitfly, San Jose Scale (Crawlers), Apply 6 to 24 qts. per acre. Do not apply within 7 days of harvest. For full coverage application.
Grapes: Grape Berry Moth, Grape Leaf Skeletonizer, Apply 4 to 10 qts. per acre. Do not apply within 14 days of harvest.
Gooseberries, Currants, Huckleberries, Pineapple: Leafhoppers, San Jose Scale (Crawlers), Apply 4 qts. per acre. Do not apply within 14 days of harvest.
Lettuce: Fall Armyworm, Flea Beetle, Apply 4 qts. per acre. Do not apply within 14 days of harvest.

CAUTION!

KEEP OUT OF REACH OF CHILDREN

HARMFUL IF SWALLOWED

AVOID CONTACT WITH SKIN

AVOID BREATHING OF SPRAY MIST

AVOID CONTAMINATION OF FEED AND FOODSTUFFS

Avoid spillage when mixing. When container is empty immediately wash thoroughly and destroy. Never re-use. Discard in a safe place.

Compatibility: Do not use this product with any spray materials which are incompatible with oil. Do not apply within 14 days of Sulfur or Sulfur-product applications.

Warning - Flammable: Do not pour, use, spill or store near heat or open flame.

This product is toxic to bees and should not be applied when bees are actively visiting the area.

This product is toxic to fish. Keep out of any body of water. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

Peaches, Nectarines, Apricots: Plum Curculio, Tent Caterpillar, Apply 6 to 24 qts. per acre. Do not apply within 21 days of harvest.

Potatoes, Sweet Potatoes, Yams: Blister Beetle, Cucumber Beetle, Fall Armyworm, Flea Beetle, Apply 4 qts. per acre. No time limitations.

Strawberries: Leafhoppers, Strawberry Weevil, Apply 4 to 10 qts. per acre. Do not apply within 14 days of harvest.

Plums, Prunes, Quince: Plum Curculio, Tent Caterpillar, Apply 6 to 24 qts. per acre. Do not apply within 7 days of harvest. For full coverage application.

Livestock (except dairy animals): For control of external parasites, use as spray only. Do not use in dipping vats. **Cattle:** Use 1 qt. per 12½ gals. of water. Apply 2 qts. of spray for large animals and proportionally less for smaller ones, for control of hornflies, for control of short-nosed and long-nosed cattle ticks, apply as a drenching spray. **Hogs:** For control of lice, use 1 qt. per 12½ gallons of water applied as a drenching spray.

Self application fly control on cattle (except dairy animals): For control of hornflies, dilute 1 gal. with 4 gals. of diesel fuel or furnace type oil. Do not use any other type of oil. Use 1 gal. of solution to saturate burlap on back rubber, soak burlap every two or three weeks.

Contact and space spray for flies: To reduce population of houseflies and stable flies in farm buildings, in barnyards, dilute 1 pt. with 2½ gals. of water and apply in space by mist blower or directly to flies by hydraulic sprayers, hand sprayer or cylindrical pressure sprayer. Do not use over 1 pt. of Methoxychlor 2 EC per 1000 cubic ft. Do not contaminate milk, feed or drinking water.

4/77 8-12-10w

OCCIDENTAL CHEMICAL COMPANY