

SUPER - SUL

MICRONIZED WETTABLE SULFUR

You Get Results With

BEST

Read This Label Carefully

ACTIVE INGREDIENT:	BY WT.
Sulfur -----	80%
INERT INGREDIENTS: -----	20%
TOTAL	100%

E.P.A. Reg. No. 7001-112-AA

Directions for Dilution

DILUTE APPLICATION: *Ground Application:* Apply specified rate in 20 to 60 gallons of water per acre. *Orchard Application:* Apply specified rate in 100 to 800 gallons of water per acre.

CONCENTRATE APPLICATION: *Ground Application:* Apply specified rate in not less than 5 gallons of water per acre. *Orchard Application:* Apply specified rate in 20 to 100 gallons of water per acre. Special concentrate equipment is necessary for these uses.

AIR APPLICATION: Apply specified rate in 5 to 20 gallons of water per acre. Orchard rates should not be applied in less than 10 gallons of water per acre.

DIRECTIONS FOR USE

Measure needed amount into bucket of water, stir to make thin slurry. Add to tank when part full and before other materials are added. Add wetting agents last to avoid foaming.

Almonds, Cherries, Plums, Prunes: *Brown Rot, Powdery Mildew, Leaf Spot, Silver Mites, Flat Mites, Red Spider.* Apply 10 to 30 lbs. per acre. Apply at bloom or early petal fall. Repeat as necessary, usually 10 to 14 days or after a period of wet weather.

Apples, Pears: *Powdery Mildew, Scab, Blister Mites, Red Spider.* Apply 10 to 30 lbs. per acre. Do not apply to D'anjou pears. Apply pre-bloom and petal fall. Apply with cover sprays throughout the season.

Beans, Peas, Broccoli, Brussels Sprouts, Cabbage, Cauliflower, Collard, Kale, Peppers, Potatoes: *Powdery Mildew, Septoria Leaf Spot, Red Spider.* Apply 3 to 5 lbs. per acre. Start in early leaf stage. Repeat application every 10 to 14 days with or without pesticides immediately after a rain.

Blackberry (Rubis Spp): *Powdery Mildew.* Apply 6 to 15 lbs. per acre. Begin when new shoots are 6 inches long and before blossoms open. Repeat at 10 day intervals or as necessary.

Citrus: *Flat Mite, Silver Mites, Red Spider.* Apply 8 to 20 lbs. per acre. Apply in any spray during season - NOT CONTAINING OIL - or following oil by 14 days!

Cotton, Seed Alfalfa: *Red Spider.* Apply 3 to 5 lbs. per acre.

Grapes: *Powdery Mildew, Bud Mite, Blister Mite, Red Spider.* Apply 2 to 5 lbs. per acre. Apply in late dormant or with cutworm spray, early shoot period and through season on 14 day schedule or with pesticides.

Peaches, Nectarines: *Brown Rot, Powdery Mildew, Silver Mites, Red Spider.* Apply 10 to 30 lbs. per acre. Apply at bloom or early petal fall. Repeat as necessary usually 10 to 14 days or after a period of wet weather.

Peanuts: *Powdery Mildew, Red Spider Mite, Leaf Spot.* Apply 3 to 5 lbs. per acre. Start in early leaf stage. Repeat applications as necessary usually 10 to 24 days

WARNING!

KEEP OUT OF REACH OF CHILDREN

Do not use on sulfur sensitive plants. Sulfur may burn foliage when temperature is excessively high. Do not apply with or following an oil application for 14 days. Do not apply oil for 14 days following treatment.

Burn empty bag - (KEEP OUT OF SMOKE)


Raspberry (Rubis Spp): *Powdery Mildew (Sphaerotheca Humuli).* Apply 6 to 15 lbs. per acre. Spray each week starting at first bloom color. Continue until all fruit is set, then repeat as weather requires.

Strawberries: *Powdery Mildew, Red Spider.* Apply 5 to 10 lbs. per acre. Start in early leaf stage, repeat applications with or without pesticides. For Powdery Mildew begin at first signs of disease and repeat as necessary.

Sugar Beets: *Red Spider (Spp).* Apply 3 to 5 lbs. per acre. Apply as needed with or without pesticides.

Tomatoes: *Russet Mite.* Apply 3 to 5 lbs. per acre. Thorough coverage is required.

Walnuts: *European Red Mite, Pacific Mite, Almond Mite, Two-spotted Mite, Red Spider Mite.* Apply 10 to 20 lbs. per acre. Apply with mite sprays during season to increase the effectiveness of Miticides. **Caution:** certain varieties of walnuts are sensitive to sulfur sprays under certain conditions. Do not apply unless the varieties are known to be tolerant of sulfur in your area.



(701:7/72) rijw

OCCIDENTAL CHEMICAL COMPANY

A DIVISION OF OCCIDENTAL PETROLEUM CORPORATION

P. O. BOX 198 • LATHROP, CALIFORNIA 95330 • PHONE (209) 858-2511

CONDITIONS OF SALE: 1. Seller warrants that this product consists of the ingredients specified and is reasonably fit for use as directed on the label. No one, other than an officer of Seller, is authorized to make any other warranty, guarantee or direction concerning this product. 2. Because the time, place, rate of application and other conditions of use are beyond Seller's control, Seller's liability from the handling, storage and use of this product is limited to replacement of product or refund of purchase price.

SUPER - SUL

MICRONIZED WETTABLE SULFUR



ACTIVE INGREDIENT	BY WT.
Sulfur	80%
INERT INGREDIENTS:	20%
TOTAL	100%

E.P.A. Reg. No. 7001-112-AA

Directions for Dilution

DILUTE APPLICATION - Ground Application: Apply specified rate in 20 to 60 gallons of water per acre.
Orchard Application: Apply specified rate in 100 to 800 gallons of water per acre.
CONCENTRATE APPLICATION - Ground Application: Apply specified rate in not less than 5 gallons of water per acre.
Orchard Application: Apply specified rate in 20 to 100 gallons of water per acre.
Special application equipment is necessary for these uses.
AIR APPLICATION: Apply specified rate in 5 to 20 gallons of water per acre. Orchard rates should not be applied in less than 10 gallons of water per acre.

DIRECTIONS FOR USE

Measure needed amount into bucket of water, stir to make thin slurry. Add to tank when part full and before other materials are added. Add wetting agents last to avoid foaming.

Almonds, Cherries, Plums, Prunes: *Brown Rot, Powdery Mildew, Leaf Spot, Silver Mites, Flat Mite, Spider Mites.* Apply 10 to 20 lbs. per acre. Apply at bloom or early petal fall. Repeat applications as necessary usually 10 to 14 days or after a period of wet weather.

Apples, Pears: *Brown Rot, Powdery Mildew, Silver Mites, Red Spider.* Apply 10 to 30 lbs. per acre. Apply at bloom or early petal fall. Repeat applications as necessary usually 10 to 14 days or after a period of wet weather.

Beans, Peas, Broccoli, Brussels Sprouts, Cabbage, Cauliflower, Collard, Kale, Peppers, Potatoes: *Powdery Mildew, Silver Mites, Leaf Spot, Red Spider.* Apply 3 to 5 lbs. per acre. Start in early leaf stage. Repeat applications as necessary usually 10 to 14 days with or without pesticides immediately after a rain.

Blackberry (Rubis Sp.): *Powdery Mildew.* Apply 10 to 15 lbs. per acre. Begin when new shoots are 6 inches long and before blossoms open. Repeat at 10 day intervals or as necessary.

Citrus: *Flat Mite, Silver Mites, Red Spider.* Apply 8 to 20 lbs. per acre. Apply in any spray during season - **NOT CONTAINING OIL** - or following oil by 14 days.

Cotton, Seed Alfalfa: *Red Spider.* Apply 3 to 5 lbs. per acre.

Grapes: *Powdery Mildew, Bad Mite, Blister Mite, Red Spider.* Apply 2 to 5 lbs. per acre. Apply in late dormant or with cutworm spray, early shoot period and through season on 14 day schedule or with pesticides.

Peaches, Nectarines: *Brown Rot, Powdery Mildew, Silver Mites, Red Spider.* Apply 10 to 30 lbs. per acre. Apply at bloom or early petal fall. Repeat as necessary usually 10 to 14 days or after a period of wet weather.

Peanuts: *Powdery Mildew, Red Spider Mite, Leaf Spot.* Apply 3 to 5 lbs. per acre. Start in early leaf stage. Repeat applications as necessary usually 10 to 24 days.

WARNING!

KEEP OUT OF REACH OF CHILDREN

Do not use on sulfur sensitive plants. Sulfur may burn foliage when temperature is excessively high. Do not apply with or following an oil application for 14 days. Do not apply oil for 14 days following treatment.

Burn empty bag - (KEEP OUT OF SMOKE)


Raspberry (Rubis Sp.): *Powdery Mildew, Silver Mites, Red Spider.* Apply 10 to 20 lbs. per acre. Spray each week starting at first bloom. Repeat applications as necessary usually 10 to 14 days or after a period of wet weather.

Strawberries: *Powdery Mildew, Red Spider.* Apply 10 to 20 lbs. per acre. Apply at bloom or early petal fall. Repeat applications with or without pesticides as necessary usually 10 to 14 days or after a period of wet weather.

Sugar Beets: *Red Spider (Str).* Apply 3 to 5 lbs. per acre. Use only with low toxicity pesticides.

Tomatoes: *Russian Mite.* Apply 3 to 5 lbs. per acre. Thorough coverage is essential.

Walnuts: *European Red Mite, Pacific Mite, Armand Mite, Two-spotted Mite, Red Spider Mite.* Apply 10 to 20 lbs. per acre. Apply with mite sprays during season to increase the effectiveness of Mitecides. **Caution:** Certain varieties of walnuts are sensitive to sulfur sprays under certain conditions. Do not apply unless the varieties are known to be tolerant of sulfur in your area.



(701:7 72) rjw

OCCIDENTAL CHEMICAL COMPANY

P.O. BOX 198 LATHROP, CALIFORNIA 95026 PHONE (209) 752-1111