




SUPER - SUL

MICRONIZED WETTABLE SULFUR



ACTIVE INGREDIENT:	BY WT.
Sulfur	80%
INERT INGREDIENTS:	
TOTAL	100%

EPA Reg. No. 7001-112-AA
EPA Est No. 14800-FL-1

KEEP OUT OF REACH OF CHILDREN WARNING

- Do not breathe dust or spray mist. Do not get in eyes, on skin or on clothing
- IF IN EYES immediately flush with plenty of water for at least 15 minutes. Call a physician
- IF INHALED and overcome by vapor or dust remove immediately from exposure and call a physician. Administer artificial respiration if breathing is irregular or has stopped
- IF LARGE QUANTITIES ARE SWALLOWED give water and induce vomiting. Get medical attention
- FOR SKIN CONTACT flush with water, use soap if available
- Do not use on sulfur sensitive plants. Sulfur may burn foliage when temperature is excessively high. Do not apply with or following an oil application for 14 days. Do not apply oil for 14 days following treatment

STORAGE & DISPOSAL

- Do not store near food or feed. To protect fish and wildlife, do not contaminate streams, lakes or ponds with this material.
- Burn empty bag (Keep Out Of Smoke)

DIRECTIONS FOR DILUTION

DILUTE APPLICATION: *Ground Application:* Apply specified rate in 20 to 60 gallons of water per acre. *Orchard Application:* Apply specified rate in 100 to 800 gallons of water per acre.
CONCENTRATE APPLICATION: *Ground Application:* Apply specified rate in not less than 5 gallons of water per acre. *Orchard Application:* Apply specified rate in 20 to 100 gallons of water per acre. Special concentrate equipment is necessary for these uses.
AIR APPLICATION: Apply specified rate in 5 to 20 gallons of water per acre. Orchard rates should not be applied in less than 10 gallons of water per acre

NET WT. 30 LBS.

DIRECTIONS FOR USE

Measure needed amount into bucket of water, stir to make thin slurry. Add to tank when part full and before other materials are added. Add wetting agents last to avoid foaming.

- Almonds, Cherries, Plums, Prunes:** *Brown Rot, Powdery Mildew, Leaf Spot, Silver Mites, Flat Mites, Red Spider Mites.* Apply 10 to 30 lbs per acre. Apply at bloom or early petal fall. Repeat as necessary, usually 10 to 14 days or after a period of wet weather
- Apples, Pears:** *Powdery Mildew, Scab, Blister Mites, Red Spider Mites.* Apply 10 to 30 lbs per acre. Do not apply to D'Anjou pears. Apply pre-bloom and petal fall. Apply with cover sprays throughout the season
- Beans, Peas, Broccoli, Brussels Sprouts, Cabbage, Cauliflower, Collard, Kale, Peppers, Potatoes:** *Powdery Mildew, Septoria, Leaf Spot, Red Spider Mite.* Apply 3 to 5 lbs per acre. Start in early leaf stage. Repeat application every 10 to 14 days with or without pesticides immediately after a rain
- Blackberry (Rubis Spp):** *Powdery Mildew.* Apply 6 to 15 lbs per acre. Begin when new shoots are 6 inches long and before blossoms open. Repeat at 10 day intervals or as necessary
- Citrus:** *Flat Mite, Scale Mites, Red Spider Mite.* Apply 8 to 20 lbs per acre. Apply in any spray during season — **NOT CONTAINING OIL** — or following oil by 14 days
- Cotton, Seed Alfalfa:** *Red Spider Mite.* Apply 3 to 5 lbs per acre
- Grapes:** *Powdery Mildew, Bug Mite, Blister Mite, Red Spider Mite.* Apply 2 to 5 lbs per acre. Apply in late dormant or with cutworm spray, early shoot period and through season on 14 day schedule or with pesticides
- Peaches, Nectarines:** *Brown Rot, Powdery Mildew, Silver Mites, Red Spider Mite.* Apply 10 to 30 lbs per acre. Apply at bloom or early petal fall. Repeat as necessary usually 10 to 14 days or after a period of wet weather
- Peanuts:** *Powdery Mildew, Red Spider Mite, Leaf Spot.* Apply 3 to 5 lbs per acre. Start in early leaf stage. Repeat applications as necessary usually 10 to 24 days
- Raspberry (Rubis Spp):** *Powdery Mildew (Sphaerotheca Humuli).* Apply 6 to 15 lbs per acre. Spray each week starting at first bloom color. Continue until all fruit is set, then repeat as weather requires
- Strawberries:** *Powdery Mildew, Red Spider Mite.* Apply 5 to 10 lbs per acre. Start in early leaf stage, repeat applications with or without pesticides. For *Powdery Mildew* begin at first signs of disease and repeat as necessary
- Sugar Beets:** *Red Spider Mite (Spp).* Apply 3 to 5 lbs per acre. Apply as needed with or without pesticides
- Tomatoes:** *Russet Mite.* Apply 3 to 5 lbs per acre. Thorough coverage is required
- Walnuts:** *European Red Mite, Pacific Mite, Almond Mite, Two spotted Mite, Red Spider Mite.* Apply 10 to 20 lbs per acre. Apply with mite sprays during season to increase the effectiveness of acaricides. **Caution:** Certain varieties of walnuts are sensitive to sulfur sprays under certain conditions. Do not apply unless the varieties are known to be tolerant of sulfur in your area.

(701:3/76)

Occidental Chemical Company

P.O. Box 198, Lathrop, California 95330 P.O. Box 1185, Houston, Texas 77001

DISCLAIMER OF WARRANTIES: Seller warrants that the chemical composition of this product conforms to the chemical description given on this label. THIS WARRANTY IS EXPRESSLY IN LIEU OF ALL OTHER WARRANTIES AND REPRESENTATIONS, EXPRESSED, IMPLIED OR STATUTORY, INCLUDING WARRANTIES OF MERCHANTABILITY AND FITNESS FOR A PARTICULAR USE. Timing, rate and method of application, weather and crop conditions, mixtures with chemicals not specifically recommended on this label or an accompanying written recommendation, are beyond the control of seller. Buyer assumes all risks of use, storage and handling of this material not in strict accordance with directions given hereon. Buyer further agrees in the event of damages arising from the use of this product to accept a replacement of the product or a refund of the purchase price of the product, at buyer's option as a charge of seller's liability. No one is authorized to make any other warranty, guarantee or directions concerning this product, and no such warranty, guarantee or directions shall be valid or binding upon seller.