

NU-LOX 5 BAIT

You Get Results With

BEST®

(CONTAINS DYLOX*)

ACTIVE INGREDIENT

Dimethyl (2,2,2-Trichloro-1-hydroxyethyl) phosphonate** ----- 5%

INERT INGREDIENTS: ----- 95%

TOTAL 100%

*Dylox is a registered trademark in U.S. Patent Offs. by Farbenfabriken Bayer A.G., Chemagro Corporation, licensee

**U.S. Patent No. 2,701,225

DIRECTIONS FOR USE

For control of surface-feeding *cutworms*, *salt marsh caterpillars*, *armyworms*, and *crickets* in the following crops. Applications may be made in a band over planted row, around the stem of individual plants or by broadcast treatment.

FIELD CROPS: Apply 2 lbs. per acre unless stated otherwise.

Alfalfa: Do not apply more than one application per cutting, nor within 14 days of cutting or grazing.

Barley, Flax, Oats, Wheat: Do not apply within 21 days of harvest. Do not graze treated fields. Do not feed treated straw to dairy or meat animals.

Corn: Apply once per season when plants are 3 to 12 inches high. Do not use corn for feed, food or ensilage within 45 days of treatment.

Cotton: Apply 5 lbs. per acre. Do not apply within 7 days of picking. Do not graze livestock in treated fields within 14 days after application.

Peanuts: Three applications may be made before digging. One additional application may be made between digging and harvest.

Sugarbeets: Apply 30 lbs. per acre. Do not apply within 28 days of harvest, if tops are to be fed to livestock. Do not apply within 14 days of harvest if tops are not to be fed to livestock.

Safflower: Apply 30 lbs. per acre. Do not apply after the plants begin to bloom. Do not apply more than twice per year.

Tobacco: Do not apply after plants are 2 feet high.

VEGETABLES

Beans (Dry & Snap): Apply 30 lbs. per acre. Do not apply within 14 days of harvest. Do not feed treated vines to meat or dairy animals.

Beans (Lima): Apply 30 lbs. per acre. Do not apply more than twice per season, nor harvest for food, feed or forage within 30 days of treatment.

Cowpeas (Southern Peas, Blackeyed Peas, Crowder Peas): Apply 30 lbs. per acre. Do not apply more than twice per season, nor within 14 days of harvest. Do not feed treated vines to meat or dairy animals.

Brussels Sprouts, Cabbage, Cauliflower, Peppers: Apply 20 lbs. per acre. Do not apply within 21 days of harvest.

Collards, Lettuce, Tomatoes: Apply 20 lbs. per acre. Do not apply within 28 days of harvest. Do not graze treated fields.

OCCIDENTAL CHEMICAL COMPANY

CAUTION:

KEEP OUT OF REACH OF CHILDREN

May be harmful if swallowed, inhaled, or absorbed through the skin. Avoid contact with skin. Do not breathe dust. Wash thoroughly with soap and warm water after handling. Wash contaminated clothing with soap and hot water before re-use.

Do not contaminate feed or food.

If poisoning occurs, obtain prompt medical aid. Prolonged exposure will result in cholinesterase depression.

TO PHYSICIAN: Atropine sulfate is antidotal. 2-Pam is also antidotal and may be administered in conjunction with atropine.

Rinse equipment and containers and dispose of waste by burying in non-crop land away from water supplies. Dispose of empty containers by burying with wastes or by burning. (KEEP OUT OF SMOKE)

This product is toxic to fish, birds, and other wildlife. Keep out of any body of water. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes.

Pumpkins: Apply 20 lbs. per acre. Do not apply more than once per season, nor within 14 days of harvest.

Table Beets: Apply 30 lbs. per acre. Do not apply within 28 days of harvest. Do not eat tops.

ACCEPTED

E.P.A. Reg. No. 7001-113-AA

(686.7 72) rjw