

ACCEPTED

August 28, 1972

UNDER THE FEDERAL INSECTICIDE ACT
FOR THE CONTROL OF PESTS OR WEEDS
EPA REG. NO. 7001-146-AA

Read all label directions carefully before applying.

GENERAL INFORMATION

Read all label directions carefully before applying. Mildew Spray is a new fungicide for the eradication and prevention of powdery mildew on roses. Used as directed, Mildew Spray will not harm rose flowers or foliage. Apply with any accurate hose-end sprayer; tank sprayer or power sprayer.

DANGER: KEEP OUT OF THE REACH OF CHILDREN. Harmful if swallowed. May cause irritation of eyes, nose, throat or skin. Avoid breathing of spray mist. Do not get in eyes, on skin or clothing. Causes eye damage. Wear goggles and rubber gloves when handling. In case of eye contact, flush with plenty of water and if irritation persists, get medical attention. Avoid contamination of feed or food-stuffs and fish ponds. Do not put concentrate or dilute into food or drink containers. When container is empty, wash thoroughly and destroy.

WARNING: Do not use, pour, spill or store near heat or open flame.

NOTICE: Buyer assumes all responsibility for safety and use, storage or handling not in accordance with directions.



Occidental Chemical Company

A DIVISION OF OCCIDENTAL PETROLEUM CORPORATION
POST OFFICE BOX 198 • LATHROP, CALIF. 95330



Professional Grade

FOR PROFESSIONAL TURF AND GROUNDS MANAGERS

MILDEW SPRAY



READ THIS LABEL CAREFULLY

EPA Reg. No. 7001-146-AA

Contains Parnon®

A fungicide for the eradication and prevention of powdery mildew on roses, zinnias and non-bearing apples and grapes

Active Ingredient:

*a,a-Bis (p-chlorophenyl)-3-pyridinemethanol 4.0%

Inert Ingredients 96.0%

*Parnon® is the trademark for Farnol, Bianco Products Co., a division of Eli Lilly and Co.

- PRODUCTS FOR:
- INDUSTRY
- SCHOOLS
- HIGHWAYS
- GOLF COURSES
- PARKS
- CEMETERIES
- ATHLETIC FIELDS
- PEST CONTROL OPERATORS

NET CONTENTS 1 GALLON

DANGER!

KEEP OUT OF REACH OF CHILDREN

SEE BACK OF LABEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

M
a
i
n
R
H
th
7

L
i
r
o
p
d
a
f
o

A
P
e
o
f
v
a
l
n
e
Z
I
1
t
a
t
e
s
o
f
a



Professional Grade

FOR PROFESSIONAL TURF AND GROUNDS MANAGERS

MILDEW SPRAY



READ THIS LABEL CAREFULLY

EPA Reg. No. 7001-146-AA

Contains Parnon[®]

A fungicide for the eradication and prevention of powdery mildew on roses, zinnias and non-bearing apples and grapes

Active Ingredient:		
*a.a-Bis (p-chlorophenyl)-3-pyridinemethanol	4.0%	
Inert Ingredients	96.0%	

*Parnon[®] is the trademark for Parnol, Cianco Products Co., a division of Eli Lilly and Co

NET CONTENTS 1 GALLON

DANGER!

KEEP OUT OF REACH OF CHILDREN

SEE BACK OF LABEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

PRODUCTS FOR:

- INDUSTRY**
- SCHOOLS**
- HIGHWAYS**
- GOLF COURSES**
- PARKS**
- CEMETERIES**
- ATHLETIC FIELDS**
- PEST CONTROL OPERATORS**

APPLICATION DIRECTIONS

Mix Mildew Spray with water at the rates listed on the Dilution Chart and apply to the plant foliage. Thoroughly cover all plant surfaces, including new growth when spraying.

ROSES:

Heavy Infestation Program: If powdery mildew is already present on the roses, spray the roses at the rate on the spray chart. Repeat every 7 days until mildew is controlled, then change to the Light Program.

Dilution Rates for Heavy Infestation Program:

1 tsp. per 1 gallon of water.

16 fl. oz. (1 pint) per 100 gallons of water.

Light Infestation Program: To insure mildew free roses, start spraying roses as soon as new growth has fully expanded in the spring at the preventive rate as shown on the Dilution Chart. Repeat every 7 to 10 days during the spring and fall. Repeat this application as required for continuous powdery mildew control.

Dilution Rates for Light Infestation Program:

1 tsp. per 2 gallons of water.

8 fl. oz. (1/2 pint) per 100 gallons of water.

APPLE TREES and GRAPEVINES (Non-Bearing): Use 8 fl. oz. (1/2 pint) per 100 gallons of water. For smaller amount, use 1 tsp. per 2 gallons of water. Begin before infection occurs. Repeat at 1 to 2 week intervals, as long as disease condition exists or as necessary to protect all new growth. Apply to insure thorough coverage of all plant surfaces.

ZINNIAS: Use 4 oz. per 125 gallons of water. For smaller amount, use 1 tsp. per 5 gallons of water. Begin before infection occurs. Repeat at 1 to 2 week intervals, as long as disease condition exists or as necessary to protect all new growth. Apply to insure thorough coverage of all plant surfaces.

re applying.

ON

ig.
on and prevention of
ildew Spray will not
accurate hose-end

CHILDREN. Harmful if
throat or skin. Avoid
on skin or clothing.
gloves when hand-
water and if irritation
tion of feed or food-
r dilute into food or
ash thoroughly and

heat or open flame.
ety and use, storage

Company
LEUM CORPORATION
IOP, CALIF. 95330