

You Get Results With

BEST[®]

MP-TOX 2-8 AQUAMUL

METHYL PARATHION MIXTURE, LIQUID

ACTIVE INGREDIENTS:

- 1. Toxaphene
- 2. O,O-Dimethyl O-p-nitrophenyl thiophosphate

INERT INGREDIENTS:

- 1. Tetrahydro-Cyclized Camphene (Chlorine content 67% to 69%)
- 2. Methyl Parathion

EPA Reg No 7001-153-AA

Directions for Dilution

DILUTE APPLICATION

- Ground Application: Apply 1.5 lbs concentrate per 100 gallons of water per acre.
- Orchard Application: Apply 1.5 lbs concentrate per 100 gallons of water per acre.

CONCENTRATE APPLICATION

- Ground Application: Apply 1.5 lbs concentrate per 100 gallons of water per acre.
- Orchard Application: Apply 1.5 lbs concentrate per 100 gallons of water per acre.

AIR APPLICATION

- Use only concentrate equipment specifically designed for these uses.
- Apply 1.5 lbs concentrate per 100 gallons of water per acre for ground use and 1.5 lbs concentrate per 100 gallons of water per acre for air use.

DIRECTIONS FOR USE

APPLES: Apply 1.5 lbs concentrate per 100 gallons of water per acre. Do not apply after harvest. Do not use after 15 days after harvest. Do not use on fruit trees that are heavily infested with codling moth.

APPLES: Apply 1.5 lbs concentrate per 100 gallons of water per acre. Do not apply after harvest. Do not use after 15 days after harvest. Do not use on fruit trees that are heavily infested with codling moth.

BROSSELETS: Apply 1.5 lbs concentrate per 100 gallons of water per acre. Do not apply after harvest. Do not use after 15 days after harvest. Do not use on fruit trees that are heavily infested with codling moth.

BRUSSELETS: Apply 1.5 lbs concentrate per 100 gallons of water per acre. Do not apply after harvest. Do not use after 15 days after harvest. Do not use on fruit trees that are heavily infested with codling moth.

CAUTION: Do not apply within 7 days of harvest. Do not use on fruit trees that are heavily infested with codling moth.

CAUTION: Do not apply within 7 days of harvest. Do not use on fruit trees that are heavily infested with codling moth.

CAUTION: Do not apply within 7 days of harvest. Do not use on fruit trees that are heavily infested with codling moth.

OCCIDENTAL CHEMICAL COMPANY

BY WT.

66.7
16.6
16.7
TOTAL 100.0%



Poisoning

Symptoms:

POISON



HEADACHE, DIARRHEA, DISCOMFORT IN CHEST, WEAKNESS, NAUSEA, CRAMPS, BLURRED VISION

CALL A PHYSICIAN IMMEDIATELY!

ANTIDOTES: If swallowed: Give at once plenty of water to drink. If inhaled: Give at once plenty of fresh air. If on skin: Wash with soap and water. If in eyes: Flush with water.

INFORMATION FOR PHYSICIAN (TREATMENT)

15-2 to 4 mg/kg of body weight of atropine sulfate should be given immediately. Repeat every 30 minutes until the patient is no longer bradycardic. Give at once plenty of fresh air. If on skin: Wash with soap and water. If in eyes: Flush with water.

DO NOT GET IN EYES, ON SKIN, OR ON CLOTHING POISONOUS IF SWALLOWED, INHALED, OR ABSORBED THROUGH SKIN OR EYES

DANGER!

KEEP OUT OF REACH OF CHILDREN

WARNING: Pesticide. Avoid contact with skin, eyes, nose, mouth, and clothing. Do not breathe dust or fumes. Do not get in eyes, on skin, or on clothing. Do not swallow. Do not use in confined spaces. Do not use near food or feed. Do not use on fruit trees that are heavily infested with codling moth.

CAUTION: Do not apply within 7 days of harvest. Do not use on fruit trees that are heavily infested with codling moth.

PEPPERS: Apply 1.5 lbs concentrate per 100 gallons of water per acre. Do not apply after harvest. Do not use after 15 days after harvest. Do not use on fruit trees that are heavily infested with codling moth.