

**BEST**

### ROSE MILDEW CONTROL

Read all label directions carefully before applying.

#### GENERAL INFORMATION

Rose Mildew Control is a new fungicide for the eradication and prevention of powdery mildew on roses. Used as directed, Rose Mildew Control will not harm rose flowers or foliage. Apply with any accurate hose-end sprayer; tank sprayer or power sprayer.

**WARNING:** KEEP OUT OF THE REACH OF CHILDREN. Harmful if swallowed. May cause irritation of eyes, nose, throat or skin. Avoid breathing of spray mist. Do not get in eyes, on skin or on clothing. In case of eye contact, flush with plenty of water and if irritation persists, get medical attention. Avoid contamination of feed or foodstuffs and fish ponds. Do not put concentrate or dilute into food or drink containers. When container is empty, wash thoroughly and destroy. Never reuse.

**WARNING:** Do not use, pour, spill or store near heat or open flame.

**NOTICE:** Buyer assumes all responsibility for safety and use, storage or handling not in accordance with directions.



**Occidental Chemical Company**

U.S. Pat. No. 7001-154-AM Label No. 1172

You get results with

**BEST**

\$2.29

# ROSE Mildew Control



Contains PARNON<sup>®</sup>\*

A fungicide for the eradication and prevention of powdery mildew on roses.

**ACTIVE INGREDIENT**

\*a Bis-p chlorophenyl-3 pyridinemethanol 1.3%

**INERT INGREDIENTS** 98.7%

\*Parnon<sup>®</sup> is the trademark for Par-nol, Eianco Products Co., a division of Eli Lilly and Co.

**WARNING:** KEEP OUT OF THE REACH OF CHILDREN. See back of label for additional precautionary statements.

**NET 8 FL. OZ. (1/2 PT.)**

#### APPLICATION DIRECTIONS

Mix Rose Mildew Control with water at the rates listed on the Dilution Chart and apply to the plant foliage. Thoroughly cover all plant surfaces, including new growth when spraying.

#### ROSES

##### Eradicant Program:

If powdery mildew is already present on the roses, spray the roses at the rate on the spray chart. Repeat every 7 days until mildew is controlled, then change to the preventive program.

##### Dilution rates for Eradicant Program:

1 oz. (2 tblsp.) per 1 gal. of water.  
6 fl. oz. (12 tblsp.) per 6 gals. of water.

##### Preventive Program:

To insure mildew free roses, start spraying roses as soon as new growth has fully expanded in the spring at the preventive rate as shown on the Dilution Chart. Repeat every 7 to 10 days during the spring and fall. Repeat this application as required for continuous powdery mildew control.

##### Dilution rates for Preventive Program:

1 tblsp. per 1 gal. of water.  
8 fl. oz. (1/2 pint) per 16 gals. of water.

ET-88

RECEIVED

APR 26 1973

PLANT FOLIAGE