

FRONT PANEL 50-A

You Get Results With
BEST

Covers up to
2500
Sq. Ft.

**TURF DISEASE
CONTROL**

For control of Large Brown Patch, Curvularia,
Gray Leaf Spot, *Diplodia* "Leaf Spot",
Melting Out, Copper Spot, Snow Mold and Stem
Rust in turf.



ACCEPTED
UNDER U.S. PATENT NOS. 3,200,153 AND 3,311,731
FUNGICIDE AND FERTILIZER
FC
ED 100-157-22
10-157-22

ACTIVE INGREDIENT	
Chlorothalonil (tetrachloroisophthalonitrile)*	2.55
INERT INGREDIENTS	97.55

*Product contains chlorothalonil covered under U. S. Patent Nos. 3,200,153 and 3,311,731
CAUTION: KEEP OUT OF THE REACH OF CHILDREN. See back panel for additional cautions.

E.P.A. Reg. No. 7001-157-AA

NET WEIGHT 25 Lbs.

FRONT PANEL 50-A

You Get Results With
BEST

Covers up to
2500
Sq. Ft.

TURF DISEASE
CONTROL

For control of Large Brown Patch, Curvularia,
Gray Leaf Spot, Leptothosporium Leaf Spot,
Melting Out, Copper Spot, Snow Mold and Stem
Rust in turf.



ACCEPTED
UNDER THE FEDERAL INSECTICIDE AND FUNGICIDE ACT
ED. 100-157-157
TO AVOID REJECTION

ACTIVE INGREDIENT
Chlorothalonil (tetrachloroisophthalonitrile)*----- 2.55
INERT INGREDIENTS-----97.55

*Product contains chlorothalonil covered under U. S. Patent
Nos. 3,290,353 and 3,221,735
CAUTION: KEEP OUT OF THE REACH OF CHILDREN. See back panel
for additional cautions.

E.P.A. Reg. No. 7001-157-AA

NET WEIGHT 20 Lbs.

You Get Results With
BEST

TURF DISEASE CONTROL

For control of Large Brown Patch, Curvularia, Gray Leaf Spot, Dollar Spot, Helminthosporium Leaf Spot, Melting Out, Copper Spot, Snow Mold and Stem Rust in turf.

DIRECTIONS FOR USE

- 1) Apply when lawn is wet.
- 2) Do not water or mow for at least 24 hours.
- 3) Adjust spreader setting to apply at proper rate.
NOTE: Always calibrate spreaders and check for accuracy before using. Age and condition of spreaders, speed of pushing, roughness of terrain may affect application rate. Operator should carefully calculate area and make certain that correct amounts of material are applied. Suggested spreader settings:

Spreader	Preventative Rate 8 lbs./1000 sq. ft.	Curative Rate 12 lbs./1000 sq. ft.
Central #1622E (Imperial Mark)	6-1/4	7-3/4
Scotts Model #35-3	6-1/2	8
Sears Model #1919	6-1/2	8
MTD #431-340	4-3/4	5-3/4
Wards Model #89-1021	4-3/4	5-3/4

- 4) Large Brown Patch, Curvularia, Gray Leaf Spot, Helminthosporium Leaf Spot, Melting Out:
As a preventative - Use 8 lbs. per 1,000 sq. ft. Begin before diseases are expected to appear. Repeat at 7 to 14 day intervals throughout season.

As a curative - Use 12 lbs. per 1,000 sq. ft. Apply at 7 day intervals.
- 5) Copper Spot, Dollar Spot: As a preventative - use 8 lbs. per 1,000 sq. ft. Begin before diseases appear. Repeat at 7 to 10 day intervals. As a curative or under severe conditions favoring development of disease - Use 12 lbs. per 1,000 sq. ft. Apply at 7 day intervals.

52-4

BACK PANEL (Cont'd.)

- 6) Snow Mold (Oregon & Washington only): As a preventative or curative -- Use 12 lbs. per 1,000 sq. ft. Apply in the fall. Repeat at 7 to 14 day intervals as necessary.
- 7) Stem Rust of Bluegrass: As a preventative - Use 8 lbs. per 1,000 sq. ft. Begin before disease appears. Repeat at 7 to 14 day intervals.
As a curative or under severe conditions - Use 12 lbs. per 1,000 sq. ft.

CAUTION: KEEP OUT OF REACH OF CHILDREN. Do not take internally. Avoid contact with skin, eyes or clothing. Can be irritating to eyes or skin. Wash thoroughly with soap and warm water after handling. In case of contact with eyes, flush with plenty of water for at least 15 minutes. Avoid breathing of dust.

Do not store near feed or foodstuffs.

This product is toxic to fish. Keep out of lakes, ponds, and streams.

Do not contaminate water by cleaning of equipment or disposal of wastes.

Apply only to areas specified on the label.

Do not reuse empty container. Destroy by burying with waste or burning in a safe place. Stay away from smoke or fumes.

USE CAUTIONS

Apply when lawn is damp. Do not water or mow for at least 24 hours after application. Do not graze treated areas. Do not feed clippings to livestock.

CONDITIONS OF SALE: Seller warrants that this product consists of the ingredients specified and is reasonably fit for use as directed on the label. No one is authorized to make any other warranty, guarantee or direction concerning this product. Because the time, place, rate of application and other conditions of use are beyond Seller's control, Seller's liability from the handling, storage and use of this product is limited to replacement of product or refund of purchase price.

OXY
LOGO

OCCIDENTAL CHEMICAL COMPANY
A Division of Occidental Petroleum Corporation
P. O. Box 198 . Bethrop, California 95330