


6451



2001-160

1-3-72

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL WARNINGS AND DIRECTIONS.

Malathion Insect Spray gives control of many common insects. Spray thoroughly to cover both upper and lower leaf surfaces and all infested portions of plants. Do not store diluted spray. Make new dilution for every use. Application may be made with any type of sprayer.

NOTICE: Buyer assumes all responsibility for safety and use not in accordance with directions.

WARNING: DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME.

WARNING: Keep out of reach of children. May be fatal if swallowed. May be absorbed through the skin. Do not breathe spray mist. Avoid contact with skin, eyes and clothing. Wash thoroughly after handling. Avoid contamination of feed and foodstuffs. Do not use on household pets or humans. Do not permit children or pets to go onto sprayed grass until spray has completely dried. Keep out of reach of domestic animals. Do not contaminate ornamental fishponds. Do not use on ferns. May injure foliage of certain varieties of apples and cherries.

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label. When container is empty, wash thoroughly and destroy. Never re-use.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

Do not use in edible products areas of food processing plants, restaurants or other areas where food is commercially prepared or processed. Do not use in serving areas while food is exposed.



Occidental Chemical Company

A DIVISION OF OCCIDENTAL PETROLEUM CORPORATION
POST OFFICE BOX 196 • LATHROP, CALIF. 95330



Pro

FOR

M

READ

EPA

ACTIVE INGREDIENTS
Malathion
Xylene

INERT INGREDIENTS
*O,O-dimethyl

For control of
White Fly and

NET C

PRODUCTS FOR:

INDUSTRY

SCHOOLS

HIGHWAYS

GOLF COURSES

PARKS

CEMETERIES

ATHLETIC FIELDS

PEST CONTROL OPERATORS

CORDANCE WITH LABEL

common insects. Spray surfaces and all infested take new dilution for every sprayer.

safety and use not in ac-

IRE NEAR HEAT OR OPEN

be fatal if swallowed. May spray mist. Avoid contact after handling. Avoid con- on household pets or hu- o sprayed grass until spray estic animals. Do not con- ferns. May injure foliage

streams or ponds. Do not disposal of wastes. Apply in container is empty, wash

o direct treatment or resi- obtained from your Coop-

rocessing plants, restaurants areared or processed. Do not

al Company

ROLEUM CORPORATION
HROP, CALIF 95330



Professional Grade

FOR PROFESSIONAL TURF AND GROUNDS MANAGERS

Malathion Insect Spray

READ THIS LABEL CAREFULLY

EPA Reg. No. 7001-160-AA

ACTIVE INGREDIENTS:

Malathion*	- - - - -	50.0%
Xylene	- - - - -	35.4%
INERT INGREDIENTS:	- - - - -	14.6%

*O,O-dimethyl dithiophosphate of diethyl mercaptosuccinate


NOT FOR HOME USE

For control of Aphids, Red Spider Mites, Thrips, Leafhoppers, Scale, White Fly and Mealybugs.

NET CONTENTS 1 GALLON

WARNING:

**KEEP OUT OF REACH OF CHILDREN
SEE ADDITIONAL CAUTIONS ON BACK PANEL**



- PRODUCTS FOR:**
- INDUSTRY**
 - SCHOOLS**
 - HIGHWAYS**
 - GOLF COURSES**
 - PARKS**
 - CEMETERIES**
 - ATHLETIC FIELDS**
 - PEST CONTROL OPERATORS**

- ① **ORNAMENTALS*** (Roses, Azalea flowers and shrubs). *Aphids:* 2 1/2 tablespoons per gallon of water. *Spider Mites, Mealybugs, Woolly A:* 1 tablespoon per gallon of water. *Pine Scale:* 1 1/2 to 2 tablespoo
 - ② **FRUITS & NUTS:** Numbers () and harvest.
 - ③ *Apples, (3), Pears (1) — Aphids:* 2 1/2 tablespoons per gallon of water. *Whi* per gallon of water. ③ *Apricots,* 2 1/2 tablespoons per gallon of water. *nuts (1) — Aphids:* 2 teaspoons
 - ④ **CITRUS FRUITS** (Grapefruit, Lemons) Apply within 7 days of harvest. *Bla* *Thrips:* 2 teaspoons per gallon
 - ⑤ **VEGETABLES:** Numbers () and harvest. Cole crops including *Cauliflower (7), Peas (3), Potat* (1) — *Aphids:* 2 teaspoons per tablespoon per gallon of water.
 - ⑥ **HOUSEHOLD PESTS** in and a *Silverfish, Clover Mites, Earwig.* Apply with a coarse spray or under sinks and to other areas, cracks and crevices. For ants i and window frames where the clover mites also treat outside areas. Avoid contamination of and water. Repeat applications
- * Dilution equivalents for large
1 tablespoon/gallon 3 qt. per
2 teaspoons/gallon 1 qt. per

Professional Grade

FOR PROFESSIONAL TURF AND GROUNDS MANAGERS

Malathion Insect Spray

READ THIS LABEL CAREFULLY

EPA Reg. No. 7001-160-AA

ACTIVE INGREDIENTS:

Malathion*	- - - - -	50.0%
Xylene	- - - - -	35.4%

INERT INGREDIENTS:	- - - - -	14.6%
--------------------	-----------	-------

*O,O-dimethyl dithiophosphate of diethyl mercaptosuccinate


NOT FOR HOME USE

For control of Aphids, Red Spider Mites, Thrips, Leafhoppers, Scale, White Fly and Mealybugs.

NET CONTENTS 1 GALLON

WARNING:

**KEEP OUT OF REACH OF CHILDREN
SEE ADDITIONAL CAUTIONS ON BACK PANEL**



① **ORNAMENTALS*** (Roses, Azaleas, Camellias, Evergreens and many other flowers and shrubs). Aphids: 2 teaspoons per gallon of water. Red Spider Mites, Mealybugs, Woolly Aphids, White Fly, Thrips, Rose Leafhoppers: 1 tablespoon per gallon of water. Black Scale, Soft Brown Scale, Monterey Pine Scale: 1/2 to 2 tablespoons per gallon of water.

FRUITS & NUTS: Numbers () indicate days required between last spray and harvest.

② Apples (3), Pears (1) — Aphids: 2 teaspoons per gallon of water. Red Spider Mites, Woolly Aphids, White Flies, Thrips, Pear Psylla: 1 tablespoon per gallon of water. Apricots (7), Cherries (3), Peaches (7) — Aphids: 2 teaspoons per gallon of water. Almonds (1), Filberts (1), Pecans (1), Walnuts (1) — Aphids: 2 teaspoons per gallon of water.

③ **CITRUS FRUITS** (Grapefruit, Lemons, Limes, Oranges, Kumquats). Do not apply within 7 days of harvest. Black Scale, California Red Scale, Purple Scale, Thrips: 2 teaspoons per gallon of water. Do not apply during fullbloom.

④ **VEGETABLES:** Numbers () indicate days required between last spray and harvest. Cole crops including Broccoli, Brussel Sprouts, Cabbage and Cauliflower (7), Peas (3), Potatoes (1), Squash (1), Melons (1), Tomatoes (1) — Aphids: 2 teaspoons per gallon of water. Beans (1) — Aphids: 1 tablespoon per gallon of water.

⑤ **HOUSEHOLD PESTS** in and about homes — Roaches, Ants, Mosquitoes, Silverfish, Clover Mites, Earwigs: 8 tablespoons (4 oz.) per gallon of water. Apply with a coarse spray or paint brush to window frames, baseboards, under sinks and to other areas frequented by pests. For roaches also treat cracks and crevices. For ants include the treatment on out trails, door sills and window frames where the pest enters the building. For mosquitos and clover mites also treat outside areas around doors, windows, and porch areas. Avoid contamination of food, dishes, utensils, milk, milk equipment and water. Repeat applications as necessary.

* **Dilution equivalents for larger sprayers:**

1 tablespoon/gallon 3 qt. per 100 gallons
2 teaspoons/gallon 1 qt. per 100 gallons