

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL WARNINGS AND DIRECTIONS.

Malathion Insect Spray gives control of many common insects. Spray thoroughly to cover both upper and lower leaf surfaces and all infested portions of plants. Do not store diluted spray. Make new dilution for every use. Application may be made with any type of sprayer.

NOTICE: Buyer assumes all responsibility for safety and use not in accordance with directions.

WARNING: DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME

WARNING: Keep out of reach of children. May be fatal if swallowed. May be absorbed through the skin. Do not breathe spray mist. Avoid contact with skin, eyes and clothing. Wash thoroughly after handling. Avoid contamination of feed and foodstuffs. Do not use on household pets or humans. Do not permit children or pets to go onto sprayed grass until spray has completely dried. Keep out of reach of domestic animals. Do not contaminate ornamental fishponds. Do not use on ferns. May injure foliage of certain varieties of apples and cherries.

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on the label. When container is empty, wash thoroughly and destroy. Never reuse.

This product is highly toxic to bees exposed to direct treatment or to the honey crop. For protective information, may be obtained from your County Extension Agent or Extension Service.

For more information on the use of this product on ornamental plants, contact your local extension agent or contact the nearest extension office. Do not use on plants where the use is prohibited by local, state or federal law.



Occidental Chemical Company

1000 WEST 10TH AVENUE, DENVER, COLORADO 80202



Professional Grade

FOR PROFESSIONAL TURF AND GROUNDS MANAGERS

Malathion Insect Spray

READ THIS LABEL CAREFULLY

EPA Reg. No 7001-160-AA

ACTIVE INGREDIENTS

Malathion*	50.0%
Xylene	35.4%

INERT INGREDIENTS

10-O-dimethyl dithiophosphate of primary amine, etc.	14.6%
--	-------

NOT FOR HOME USE

For control of Aphids, Beet Beetles, Fleas, Ground Squirrels, Japanese Beetles, Leaf Beetles, Mosquitoes, White Fly and Mealybugs.

NET CONTENTS 1 GALLON

WARNING.

KEEP OUT OF REACH OF CHILDREN

PRODUCTS FOR
INDUSTRY
SCHOOLS
HIGHWAYS
GOLF COURSE
PARKS
CEMETERIES
ATHLETIC FIELDS
RESIDENTIAL AREAS

ORNAM
flowers
der Mit
1 table
Pine Sc
FRUITS
and hal
Apples
der Mit
per gal
teaspoc
nuts (1)
CITRUS
piy with
Thrips:
VEGET
harvest
Caulflo
1 A
tablesp
HOUSE
Silverfis
Apply v
water:
ticks
and win
and
and wal
Dilutio
table
and



Professional Grade

FOR PROFESSIONAL TURF AND GROUNDS MANAGERS

Malathion Insect Spray

READ THIS LABEL CAREFULLY

EPA Reg. No 7001-160-AA

ACTIVE INGREDIENTS:		
Malathion*	- - - - -	50.0%
Xylene	- - - - -	35.4%
INERT INGREDIENTS - - - - - 14.6%		
*O,O-dimethyl dithiophosphate of diethyl mercaptosuccinate		

NOT FOR HOME USE

For control of Aphids, Red Spider Mites, Thrips, Leafhoppers, Scale, White Fly and Mealybugs

NET CONTENTS 1 GALLON

WARNING:

KEEP OUT OF REACH OF CHILDREN
Hazardous to Aquatic Life



- PRODUCTS FOR**
INDUSTRY
SCHOOLS
HIGHWAYS
GOLF COURSES
PARKS
CEMETERIES
ATHLETIC FIELDS
PEST CONTROL OPERATORS

ORNAMENTALS* (Roses, Azaleas, Camellias, Evergreens and many other flowers and shrubs). *Aphids*: 2 teaspoons per gallon of water. *Red Spider Mites, Mealybugs, Woolly Aphids, White Fly, Thrips, Rose Leafhoppers*: 1 tablespoon per gallon of water. *Black Scale, Soft Brown Scale, Monterey Pine Scale*: 1/2 to 2 tablespoons per gallon of water.

FRUITS & NUTS: Numbers () indicate days required between last spray and harvest.

Apples (3), Pears (1) — *Aphids*: 2 teaspoons per gallon of water. *Red Spider Mites, Woolly Aphids, White Flies, Thrips, Pear Psylla*: 1 tablespoon per gallon of water. Apricots (7), Cherries (3), Peaches (7) — *Aphids*: 2 teaspoons per gallon of water. Almonds (1), Filberts (1), Pecans (1), Walnuts (1) — *Aphids*: 2 teaspoons per gallon of water.

CITRUS FRUITS (Grapefruit, Lemons, Limes, Oranges, Kumquats): Do not apply within 7 days of harvest. *Black Scale, California Red Scale, Purple Scale, Thrips*: 2 teaspoons per gallon of water. Do not apply during fullbloom.

VEGETABLES: Numbers () indicate days required between last spray and harvest. Cold crops including Broccoli, Brussel Sprouts, Cabbage and Cauliflower (7), Peas (3), Potatoes (1), Squash (1), Melons (1), Tomatoes (1) — *Aphids*: 2 teaspoons per gallon of water. Beans (1) — *Aphids*: 1 tablespoon per gallon of water.

HOUSEHOLD PESTS in and about homes: *Roaches, Ants, Mosquitoes, Silverfish, Clover Mites, Earwigs*: 8 tablespoons (4 oz) per gallon of water. Apply with a coarse spray or paint brush to window frames, baseboards, under sinks and to other areas frequented by pests. For roaches, also treat cracks and crevices. For ants include the treatment of out doors, door mats and window frame where the pest enters the building. For earwigs and clover mites also treat outdoors in groups, door, windows and porch areas. Avoid contamination of food, dishes, garden and lawns with spray and water. Repeat application as necessary.

* Dilution equivalents for larger sprayers:

1 tablespoon/gallon = 1 qt per 100 gal in
1 teaspoon/gallon = 1 qt per 1000 gal in