

SEVIN 5 BAIT

CRUMBLES

You Get Results With

BEST®

ACTIVE INGREDIENTS:

Carbaryl (1-naphthyl N-methylcarbamate) ----- 5%

INERT INGREDIENTS: ----- 95%

TOTAL 100%

BY WT.

CAUTION

KEEP OUT OF REACH OF CHILDREN

HARMFUL IF INHALED OR SWALLOWED

*Trademark of Union Carbide Corp. for the active ingredients
1-naphthyl N-methylcarbamate

E.P.A. Reg. No. 7001-168-AA

DIRECTIONS FOR USE

Broadcast application may be made by ground equipment or aircraft. Treatments may be repeated as necessary. Apply recommended dosage for control of cutworms, armyworms, crickets, darkling ground beetles, grasshoppers and saw bugs.

Alfalfa, Peas: Apply 50 lbs. per acre. No time limitation.

Asparagus, Strawberries: Apply 40 lbs. per acre. Do not apply within 1 day of harvest.

Beans, Carrots, Corn, Eggplant, Onions, Peppers, Potatoes, Tomatoes, Sweet Corn: Apply 40 lbs. per acre. No time limitation.

Broccoli, Brussels Sprouts, Cabbage, Cauliflower, Head Lettuce, Garden Beets (roots), Horseradish, Parsnips, Radishes, Rutabagas, Turnips (roots): Apply 40 lbs. per acre. Do not apply within 3 days of harvest.

Cotton: Apply 30 to 40 lbs. per acre. No time limitation.

Cucumbers, Squash, Melons, Peanuts: Apply 20 lbs. per acre. No time limitation.

Sugar Beets, Chard, Endive, Escarole, Garden Beets (leaves), Kale, Leaf Lettuce, Parsley, Spinach, Swiss Chard, Turnips (tops): Apply 40 lbs. per acre. Do not apply within 14 days of harvest.

Flower Trees and Vineyards: Apply 40 to 45 lbs. per acre to crops listed below to control ants, crickets, cutworms, grasshoppers, saw bugs. A broadcast application may be made by ground equipment or aircraft.

Apples	Nectarines
Apricots	Peaches
Cherries	Pears
Citrus	Plums
Grapes	

Do not apply within 5 days of harvest in citrus, 7 days in apricots, nectarines and 1 day in apples, cherries, peaches, pears, plums. No time limitation on grapes.

Avoid breathing of dust. Do not take internally. Avoid contact with skin. Wear regular long-sleeved work clothing. Change to clean clothing daily. Wash hands and face before eating. Wash thoroughly after handling.

NOTE TO PHYSICIAN: Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal.

Avoid contamination of food, feed, water supplies, streams and ponds. Keep pets off treated areas.

Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.



Label on container: Apply 40 to 45 lbs. per acre to crops listed below to control ants, crickets, cutworms, grasshoppers, saw bugs. A broadcast application may be made by ground equipment or aircraft.

OCCIDENTAL CHEMICAL COMPANY

P. O. BOX 178 LATHROP, CALIFORNIA 95330 PHONE (916) 858-2511