

READ ENTIRE LABEL. USE STRICTLY ACCORDANCE WITH LABEL CAUTIONS, WARNINGS AND DIRECTIONS.

Keep pesticide in original container. Do not put concentrate or dilute into food or drink containers.

DIRECTIONS

For best results spray to cover all exposed plant parts and repeat as necessary. Do not spray when temperature is over 80°. All spraying should be done early in the morning to avoid burning. After spraying remove excess spray material from the foliage by tapping branches with 2 thin sticks. Do not store diluted spray. Make new dilution for each use. HORTICULTURAL CAUTION: Do not use on apricots as they are susceptible to sulfur injury. Do not use on evergreen plants. Do not apply in freezing weather.

DORMANT SPRAYING

ROSES: Rose Rust, Black Spot—Prune, destroy all old leaves. Spray bushes at 10 tablespoonfuls (5 oz.) per gallon of water.

PEACH LEAF CURL, PEACH BLIGHT: Peaches—10 tablespoonfuls (5 oz.) per gallon of water in late fall before heavy rains—also before buds begin to swell in late winter.

BLISTER MITE: Pears—10 tablespoonfuls (5 oz.) per gallon of water just as buds begin to swell in spring and late fall before leaves start to turn color.

LEAF SPOT, CANE BLIGHT, RED BERRY TROUBLE (Recognized by Dwarf Berries): Caneberries, BLACKBERRIES, GOOSEBERRIES, RASPBERRIES—10 tablespoonfuls (5 oz.) per gallon of water in late fall, followed by another treatment in spring as buds begin to swell.

COMBINATION WINTER SPRAY

DILUTION: Use 1½ pints of spray and ½ pint of BEST Oil Spray in 6 gallons of water (7½ tablespoonfuls BEST Lime Sulfur Spray and 2½ tablespoonfuls BEST Oil Spray per gallon). Keep agitated during spraying.

ROSES AND ORNAMENTALS: (Dormant, deciduous only)—rose scale, oyster shell scale, brown apricot scale, black scale, over-wintering spider mite and aphid eggs, many over-wintering fungus spores or plant diseases.

FRUIT TREES (APPLES, CHERRIES, NECTARINES, PEACHES, PLUMS, PRUNES, QUINCE):—Dormant, deciduous only—San Jose scale, brown apricot scale, over-wintering spider mite and aphid eggs, many over-wintering fungus spores or plant diseases.

GROWING SEASON SPRAYING

POWDERY MILDEW: Roses, delphiniums, lilacs, euonymus, columbine, crepe myrtle, sweet peas, zinnias, western grapes, fruits, ornamentals, tuberoses, begonias—2 teaspoonfuls per gallon of water on tender new growth and 3 teaspoonfuls per gallon of water on mature foliage. Apply every 1 to 2 weeks.

LAWN AND SOIL TREATMENT

LAWNS, GARDENS: To condition alkaline lawn and garden soils and improve water penetration and moisture retention in heavy soils, apply 1 pint in 7½ to 15 gallons of water on 500 sq. ft. of area. Apply with sprayer or watering can. After application, water lawn immediately. Apply every 60 days throughout spring and summer season as needed.

SOIL CORRECTION FOR ACID LOVING PLANTS:—Rhododendrons, Azaleas, Camellias, 1 teaspoonful per gallon of water. Apply ½ gallon diluted material per plant to surface of soil around plant. Repeat treatment every 2 to 3 months until plants regain proper growth.

CAUTION: KEEP OUT OF THE REACH OF CHILDREN. May cause skin irritation. Avoid contact with skin. Wash hands after use. If material gets into eyes immediately wash with water. If eye irritation persists contact doctor at once.

Pick open flowers before application to avoid discoloration. Avoid spraying on painted surfaces to avoid staining. However, if accidentally sprayed on painted surfaces, considerable bleaching out will occur. If crust occurs on surface, break with a stick. Crystallization will not impair product quality. Keep bottle tightly closed when not in use. Do not use 30 days before or after using an oil spray during growing season or on plants in leaf.

Lime sulfur should not be used within 10-14 days of an oil spray at any stage. When container is empty, immediately wash thoroughly and destroy. Never reuse.

NOTICE: Buyer assumes all responsibility for safety and use not in accordance with directions. Seller makes no warranty express or implied concerning the use of this product other than indicated on the label. Because the time, place, rate of application and other conditions are beyond the seller's control, seller's liability from handling and use of this product is limited to replacement of product or refund of purchase price.

E.P.A. Reg. No. 7001-172-AA

E.P.A. Est. No. 7001-CA-1



Occidental Chemical Company

DIVISION OF OCCIDENTAL PETROLEUM CORPORATION
POST OFFICE BOX 198 • LATHROP, CALIF. 95330

Label No. 214

You Get Results With
BEST

\$1.98

ACCIDENTAL Sulfur Spray



Aug 2, 1974
UNDER THE FEDERAL INSECTICIDE ACT
FUNGICIDE AND RODENTICIDE
FOR ECONOMIC POISON
E.P.A. REG. NO. 7001-172-AA

LIME-SULFUR FUNGICIDE
For growing season and dormant spraying to control certain plant diseases. Controls Rose powdery mildew and Peach leaf curl. Also highly effective as soil conditioner and corrective.

ACTIVE INGREDIENT:
Calcium Polysulfide 29.0%
INERT INGREDIENTS: 71.0%
DENSITY: Baume at 60 F. 31
Lbs. per gal. at 68 F. 10.6 lbs.

CAUTION: KEEP OUT OF REACH OF CHILDREN. See back panel for additional cautions.

NET CONTENTS 16 FL. OZS. (1 PT.)

4558