

CAUTION

KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed. Avoid inhaling spray mist. This material may cause skin irritation. Wash hands and face thoroughly after using. Do not store near heat or open flame. Use with adequate ventilation. Keep container closed. Avoid contamination of feed and foodstuffs.

CAUTION COMBUSTIBLE LIQUID

DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME

DESTROY THIS CONTAINER WHEN EMPTY - NEVER REUSE

Use Cautions - This product is toxic to fish. Keep out of lakes, streams, or ponds. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

Do not plant any root crop not registered for PCNB in rotation on PCNB treated soil

DIRECTIONS FOR USE

Beans: Root & Stem Rot - *Rhizoctonia solani*. Furrow - Application rates are to 14,500 linear feet of row for bush type beans and to 8,430 linear feet of row for pole beans. Apply 2 to 3 quarts in 8 to 10 gallons of water. Spray planting furrow and covering soil at planting time. Avoid applying directly to seed; otherwise delayed emergence may occur. In California and Arizona apply 3 to 4 pints in 8 to 10 gallons of water. Apply in furrow and incorporate with covering soil at planting. Use lower rate for lighter soils. Avoid application to bare seed.

Snap & Dry Beans: White Mold - *Sclerotinia sclerotiorum*. Use 1 gallon mixed in 8 to 10 gallons of water and apply to 14,500 linear feet of row for bush type beans or to 8,430 linear feet of row for pole beans. Spray in an 8 inch wide band, centered on the row, immediately after or at time of seeding. Avoid applying directly to seed; otherwise delayed emergence may occur. (Where dinitro type pre-emergence herbicides are used, add 1 gallon to the final herbicide spray mix and apply simultaneously). If disease is severe, application may be repeated at 2 to 3 week intervals, using 1 or 2 nozzles per row and directly spray at base of plants. Do not apply after first bloom. Soil should remain undisturbed after treatment. Do not feed treated vines to livestock.

Cotton: Damping Off - *Rhizoctonia solani*. Use 1 to 1 gallon per acre in at least 15 gallons of water (per 12,500 linear feet of row based on 40 inch spacings). Spray on seed and

You Get Results With

BEST[®]

TERRACLOR^{*} 2 EC

READ THIS LABEL CAREFULLY

ACTIVE INGREDIENT:

Pentachloronitrobenzene 24.0%

INERT INGREDIENTS: 76.0%

TOTAL 100.0%

Contains 2 lbs. Pentachloronitrobenzene per U.S. gallon at 68°F

^{*}Registered trademark of Olin Corporation

E.P.A. Reg. No. 7001-175-AA

E.P.A. Est. No. 7001-CA-1

CAUTION: KEEP OUT OF REACH OF CHILDREN

PROTECT FROM TEMPERATURES BELOW 20°F

SEE ADDITIONAL CAUTIONS ON SIDE PANEL

NET CONTENTS 1 GALLON OCCIDENTAL CHEMICAL COMPANY

DIVISION OF OCCIDENTAL PETROLEUM CORPORATION

LATHROP, CALIFORNIA

surrou
by liv
Garlic
and a
chine
bottom
the co
Potato
100 to
disc a
the di
tuber
moist.
ed in
points
and pe
placed
acre);
Spray
ORNAM
Azalea
plants
to oper
Lawns:
August
t. 1,000
obtain
to insu
unusua
should
discol
appear
For dic
sq. ft.

CONDITIO
reasonabl
make any
rate of
handling,

You Get Results With

BEST®

TERRACLOR® 2 EC

READ THIS LABEL CAREFULLY

ACTIVE INGREDIENT:

Pentachloronitrobenzene 24.0%

INERT INGREDIENTS: 76.0%
TOTAL 100.0%

Contains 2 lbs. Pentachloronitrobenzene per U.S. gallon at 68°F

*Registered trademark of Occ. Corporation

E.P.A. Reg. No. 7001-175-AA

E.P.A. Est. No. 7001-CA-1

CAUTION: KEEP OUT OF REACH OF CHILDREN

PROTECT FROM TEMPERATURES BELOW 20°F

SEE ADDITIONAL CAUTIONS ON SIDE PANEL

NET CONTENTS 1 GALLON
OCCIDENTAL CHEMICAL COMPANY

DIVISION OF OCCIDENTAL PETROLEUM CORPORATION

LATHROP, CALIFORNIA

surrounding soil at planting time. Do not allow the feeding or grazing of cotton foliage by livestock.

Garlic: White Rot - *Sclerotium cepivorum*. Use 10 gallons mixed in 100 gallons of water and apply to 21,800 linear feet of row at planting time. Attach a suitable rig to the machine planter, using 2 nozzles per planting furrow. Direct the front nozzle to spray the bottom of the open furrow and the "seed" as it is dropped. Direct the rear nozzle to spray the covering soil.

Potatoes: *F. zootonia Solani*. Broadcast Application - Apply 9 to 12½ gallons mixed with 100 to 150 gallons of water to one acre. Apply to the soil surface, prior to planting and disc and cross disc to a depth of 4 to 6 inches. A drag harrow is usually attached behind the discs on the last discing. It is very important to thoroughly mix the Terraclor into the tuber forming zone of the soil. It is desirable to make treatment when the soil is slightly moist. Row (in the furrow) Application - Use 5 gallons per 12,400 linear feet of row mixed in sufficient water to total approximately 20 gallons of finished spray. Spray several points in the planting row to treat all the soil that surrounds and covers the seed piece and potato forming zone. Three nozzles attached to the planter work best; one fan nozzle placed above and slightly ahead of closing discs (approximately 10 gals. of spray per acre); one fan nozzle to rear of closing discs (approximately 5 gals. of spray per acre). Spray pattern should treat 12 to 14 inch band centered on the row.

ORNAMENTALS

Azalea & Camellia: Azalea Petal Blight, Camellia Flower Blight. Apply as a drench under plants at a rate of 1½ quarts in 30 gallons of water per 150 sq. ft. Begin application prior to opening buds and repeat every 3 to 4 weeks through the bloom period.

Lawns: Brown Patch - *Rhizoctonia Solani*. For Southern or warm season lawns such as St. Augustine, Bermuda and similar grasses - make one application in the fall or spring at first indication of infection. Mix 1 quart in sufficient water (10 to 15 gals.) to cover 1,000 sq. ft. Apply by means of a sprinkler can, spray or any other convenient method of obtaining even coverage. Following application, the treated area should be lightly watered to insure movement through the grass to the soil level. If the treated area is subjected to unusually heavy rainfall, or flooded, or if the disease is severe or reappears, the area should be retreated 3 to 4 weeks later. Under certain growing conditions, a temporary discoloration of the grass occasionally may occur. This causes no harm and will disappear in a short time. Do not graze treated areas. Do not feed clippings to livestock.

For dichondra: Treat as above, using 1½ quarts mixed in 40 gallons of water per 1,000 sq. ft.

CONDITIONS OF SALE: 1. Seller warrants that this product consists of the ingredients specified and is reasonably fit for use as directed on the label. No one, other than an officer of Seller, is authorized to make any other warranty, guarantee or direction concerning this product. 2. Because the time, place, rate of application and other conditions of use are beyond Seller's control, Seller's liability from the handling, storage and use of this product is limited to replacement of product or refund of purchase price.