

You Get Results With

BEST

DUSTING SULFUR

Read This Label Carefully

ACTIVE INGREDIENTS:

Sulfur

INERT INGREDIENTS:

BY WT.

98%

TOTAL 2%
100%

E.P.A. Reg. No. 7001-181-AA

E.P.A. Est. No. 7001-CA-1

CAUTION

KEEP OUT OF REACH OF CHILDREN

Sulfur may cause severe fruit and foliage injury to certain crops. Sulfur sensitive plants include anjou and comice pears, apricots, cranberries, certain varieties of cucurbits, filberts, spinach, tung, walnuts, and certain ornamentals. Apples, pears and strawberries are susceptible to foliage burn with sulfur under certain climatic conditions. Sulfur may burn foliage when temperature is high. Do not apply if temperatures during or within 3 days after application are expected to exceed 90°F in the area to be treated. Do not use within 2 weeks of an oil spray treatment, on citrus within 3 to 6 weeks. When growing crops for processing consult the processor before applying sulfur.

Do not reuse container. Destroy when empty.

DIRECTION FOR USE

Can be applied by ground or air application.

FRUIT & NUT CROPS

Almonds: brown rot, scab, shothole, mite suppression - Apply 40-45 lbs. per acre. Begin when disease threatens or in pre-bloom stage and repeat at 7 day intervals or as necessary.

Apples: Apply 30-40 lbs. per acre in pre-bloom, bloom and cover periods as necessary.

Apples: apple cedar rust, black rot, brown rot, frog eye leaf spot, powdery mildew, quince rust, scab, sooty blotch - apply 30-60 lbs. per acre from pre-pink through cover. Caution: MacIntosh, golden delicious, Jonathan and certain other varieties may be injured by sulfur application under certain conditions.

Avocado: mite suppression - apply 40 to 50 lbs. per acre.

Blackberries, Boysenberries, Dewberries, Loganberries, Raspberries: powdery mildew - apply 20-30 lbs. per acre. Begin when new shoots are 6 inches long and before blossoms open. Repeat at 10 day intervals or as necessary.

Currants, Gooseberries, Blueberries & Huckleberries: powdery mildew - apply 20-30 lbs. per acre. Begin when new shoots are 6 inches long and before blossoms open. Repeat at 10 day intervals or as necessary.

Cherries: brown rot, leaf spot, powdery mildew, rust, scab, stem blight - apply 35-60 lbs. per acre. Begin in pink stage. Repeat at bloom, petal fall and in cover stages at 7-14 day intervals and as a post harvest foliar treatment for leaf spot.

Citrus: citrus flat mite (Brevipalpus) - Apply 100 lbs. per acre. Citrus red spider mite - apply 80 lbs. per acre. Two spotted mite - apply 50-60 lbs. per acre.

Figs: fig rust mite - apply 40-50 lbs. per acre.

Grapes: powdery mildew - apply 10-40 lbs. per acre. Rust - apply 15-40 lbs. per acre. Begin when new shoots are 6 inches long. Repeat before blossoms open and continue as necessary. Caution: Concord and other labrusca type grapes may be injured by sulfur.

Nectarines & Peaches: brown rot, leaf spot, powdery mildew, rust, scab, shothole - apply 35-60 lbs. per acre. Apply at pink, bloom, petal fall, shuck split, shuck fall and at 10-14 day intervals as long as necessary. For brown rot of fruit, apply when fruit starts to ripen.

Pears: brown rot, leaf spot, powdery mildew, rust, scab - apply 20-60 lbs. per acre at pre-bloom, bloom, petal fall and cover periods.

Plums & Prunes: brown rot, leaf spot, powdery mildew, rust, scab, shothole - apply 35-60 lbs. per acre beginning at late dormant stage for shothole, at pink stage for other diseases. Repeat at bloom, petal fall, shuck fall and in cover periods as necessary. For brown rot of fruit, dust when fruit starts to ripen.

Quince: brown rot, powdery mildew, scab - apply 35-60 lbs. per acre. Begin before diseases are expected to appear. Repeat at 7 to 10 day intervals or as necessary.

Strawberries: powdery mildew - apply 20-50 lbs. per acre. Caution: sulfur may injure strawberries under certain climatic conditions. Begin at first sign of disease and repeat at 7 to 10 day intervals or as necessary.

VEGETABLE CROPS

Artichokes: leaf spot - apply 25-35 lbs. per acre. Begin when disease first appears and repeat at 7 to 10 day intervals or as necessary.

Asparagus: rust - apply 20-35 lbs. per acre. Begin after cutting stops. Repeat at 7-10 day intervals through season.

Beans: leaf spot, powdery mildew, rust - apply 20-35 lbs. per acre. Begin when first true leaves form or at first sign of disease. Repeat at 7-10 day intervals. Caution: sulfur may injure certain varieties.

Beets: rust - apply 20 lbs. per acre. Apply at first signs of disease. Repeat at 7 to 10 day intervals or as necessary.

Broccoli, Brussels Sprouts, Cabbage, Cauliflower, Collards, Kale, Rutabagas, Turnips: powdery mildew - apply 20-25 lbs. per acre. Begin at first sign of disease. Repeat at 7 to 10 day intervals or as necessary.

Carrots: powdery mildew, rust - apply 20-50 lbs. per acre. Begin at first sign of disease and repeat at 7-10 day intervals.

Celery: early and late blight - apply 6-12 lbs. per acre. Begin as soon as plant emerges. Repeat at 7 to 10 day intervals or as necessary.

Eggplants: powdery mildew - apply 20-35 lbs. per acre. Begin when first true leaves appear. Repeat at weekly intervals.

Garlic: downy and powdery mildew - apply 20-35 lbs. per acre before disease is expected to appear. Repeat at 7 to 10 day intervals or as necessary.

Lettuce: powdery mildew - apply 10 to 15 lbs. per acre. Begin at first sign of disease. Repeat as necessary.

Onions: botrytis blight, downy and powdery mildew - apply 20 to 50 lbs. per acre. Begin before disease is expected to appear. Repeat as necessary.

Pears: powdery mildew, rust - apply 20 to 35 lbs. per acre. Begin early in season before disease appears. Repeat at 7 to 10 day intervals.

Peppers: powdery mildew - apply 20-35 lbs. per acre. Begin before disease appears. Repeat at weekly intervals.

Potatoes: powdery mildew - apply 5-15 lbs. per acre. Apply before disease appears. Repeat at 7 to 10 day intervals or as necessary.

Spinach: powdery mildew, rust - apply 8-30 lbs. per acre. Apply when disease first appears. Repeat at 7-10 day intervals.

Tomatoes: russet mite, two spotted mite - apply 20-50 lbs. per acre.

FIELD CROPS

Alfalfa: powdery mildew, rust - apply 20-25 lbs. per acre. 6 inches high or when disease appears. Repeat at 7 to 14 days.

Barley, Oats, Wheat: rust - apply 25 to 40 lbs. per acre. 10 day intervals.

Clover: powdery mildew, rust - apply 20 lbs. per acre before disease. Repeat at 7 to 10 day intervals or as necessary.

Cotton: rust - apply 20-30 lbs. per acre. Begin when first signs appear. Repeat at 7-10 day intervals.

Cowpeas: rust - apply 20-25 lbs. per acre. Begin soon after through season.

Flax: powdery mildew - apply 20-30 lbs. per acre. Begin intervals or as necessary.

Hops: mite suppression - apply 30-40 lbs. per acre.


Peanuts: leaf spot, powdery mildew - apply 20 to 25 lbs. per acre. 2 days old or when first spots appear on lower leaves after rains. For rust apply 20 to 30 lbs. per acre, 7 to 10 day intervals as necessary.

Soybeans: leaf spot, powdery mildew - apply 8-15 lbs. per acre. 7-10 day intervals.

Vetch: rust - apply 25-40 lbs. per acre. Begin at first signs.

ORNAMENTAL CROPS

Alder, Azalea, Buttonbush, Catalpa, Chrysanthemum, flowers, Sycamore, Zinnias: powdery mildew - apply 20-30 lbs. per acre. Repeat at 5 to 10 day intervals and after rain or as necessary.



P. O. BOX 198 • LATHROP, CALIFORNIA

CONDITIONS OF SALE: 1. Seller warrants ingredients specified and is reasonably fit for use. No one, other than an officer of Seller, is authorized to make any warranty, guarantee or direction concerning time, place, rate of application and other conditions of use. Seller's liability from the use of this product is limited to replacement of product.

DUSTING SULFUR

Read This Label Carefully

BY WT.
98%

2%
TOTAL 100%

CAUTION KEEP OUT OF REACH OF CHILDREN

Injury to certain crops. Sulfur sensitive plants include anjou and varieties of cucurbits, figs, figs, spinach, tung, walnuts and certain varieties are susceptible to foliage burn with sulfur under certain conditions when temperature is high. Do not apply if temperatures during or after application exceed 90°F in the area to be treated. Do not use within 2 weeks within 3 to 6 weeks. When growing crops for processing, consult

INSTRUCTIONS FOR USE

FRUIT & NUT CROPS

Suppression. Apply 40-45 lbs. per acre. Begin when disease first appears. Repeat at 7 day intervals or as necessary.

Apply at bloom and cover periods as necessary.

Apple, brown rot, leaf spot, powdery mildew, quince rust, scab, sooty blotch, pink through cover. Caution: Mac Intosh, golden delicious, Jonared are injured by sulfur application under certain conditions.

Apply 50 lbs. per acre.

Blackberries, Loganberries, Raspberries: powdery mildew. Apply 20 lbs. per acre. Begin when disease first appears. Repeat at 10 day intervals or as necessary.

Huckleberries: powdery mildew. Apply 20-30 lbs. per acre. Begin when disease first appears. Repeat at 10 day intervals or as necessary.

Peaches, Nectarines: brown rot, leaf spot, powdery mildew, rust, scab, shothole. Apply 35-60 lbs. per acre. Begin when disease first appears. Repeat at 7-14 day intervals and as a post harvest application.

Apply 100 lbs. per acre. Citrus red spider mite. Apply 80 lbs. per acre. Repeat at 7-10 day intervals or as necessary.

Figs: fig rust mite. Apply 40-50 lbs. per acre.

Grapes: powdery mildew. Apply 10-40 lbs. per acre. Rust. Apply 15-40 lbs. per acre. Begin when new shoots are 6 inches long. Repeat before blossoms open and continue as necessary. Caution: Concord and other labrusca type grapes may be injured by sulfur.

Nectarines & Peaches: brown rot, leaf spot, powdery mildew, rust, scab, shothole. Apply 35-60 lbs. per acre. Apply at pink, bloom, petal fall, shuck split, shuck fall and at 10-14 day intervals as long as necessary. For brown rot of fruit, apply when fruit starts to ripen.

Pears: brown rot, leaf spot, powdery mildew, rust, scab. Apply 20-60 lbs. per acre at pre bloom, bloom, petal fall and cover periods.

Plums & Prunes: brown rot, leaf spot, powdery mildew, rust, scab, shothole. Apply 35-60 lbs. per acre. Begin at late dormant stage for shothole, at pink stage for other diseases. Repeat at bloom, petal fall, shuck fall and in cover periods as necessary. For brown rot of fruit, dust when fruit starts to ripen.

Quince: brown rot, powdery mildew, scab. Apply 35-60 lbs. per acre. Begin before diseases are expected to appear. Repeat at 7 to 10 day intervals or as necessary.

Strawberries: powdery mildew. Apply 20-50 lbs. per acre. Caution: sulfur may injure strawberries under certain climatic conditions. Begin at first sign of disease and repeat at 7 to 10 day intervals or as necessary.

VEGETABLE CROPS

Artichokes: leaf spot. Apply 25-35 lbs. per acre. Begin when disease first appears and repeat at 7 to 10 day intervals or as necessary.

Asparagus: rust. Apply 20-35 lbs. per acre. Begin after cutting stops. Repeat at 7-10 day intervals through season.

Beans: leaf spot, powdery mildew, rust. Apply 20-35 lbs. per acre. Begin when first true leaves form or at first sign of disease. Repeat at 7-10 day intervals. Caution: sulfur may injure certain varieties.

Beets: rust. Apply 20 lbs. per acre. Apply at first signs of disease. Repeat at 7 to 10 day intervals or as necessary.

Broccoli, Brussels Sprouts, Cabbage, Cauliflower, Collards, Kale, Rutabagas, Turnips: powdery mildew. Apply 20-25 lbs. per acre. Begin at first sign of disease. Repeat at 7 to 10 day intervals or as necessary.

Carrots: powdery mildew, rust. Apply 20-50 lbs. per acre. Begin at first sign of disease and repeat at 7-10 day intervals.

Celery: early and late blight. Apply 6-12 lbs. per acre. Begin as soon as plant emerges. Repeat at 7 to 10 day intervals or as necessary.

Eggplants: powdery mildew. Apply 20-35 lbs. per acre. Begin when first true leaves appear. Repeat at weekly intervals.

Garlic: downy and powdery mildew. Apply 20-35 lbs. per acre before disease is expected to appear. Repeat at 7 to 10 day intervals or as necessary.

Lettuce: powdery mildew. Apply 10 to 15 lbs. per acre. Begin at first sign of disease. Repeat as necessary.

Onions: botrytis blight, downy and powdery mildew. Apply 20 to 50 lbs. per acre. Begin before disease is expected to appear. Repeat as necessary.

Peas: powdery mildew, rust. Apply 20 to 35 lbs. per acre. Begin early in season before disease appears. Repeat at 7 to 10 day intervals.

Peppers: powdery mildew. Apply 20-35 lbs. per acre. Begin before disease appears. Repeat at weekly intervals.

Potatoes: powdery mildew. Apply 5-15 lbs. per acre. Apply before disease appears. Repeat at 7 to 10 day intervals or as necessary.

Spinach: powdery mildew, rust. Apply 8-30 lbs. per acre. Apply when disease first appears. Repeat at 7-10 day intervals.

Tomatoes: russet mite, two spotted mite. Apply 20-50 lbs. per acre.

FIELD CROPS

Alfalfa: powdery mildew, rust. Apply 20-25 lbs. per acre. Apply to stubble or when crop is not over 3 to 6 inches high or when disease appears. Repeat at 7 to 14 day intervals or as necessary.

Barley, Oats, Wheat: rust. Apply 25 to 40 lbs. per acre. Apply at first sign of disease. Repeat at 7 to 10 day intervals.

Clover: powdery mildew, rust. Apply 20 lbs. per acre beginning when crop is in bud stage or at first sign of disease. Repeat at 7 to 10 day intervals or as necessary.

Cotton: rust. Apply 20-30 lbs. per acre. Begin when first true leaves form or when disease threatens. Repeat at 7-10 day intervals.

Cowpeas: rust. Apply 20-25 lbs. per acre. Begin soon after seedling emerge. Repeat at 7-10 day intervals through season.

Flax: powdery mildew. Apply 20-30 lbs. per acre. Begin at first sign of disease. Repeat at 7 to 10 day intervals or as necessary.

Hops: mite suppression. Apply 30-40 lbs. per acre.


Peanuts: leaf spot, powdery mildew. Apply 20 to 25 lbs. per acre. Begin when plants are 40 days old or when first spots appear on lower leaves. Repeat at 10 to 14 day intervals and after rains. For rust, apply 20 to 30 lbs. per acre. Apply when disease threatens. Repeat at 7 to 10 day intervals as necessary.

Soybeans: leaf spot, powdery mildew. Apply 8-15 lbs. per acre. Apply at first signs of disease. Repeat at 7-10 day intervals.

Vetch: rust. Apply 25-40 lbs. per acre. Begin at first signs of disease. Repeat at 7-10 day intervals.

ORNAMENTAL CROP USE

Alder, Azalea, Buttonbush, Catalpa, Chrysanthemums, Gladiolus, Honeysuckle, Roses, Sunflowers, Sycamore, Zinnias: powdery mildew. Apply 20-40 lbs. per acre. Begin when disease appears and repeat at 5 to 10 day intervals and after rain or as necessary.



(19:9/74)

OCCIDENTAL CHEMICAL COMPANY

A DIVISION OF OCCIDENTAL PETROLEUM CORPORATION

P. O. BOX 198 • LATHROP, CALIFORNIA 95330 • (209) 858-2511

CONDITIONS OF SALE: 1. Seller warrants that this product consists of the ingredients specified and is reasonably fit for use as directed on the label. No one, other than an officer of Seller, is authorized to make any other warranty, guarantee or direction concerning this product. 2. Because the time, place, rate of application and other conditions of use are beyond Seller's control, Seller's liability from the handling, storage and use of this product is limited to replacement of product or refund of purchase price.