

PRECAUTIONS: Do not use Simazine 80 WP on Bermudagrass in New Mexico and the High Plains of Texas. Do not apply within 7 days following cutting. Do not graze dairy or beef cattle or sheep for 30 days following 1-3 lbs. per acre application or 60 days following 4 lbs. application. Do not cut for hay 60 days following 1-3 lbs. per acre application or 90 days following 4 lbs. application.

Corn: For ground applications use a minimum of 20 gals. of water per acre. Aerial applications may be made using the broadcast rates listed below and 1 gallon of water for each 1 lb. to 1 1/2 lbs. of wettable powder to be applied per acre. Nitrogen solutions or complete liquid fertilizers may replace all or part of the water as a carrier for Simazine 80 WP. Do not apply after corn has emerged as there is danger of liquid fertilizers causing crop injury.

Pre-emergence: Apply before weeds and corn emerge. Use rates listed below.

Preplant: Broadcast in the spring after plowing at the broadcast rates indicated in the table below. Application may be made before, during, or after final seedbed preparation. If soil is tilled or worked after application, avoid deep incorporation of Simazine 80 WP. Best results have been obtained when it is applied within 2 weeks prior to planting.

NOTE: Under dry weather conditions, preplant applications may give better weed control.

SOIL TYPE	RATE PER ACRE	
	BROADCAST	13 INCH BAND ON 40-INCH ROWS*
Light soils Sands, silts, and loams that are low in organic matter	2 1/2 lbs.	13 ozs.
Medium soils Soils containing moderate amounts of clay and organic matter	3 lbs.	1 lb.
Heavy soils Loams that are high in organic matter and clays (including dark prairie soils of Corn Belt)	3 1/2 lbs.	1 1/4 lbs.
Organic soils Peats, mucks, and high organic clays	5 lbs.	1 lb. 10 ozs.

*For other band or row widths adjust the rate in proportion to area actually treated.

To control quackgrass: Broadcast 3 3/4-5 lbs. per acre in the fall. Plow two to three weeks later or if erosion is a problem, delay plowing until spring. Do not plant any crop except corn in the spring following treatment. Do not graze treated area.

NOTE: Should weeds develop, particularly under dry or minimal moisture conditions, a shallow cultivation will generally result in better weed control.

Precautions: Do not apply more than 5 lbs. of Simazine 80 WP to corn in any one year. Land treated with it should not be planted to any crop except corn until the following year or injury may occur. Following harvest of a treated crop, plow and thoroughly till the soil in fall or spring to minimize possible injury to rotational, spring-seeded crops, regardless of the rate used. For any reason, Simazine 80 WP is used at broadcast rate higher than 3 3/4 lbs. per acre (or comparable rates in a band application), a crop of untreated corn should precede the next rotational crop. Do not apply Simazine 80 WP preplant incorporated for weed control in corn in the High Plains and Intermountain areas of the West (including central and western Kansas, western Nebraska, western Oklahoma, and the Panhandle of Texas) where rainfall is sparse and erratic or where irrigation is required. In the High Plains and Intermountain areas of the West where rainfall is sparse and erratic or where irrigation is required, use Simazine 80 WP to control weeds in corn only when corn is to follow corn or a crop of untreated corn is to precede other rotational crops. In western Minnesota and eastern parts of the Dakotas, Nebraska, and Kansas, corn treated with this product should not be followed by Soybeans if the broadcast rate applied was more than 2 1/2 lbs. per acre (or comparable rate in a band) or injury may occur. Do not plant sugar beets, tobacco, vegetables (including dry beans), spring-seeded small grains, or small-seeded legumes and grasses the year following Simazine 80 WP application or injury may occur.

Grasses Grown for Seed (Pacific Northwest Only): Perennial Ryegrass, Highland or Astoria Bentgrasses, Orchardgrass, Tall Fescue and Fine Fescues such as Illahep, Ranier, Pennlawn, Chewings and related species. For the control of broadleaf weeds and annual grasses such as annual ryegrass, rattail fescue, silver hairgrass and downy brome (cheat), apply 2 1/2 lbs. of Simazine 80 WP broadcast in 15 or more gallons of water per acre as soon as fall rains start. Early application is important. Use a spray rig having mechanical agitation.

Precautions: Apply only to grasses from which at least one seed crop has been cut. Allow 30 days between last application and grazing to dairy and beef cattle and sheep, and 60 days between the last application and cutting for hay.

NURSERIES, CHRISTMAS TREE PLANTINGS, SHELTER BELTS

Nurseries: (see list below) Broadcast 2 1/2-3 1/2 lbs. in at least 25 gallons of water per acre in fall or spring to nursery stock established at least one year after transplanting.

Christmas Tree Plantings and Shelter Belts: (see list below) Broadcast 2 1/2-5 lbs. in at least 25 gals. of water per acre after transplanting. For band applications reduce the rate of Simazine 80 WP in proportion to the area treated. For example: Apply at the rate of 1 1/2 lbs. per acre on heavy soil type in a 4-foot band one mile long or 1 1/2 oz. per 1,000 sq. ft. (equivalent to 2 1/2 lbs. broadcast per acre). Remove weed growth before application. Use the same rate for annual maintenance applications.

For quackgrass control: broadcast 5 lbs. per acre in the fall or make a split application (apply 2 1/2 lbs. per acre in the fall and 2 1/2 lbs. per acre in early spring after quackgrass begins growth).

Precautions: Do not use Simazine 80 WP on seedbeds or cutting beds. Do not apply to Christmas tree shelter belt transplants less than three years of age. Do not apply it more than once a year, even if it is directed for quackgrass control.

Apply Simazine 80 WP to these species of trees and shrubs as recommended above: American Elm, Austrian Pine, Arborvitae, Balsam Fir, Barberry, Blue Spruce, Boxelder, Bush Honeysuckle, Caragana, Colorado Blue Spruce, Douglas Fir, Fraser Fir, Hemlock, Honey Locust, Juniper, Mugo Pine, Norway Spruce, Oregon Grape (Mahonia spp.), Red Cedar, Red Oak, Red Pine, Norway Pine, Red Spruce, Russian Olive, Scotch Pine, Siberian Elm, White Cedar, White Pine, White Spruce, Yew, Taxus spp.

TURF GRASSES FOR SOD

St. Augustine, Centipede, and Zoysia Grass: Broadcast 2 1/2-5 lbs. per acre in spring or fall as indicated below. Apply in sufficient water for thorough coverage.

Muck or peat: 5 lbs. *Old Beds* Within 2 days after lifting of sod
New Beds 3-4 days after sprigging or plugging

Sandy soils: 2 1/2 lbs. *Old Beds* Within 2 days after lifting of sod
New Beds 7-10 days after sprigging or plugging

Apply an additional 2 1/2 lbs. on muck or peat, or 1 1/2 lbs. on sandy soils if weed growth recurs.

Precautions: Do not apply within 30 days before cutting or lifting. Do not use north of 37° latitude.

VEGETABLE CROPS

Asparagus (Established): Broadcast 2 1/2-5 lbs. per acre to beds established one year or more. Make one application per year either after disking in the spring and before weeds emerge, or following harvest after removal of existing weed growth. Spring application must be made at least 3 days before first cutting. Precautions for Washington and Oregon only. Do not use on sand or loamy sand soils or gravelly areas of fields as crop injury may result. Do not allow irrigation water to stand or to form pools in the field or injury may occur in these areas.

Artichokes: Broadcast 2 1/2-5 lbs. per acre after last fall tillage. Apply as a directed spray.

Precautions: Do not apply to sandy soils. Do not make more than one application per year.

NONSELECTIVE WEED CONTROL ON NONCROP LAND

Apply Simazine 80 WP at rates recommended below. Use the higher rates on heavy clay and muck soils. Best results are obtained when Simazine 80 WP is applied before weeds emerge. If weed growth has begun and removal is not practicable, include a contact or translocated herbicide in the spray. Do not contaminate domestic irrigation water supplies, or lakes, streams, or ponds.

It can be used to provide long-term weed control on industrial sites, highway medians and shoulders, railroad rights of way, lumberyards, petroleum tank farms, and in non-crop areas on farms, such as around buildings, equipment and fuel storage areas, along fences, roadsides, and lanes. Aerial applications may be made where it is practicable.

Use sufficient water to assure thorough coverage. Use at least 1 gal. of water for each 1 lb. of Simazine 80 WP, more if practical for both ground and aerial applications. Mechanical, by-pass, or jet agitation is necessary to keep this product in suspension during application.

To control most annual and many perennial broadleaf and grassy weeds (including quackgrass, bluegrass, redtop, burdock, Canada thistle, orchardgrass, purple top, smooth brome, dog fennel and plantain): Broadcast 12 1/2-25 lbs. per acre.


To control hard to kill perennial weeds (such as bull thistle and sow thistle): Broadcast 25-50 lbs. per acre. For longer residual control in regions of high rainfall and a long growing season, broadcast 25-50 lbs. per acre.

For small areas, 4 1/2 oz. per 1,000 sq. ft. is equivalent to 1 1/2 lbs. per acre.

OCCIDENTAL CHEMICAL COMPANY

P. O. BOX 198 - LATHROP, CALIFORNIA 95130 - PHONE (209) 868-2511

DISCLAIMER OF WARRANTIES: Seller warrants that the chemical composition of this product conforms to the chemical description given on this label. THIS WARRANTY IS EXPRESSLY IN LIEU OF ALL OTHER WARRANTIES AND REPRESENTATIONS, EXPRESSED, IMPLIED OR STATUTORY, INCLUDING WARRANTIES OF MERCHANTABILITY AND FITNESS FOR A PARTICULAR USE. Timing, rate and method of application, weather and crop conditions, mixtures with chemicals not specifically recommended on this label or an accompanying written recommendation, are beyond the control of seller. Buyer assumes all risks of use, storage and handling of this material not in strict accordance with directions given herewith. Buyer further agrees in the event of damages arising from the use of this product to accept a replacement of the product or a refund of the purchase price of the product at buyer's option as full discharge of seller's liability. No one is authorized to make any other warranty, guarantee or directions concerning this product and no such warranties, guarantees or directions shall be valid or binding upon seller.



PRECAUTIONS: Do not use Simazine 80 WP on Bermudagrass in New Mexico and the High Plains of Texas. Do not apply within 7 days following cutting. Do not graze dairy or beef cattle or sheep for 30 days following 1.3 lbs. per acre application or 60 days following 4 lbs. application. Do not cut for hay 60 days following 1.3 lbs. per acre application or 90 days following 4 lbs. application.

Corn: For ground applications use a minimum of 20 gals. of water per acre. Aerial applications may be made using the broadcast rates listed below and 1 gallon of water for each 1 lb. to 1.5 lbs. of wettable powder to be applied per acre. Nitrogen solutions or complete liquid fertilizers may replace all or part of the water as a carrier for Simazine 80 WP. Do not apply after corn has emerged as there is danger of liquid fertilizers causing crop injury.

Pre-emergence: Apply before weeds and corn emerge. Use rates listed below.

Preplant: Broadcast in the spring after plowing at the broadcast rates indicated in the table below. Application may be made before, during, or after final seedbed preparation. If soil is tilled or worked after application, avoid deep incorporation of Simazine 80 WP. Best results have been obtained when it is applied within 2 weeks prior to planting.

NOTE: Under dry weather conditions preplant applications may give better weed control.

SOIL TYPE	RATE PER ACRE	
	BROADCAST	13 INCH BAND ON 40 INCH ROWS*
Light soils Sands, silts, and loams that are low in organic matter	2.5 lbs.	1.5 lbs.
Medium soils Soils containing moderate amounts of clay and organic matter	3 lbs.	1.5 lbs.
Heavy soils Loams that are high in organic matter and clays (including dark prairie soils of Corn Belt)	3.5 lbs.	1.5 lbs.
Organic soils Peats, mucks, and high organic clays	5 lbs.	1 lb. 10 lbs.

*For other band or row widths adjust the rate in proportion to area actually treated.

To control quackgrass: Broadcast 3-4.5 lbs. per acre in the fall. Plow two to three weeks later or if erosion is a problem, delay plow until spring. Do not plant any crop except corn in the spring following treatment. Do not graze treated area.

NOTE: Should weeds develop, particularly under dry or minimal moisture conditions, a shallow cultivation will generally result in better weed control.

Precautions: Do not apply more than 5 lbs. of Simazine 80 WP to corn in any one year. Land treated with it should not be planted to any crop except corn until the following year or injury may occur. Following harvest of a treated crop, plow and thoroughly till the soil in fall or spring to minimize possible injury to rotational spring seeded crops. Regrowth of the treated area may be delayed. In the West, use of broadcast rate higher than 3-4 lbs. per acre (or comparable rates in a band application) a crop of untreated corn should precede the next rotational crop. Do not apply Simazine 80 WP preplant incorporated for weed control in corn in the High Plains and Intermountain areas of the West (including central and western Kansas, western Nebraska, western Oklahoma, and the Panhandle of Texas) where rainfall is sparse and erratic or where irrigation is required. In the High Plains and Intermountain areas of the West where rainfall is sparse and erratic or where irrigation is required, use Simazine 80 WP to control weeds in corn only when corn is to follow corn or a crop of untreated corn is to precede other rotational crops. In western Minnesota and eastern parts of the Dakotas, Nebraska, and Kansas, corn treated with this product should not be followed by soybeans if the broadcast rate applied was more than 2.5 lbs. per acre (or comparable rate in a band) or injury may occur. Do not plant sugar beets, tobacco, vegetables (including dry beans), spring-seeded small grains or small-seeded legumes and grasses the year following Simazine 80 WP application or injury may occur.

Grasses Green for Seed: Pacific Northwest Only: Perennial Ryegrass, Highland or Astoria Bentgrasses, Orchardgrass, Tall Fescue and Fine Fescues such as Illahee, Ramier, Pentstemon, Chewings and related species. For the control of broadleaf weeds and annual grasses such as annual ryegrass, rattail fescue, silver hairgrass and downy brome (cheat), apply 2.5 lbs. of Simazine 80 WP broadcast in 15 or more gallons of water per acre as soon as fall rains start. Early application is important. Use a spraying having mechanical agitation.

Precautions: Apply only to grasses from which at least one seed crop has been cut. Allow 30 days between last application and grazing to dairy and beef cattle and sheep, and 60 days between the last application and cutting for hay.

NURSERIES, CHRISTMAS TREE PLANTINGS, SHELTER BELTS

Nurseries (see list below): Broadcast 2.5-3 lbs. in at least 25 gallons of water per acre in fall or spring to nursery stock established at least one year after transplanting.

Christmas Tree Plantings and Shelter Belts (see list below): Broadcast 2.5-5 lbs. in at least 25 gallons of water per acre after transplanting. For hand applications reduce the rate of Simazine 80 WP and volume of water in proportion to the area treated. For example, apply at the rate of 1.5-2 lbs. according to your soil type in a 4-foot band one mile long or 1.2 oz. per 1,000 sq. ft. (equivalent to 2.5 lbs. broadcast per acre). Remove weed growth before application. Use the same rate for annual maintenance applications.

For quackgrass control: broadcast 5 lbs. per acre in the fall or make a split application applying 2.5 lbs. per acre in the fall and 2.5 lbs. per acre in early spring after quackgrass begins growth.

Precautions: Do not use Simazine 80 WP on seedbeds or cutting beds. Do not apply to Christmas tree or shelter belt transplants less than three years of age. Do not apply it more than once a year, except as directed for quackgrass control.

Apply Simazine 80 WP to these species of trees and shrubs as recommended above: American Elm, Austrian Pine, Arborvitae, Balsam Fir, Barberry, Blue Spruce, Boxelder, Bush Honeysuckle, Caragana, Cotton-wool, Dogwood, Douglas Fir, Fraser Fir, Hemlock, Honey Locust, Juniper, Mugo Pine, Norway Spruce, Oregon Grape, Mahonia spp., Red Cedar, Red Oak, Red Pine, Norway Pine, Red Spruce, Russian Olive, Scotch Pine, Siberian Elm, White Cedar, White Pine, White Spruce, Yew (Taxus spp.).

TURF GRASSES FOR SOD

St. Augustine, Centipede, and Zoysia Grass: Broadcast 2.5 lbs. per acre according to soil type as indicated below. Apply in sufficient water for thorough coverage.

Muck or peat: 5 lbs. *Old Beds* Within 2 days after lifting of sod
New Beds 3-4 days after springing or plugging

Sandy soils: 2.5 lbs. *Old Beds* Within 2 days after lifting of sod
New Beds 7-10 days after springing or plugging

Apply an additional 2.5 lbs. on muck or peat or 1.5 lbs. on sandy soils if weed growth recurs.

Precautions: Do not apply within 30 days before cutting or lifting. Do not use north of North Carolina.

VEGETABLE CROPS

Asparagus (Established): Broadcast 2.5-5 lbs. per acre to beds established one year or more. Make one application per year either after disking in the spring and before weeds emerge, or following harvest after removal of existing weed growth. Spring application must be made at least 3 days before first cutting. Precautions for Washington and Oregon only. Do not use on sand or loamy sand soils or gravelly areas of fields as crop injury may result. Do not allow irrigation water to stand or to form pools in the field or injury may occur in these areas.

Artichokes: Broadcast 2.5 lbs. per acre after last fall tillage. Apply as a directed spray.

Precautions: Do not apply to sandy soils. Do not make more than one application per year.

NONSELECTIVE WEED CONTROL ON NONCROP LAND

Apply Simazine 80 WP at rates recommended below. Use the higher rates on heavy clay and muck soils. Best results are obtained when Simazine 80 WP is applied before weeds emerge. If weed growth has begun and removal is not practicable, include a contact or translocated herbicide in the spray. Do not contaminate domestic or irrigation water supplies, or lakes, streams, or ponds.

It can be used to provide long term weed control on industrial sites, highway medians and shoulders, railroad rights of way, lumberyards, petroleum tank farms, and in non crop areas on farms, such as around buildings, equipment and fuel storage areas, along fences, roadsides, and lanes. Aerial applications may be made where it is practicable.

Use sufficient water to assure thorough coverage. Use at least 1 gal. of water for each 1 lb. of Simazine 80 WP, more if practical for both ground and aerial applications. Mechanical, by pass, or jet agitation is necessary to keep this product in suspension during application.

To control most annual and many perennial broadleaf and grass weeds (including quackgrass, bluegrass, redtop, burdock, Canada thistle, orchardgrass, purple top, smooth brome, dog fennel and plantain): Broadcast 12.5-25 lbs. per acre.

To control hard to kill perennial weeds (such as bull thistle and sow thistle): Broadcast 25-50 lbs. per acre. For longer residual control in regions of high rainfall and a long growing season, broadcast 25-50 lbs. per acre.

For small areas 4.5 oz. per 1,000 sq. ft. is equivalent to 12.5 lbs. per acre.

OCCIDENTAL CHEMICAL COMPANY

P. O. BOX 110 - LATHROP, CALIFORNIA 95230 - PHONE (209) 866-2511

DISCLAIMER OF WARRANTIES: Seller warrants that the chemical composition of this product conforms to the chemical description given on the label. THIS WARRANTY IS EXPRESSLY IN LIEU OF ALL OTHER WARRANTIES AND REPRESENTATIONS, EXPRESSED, IMPLIED OR STATUTORY, INCLUDING WARRANTIES OF MERCHANTABILITY AND FITNESS FOR A PARTICULAR USE. Timing, rate and method of application, weather and crop conditions, mixtures with chemicals not specifically recommended on this label or an accompanying written recommendation, are beyond the control of seller. Buyer assumes all risks of use, storage and handling of this material not in strict accordance with directions given hereon. Buyer further agrees in the event of damages arising from the use of this product to accept a replacement of the product or a refund of the purchase price of the product, at buyer's option, as full discharge of seller's liability. No one is authorized to make any other warranty, guarantee or directions concerning this product. Such warranties, guarantees or directions shall be valid or binding upon seller.

(68)

BEST DOCUMENT AVAILABLE

NURSERIES, CHRISTMAS TREE PLANTINGS SHELTER BELTS

Nurseries (see list below): Broadcast 2 1/2 - 3 1/2 lbs. in at least 25 gallons of water per acre in fall or spring to nursery stock established at least one year after transplanting.

Christmas Tree Plantings and Shelter Belts (see list below): Broadcast 2 - 5 lbs. in at least 25 gallons of water per acre after transplanting. For hand applications reduce the rate of Simazine 80 WP and volume of water in proportion to the area treated. For example: Apply at the rate of 1 - 2 1/2 lbs. according to your soil type in a 4 foot band one mile long or 1/2 oz. per 1 000 sq. ft. (equivalent to 2 1/2 - 5 lbs. broadcast per acre). Remove weed growth before application. Use the same rate for annual maintenance applications.

For quackgrass control broadcast 5 lbs. per acre in the fall or make a split application applying 2 1/2 lbs. per acre in the fall and 2 1/2 lbs. per acre in early spring after quackgrass begins growth.

Precautions: Do not use Simazine 80 WP on seedbeds or cutting beds. Do not apply to Christmas tree or shelter belt transplants less than three years of age. Do not apply if more than once a year except as directed for quackgrass control.

Apply Simazine 80 WP to these species of trees and shrubs as recommended above: American Elm, Austrian Pine, Arborvitae, Balsam Fir, Barberry, Blue Spruce, Boxelder, Bush Honeysuckle, Caragana, Cotton-aster, Dogwood, Douglas Fir, Fraser Fir, Hemlock, Honey Locust, Juniper, Mugho Pine, Norway Spruce, Oregon Grape (Mahonia spp.), Red Cedar, Red Oak, Red Pine (Norway Pine), Red Spruce, Russian Olive, Scotch Pine, Siberian Elm, White Cedar, White Pine, White Spruce, Yew (Taxus spp.).

TURF GRASSES FOR SOD

St. Augustine, Centipede, and Zoysia Grass: Broadcast 2 1/2 - 5 lbs. per acre according to soil type as indicated below. Apply in sufficient water for thorough coverage.

Muck or peat 5 lbs. *Old Beds* Within 2 days after lifting of sod
New Beds 3 - 4 days after sprigging or plugging

Sandy soils 2 - 3 lbs. *Old Beds* Within 2 days after lifting of sod
New Beds 7 - 10 days after sprigging or plugging

Apply an additional 2 1/2 lbs. on muck or peat or 1 1/2 lbs. on sandy soils if weed growth recurs.

Precautions: Do not apply within 30 days before cutting or lifting. Do not use north of North Carolina.

VEGETABLE CROPS

Asparagus (Established): Broadcast 2 - 5 lbs. per acre to beds established one year or more. Make one application per year either after disking in the spring and before weeds emerge or following harvest after removal of existing weed growth. Spring application must be made at least 3 days before first cutting. Precautions for Washington and Oregon only. Do not use on sand or loamy sand soils or gravelly areas of fields as crop injury may result. Do not allow irrigation water to stand or to form pools in the field or injury may occur in these areas.

Artichokes: Broadcast 2 - 5 lbs. per acre after last fall tillage. Apply as a directed spray.

Precautions: Do not apply to sandy soils. Do not make more than one application per year.

NONSELECTIVE WEED CONTROL ON NONCROP LAND

Apply Simazine 80 WP at rates recommended below. Use the higher rates on heavy clay and muck soils. Best results are obtained when Simazine 80 WP is applied before weeds emerge. If weed growth has begun and removal is not practicable, include a contact or translocated herbicide in the spray.

Do not contaminate domestic or irrigation water supplies or lakes, streams or ponds.

It can be used to provide long term weed control on industrial sites, highway medians and shoulders, railroad rights of way, lumberyards, petroleum tank farms, and in non crop areas on farms, such as around buildings, equipment and fuel storage areas, along fences, roadsides, and lanes. Aerial applications may be made where it is practicable.

Use sufficient water to assure thorough coverage. Use at least 1 gal. of water for each 1 lb. of Simazine 80 WP, more if practical for both ground and aerial applications. Mechanical by pass or jet agitation is necessary to keep this product in suspension during application.

To control most annual and many perennial broadleaf and grass weeds (including quackgrass, bluegrass, redtop, burdock, Canada thistle, orchardgrass, purple top, smooth brome, dog fennel and plantain): Broadcast 12 1/2 - 25 lbs. per acre.

To control hard to kill perennial weeds (such as bull thistle and sow thistle): Broadcast 25 - 50 lbs. per acre. For longer residual control in regions of high rainfall and a long growing season, broadcast 25 - 50 lbs. per acre.

For small areas, 4 - 6 oz. per 1 000 sq. ft. is equivalent to 12 1/2 lbs. per acre.

OCCIDENTAL CHEMICAL COMPANY

P. O. BOX 198 - LATHROP, CALIFORNIA 95128 - PHONE (209) 866-2611

DISCLAIMER OF WARRANTIES: Seller warrants that the chemical composition of this product conforms to the chemical description given on this label. THIS WARRANTY IS EXPRESSLY IN LIEU OF ALL OTHER WARRANTIES AND REPRESENTATIONS, EXPRESSED, IMPLIED OR STATUTORY, INCLUDING WARRANTIES OF MERCHANTABILITY AND FITNESS FOR A PARTICULAR USE. Timing, rate and method of application, weather conditions, mixtures with chemicals not specifically recommended on this label, or an assumption of the recommendation, are beyond the control of seller. Buyer assumes all risks of use, storage and handling of this material not in strict accordance with directions given hereon. Buyer further agrees, in the event of any claim arising from the use of this product, to accept a replacement of the product or a refund of the purchase price of the product, at buyer's option, as full discharge of seller's liability. No one is authorized to make any warranty, guarantee or directions concerning this product, and no such warranties, guarantees or directions shall be valid or binding upon seller.

(68)

(69)

