


# DIAZINON 5 GRANULES

<b>ACTIVE INGREDIENT:</b>		<b>BY WT.</b>
O,O-diethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate		5.0%
<b>INERT INGREDIENTS:</b>		95.0%
	<b>TOTAL</b>	<b>100.0%</b>

EPA Reg. No. 7001-197-AA  
EPA Est. No. 7001-CA-1

Diazinon® trademark of Ciba-Geigy Corp.



## CAUTION

### KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed. Avoid contamination of food and feed. Do not store near food or feed products. Avoid inhalation of dust. Avoid contact with skin, eyes and clothing. Wash thoroughly after handling. May be absorbed through skin.

This product is toxic to fish, birds, and other wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, streams or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

Do not reuse container. Destroy it by burying or burning. Avoid smoke or fumes.

### DIRECTIONS FOR USE

**LAWN INSECTS:** *Lawn Chinch Bugs*. Apply 1 1/3 to 4 lbs. per 1000 sq. ft. Application should be repeated in 7 to 10 days if necessary.

*Chinch Bugs, Earwigs, Ticks, Ants, Lawn Moths (Sod Webworms), Bermuda Mites, Crickets, Armyworms, Brown Dog Ticks, Chiggers, Clover Mites, Cutworms, Fleas, Leafhoppers, Sowbugs, Millipedes, Springtails (Collembola)*. Apply Diazinon® 5 Granules to your lawn with a fertilizer spreader. Application should be made when grass is dry. Water lawn lightly after application. Repeat as necessary.

For application with other types of equipment, or by hand, apply at the rate of 2 lbs. per 1000 sq. ft. (an area 20' by 50') or 3.2 ounce per 100 sq. ft. (an area 10' by 10'). For heavy lawns, such as St. Augustine lawns, application should be increased to 3 lbs. per 1000 sq. ft. (or 4.8 ounce per 100 sq. ft.).

*Bill Bugs and White Grubs of Japanese Beetle, European Chafer, Southern Chafer.* Apply 2.5 lbs. per 1000 sq. ft. of lawn. Water grass thoroughly after application. For grub control make application anytime between late July and early October. For *bill bug* control, apply when activity is first observed or when chewed or brown grass indicated damage from these insects. Repeat as necessary.

#### SPOT TREATMENT

*Ants:* Apply 1½ teaspoons of Diazinon® 5 Granules over and around each individual ant hill.

*Digger Wasps:* Apply 1½ teaspoons of Diazinon® 5 Granules in and around each individual ground nest opening. Make the application about dusk when the wasps are not active.

#### BAND TREATMENT AROUND FOUNDATION

The following insects are commonly found around or near the foundation of houses as well as in the lawn. Certain of these insects may even enter the house. *Ants, Brown Dog Ticks, Clover Mites, Crickets, Earwigs, Fleas, Millipedes, Sowbugs, Springtails (Collembola).* For control of these insects, apply a five foot band of Diazinon® 5 Granules next to the foundation of the house. Distribute the granules uniformly using a rate of 3.2 ounces for each 5' by 20' area to be treated.

(GL-10:8/76)

# NET WEIGHT 35 POUNDS