L'YPREX 4 DUST

Read This Label Carefully

ACTIVE INGREDIENT:	BY WT.
Dodine (Dodecylguanidine acetate)	4%
NERT INGREDIENTS:	96%
TOTAL	100 %

EPA Reg. No. 7001-205-AA EPA Est. No. 7001-CA-1

You Get Results With

WARNING

KEEP OUT OF REACH OF CHILDREN

Causes eye burn, harmful if swallowed. Do not get in eyes or on skin. Wear gloves and goggles when handling. Avoid breathing dust. Keep container closed. Use with adequate ventilation. Wash thoroughly after handling. Do not contaminate food or feed products.

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning equipment or disposal of wastes. Apply this product only as specified on this label.

Destroy empty bags by burning. Keep out of smoke.

Che

firs' nuriii

Pears (Western states only): Scab. Apply 50 lbs per acre Begin in pre-bloom stages. Repeat at 5 to 7 days through first cover spray. Do not apply within 5 days of harvest. Do not graze cover crop in treated orchards. Strawberries: Leaf Blight, Leaf Scorch, Leaf Spot. Apply 25 to 30 lbs/per acre. Begin

within 14 days of harvest.

Roses: Black Spot. Apply 8 to 10 lbs. per acre. Make first application when young leaves first appear. Repeat at 7 to 10 day intervals throughout season. Caution: Higher dosage may cause stunting and defoliation on first year plants and chlorosis on older established plants

OCCIDENTAL CHEMICAL COMPANY

• LATHROP, CALIFORNIA 95330 • PHONE (209) 858-2511 P. O. BOX 198


CONDITIONS OF SALE: 1. Seller warrants that this product consists of the ingredients specified and is reasonably fit for use as directed on the label. No one, other than an officer of Seller, is authorized to make any other warranty, guarantee or direction concerning this product. 2. Because the time, place, rate of application and other conditions of use are beyond Seller's control, Seller's liability from the handling, storage and use of this product is limited to replacement of product or refund of purchase price.

DIRECTIONS FOR USE

Apples: Scab. Apply 20 to 40 lbs per acre Apply in late cover application or as necessary. Note: May cause russeting on golden delicious and grimes golden. Do not apply within 5 days of harvest. Do not graze cover crop in treated orchards. Do not use treated apple pomace for use in livestock feeds

'.eaf Spot. Apply 40 to 60 lbs. per acre. Apply at petal fall or when leaves Repeat at 7 to 10 day intervals. Do not apply when temperature and Lie excessively high.

as soon as new growth starts. Repeat at weekly intervals as necessary. Do not apply



(737:9,75)