



DIBROM 4 DUST

ACCEPTED

MAR 22 1976

Under the Federal Insecticide, Fungicide, and Rodenticide Act, 1910
EPA Reg. 950.00000000

ACTIVE INGREDIENT:

Naled* 4

INERT INGREDIENTS:

TOTAL 100%

* Dibrom is Chevron Chemical Company's registered trademark for Naled (1,2-dibromo-2,2-dichloroethyl dimethyl phosphate)
U.S. Patent No. 2,971,882

EPA Reg. No. 7001-219-AA

EPA Est. No. 7001-CA-1

WARNING

KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed. May cause irritation of eyes, nose and throat. Avoid breathing dust. Avoid contact with skin and eyes. Wash thoroughly after using. Do not store or transport near feed or food.

Do not use on cherries as injury may result.

May corrode metal dust equipment. Clean equipment after use.

This product is toxic to fish, shrimp, crab, and other wildlife. Keep out of lakes, streams, ponds, bays, marshes and estuaries. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

Use within 60 days of manufacture only, as this product decomposes.

DATE OF MANUFACTURE

Do not allow this product to drift onto adjoining food or feed crops as residues may contaminate these crops making them unfit for sale, feed or human consumption. Keep clear of food or dairy animals, streams, fish ponds and wildlife areas.

STORE IN COOL, DRY PLACE!

BURN BAG IMMEDIATELY WHEN EMPTY

STAY OUT OF SMOKE!

DO NOT APPLY TO SULFUR SENSITIVE CROPS OR VARIETIES

DO NOT APPLY DURING PERIODS OF EXTREME HEAT

DIRECTIONS FOR USE

This product is formulated for application by ground or air equipment. For airplane application, use only the dosages. Begin applications at first sign of insects. Repeat as necessary. Apply for contact and thorough coverage to obtain control of insects.

Adult Mosquito Control: Corral, feed lots, pens. Apply 8 to 10 lbs. per acre. Apply in ground fast. Repeat as needed. Avoid farm buildings including dairy barns. Do not pasture animals within 24 hours.

Beans: Apply 30 to 50 lbs. per acre. Do not apply within 4 days of harvest.

Broccoli, Brussels Sprouts, Cabbage, Cauliflower, Collards, Kale: Apply 30 to 50 lbs. per acre. Do not apply within 1 day of harvest on collards, brussels sprouts, cabbage and cauliflower. Do not apply to collards and kale within 4 days of harvest.

Citrus (Grapefruit, Lemons, Oranges, Tangerines): Apply 30 to 50 lbs. per acre. Do not apply within 1 day of harvest in Arizona and California. Within 7 days of harvest in other areas.

Cotton: Apply 30 to 50 lbs. per acre. Do not apply within 4 days of harvest.

Cucumbers, Summer Squash: Apply 30 to 50 lbs. per acre. May be used up to day of harvest.

Eggplant: Apply 30 to 50 lbs. per acre. Do not apply within 1 day of harvest.

Grapes: Apply 30 to 50 lbs. per acre. Do not apply within 4 days of harvest.

Hops: Apply 30 to 50 lbs. per acre. Do not apply within 4 days of harvest.

Lettuce, Spinach, Celery, Chard, Turnip Greens: Apply 30 to 50 lbs. per acre. Do not apply within 1 day of harvest on lettuce, spinach, chard and turnip greens. Do not apply to celery within 4 days of harvest.

Melons (Cantaloupes, Honeydews, Muskmelons, Watermelons), Pumpkins, Winter Squash: Apply 30 to 50 lbs. per acre. Do not apply within 1 day of harvest.

Ornamentals: Apply 30 to 50 lbs. per acre.

Peaches: Apply 30 to 50 lbs. per acre. Do not apply within 4 days of harvest. Restricted to the Western United States.

Peas: Apply 30 to 50 lbs. per acre. Do not apply within 4 days of harvest.

Peppers: Apply 30 to 50 lbs. per acre. Do not apply within 1 day of harvest.

Roses (Field Grown): Apply 30 to 50 lbs. per acre.

Strawberries: Apply 30 to 50 lbs. per acre. Do not apply within 1 day of harvest.

Tomatoes (Field): Apply 30 to 50 lbs. per acre. Do not apply within 1 day of harvest.

Walnuts: Apply 30 to 50 lbs. per acre. Do not apply within 10 days of harvest.

(20:176)

Occidental Chemical Company

P.O. Box 198 Lathrop, California 95330



DIBROM 4 DUST

ACTIVE INGREDIENT:

Naled*.....

BY WT.

4%

INERT INGREDIENTS:

TOTAL 100%

96%

* Dirbom is Chevron Chemical Company's registered trademark for Naled
(1,2-dibromo-2,2-dichloroethyl dimethyl phosphate)

U S Patent No. 2,971,882

EPA Reg. No. 7001-219-AA

EPA Est. No. 7001-CA-1

WARNING

KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed. May cause irritation of eyes, nose and throat. Avoid breathing dust. Avoid contact with skin and eyes. Wash thoroughly after using. Do not store or transport near feed or food.

Do not use on cherries as injury may result.

May corrode metal dust equipment, clean equipment after use.

This product is toxic to fish, shrimp, crab, and other wildlife. Keep out of lakes, streams, ponds, tidal marshes and estuaries. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

Use within 60 days of manufacture only, as this product decomposes

DATE OF MANUFACTURE: _____

Do not allow this product to drift onto adjoining food or feed crops as residues may contaminate these crops making them unfit for sale, feed or human consumption. Keep clear of food or dairy animals, streams, fish ponds and wildlife areas

STORE IN COOL, DRY PLACE!

BURN BAG IMMEDIATELY WHEN EMPTY

STAY OUT OF SMOKE!

DO NOT APPLY TO SULFUR-SENSITIVE CROPS OR VARIETIES

DO NOT APPLY DURING PERIOD OF EXTREME HEAT

DIRECTIONS FOR USE

This product is formulated for application by ground or air equipment. For airplane application, use heavier dosages. Begin applications at first sign of insects. Repeat as necessary. Apply for contact and thorough coverage to obtain control of insects.

Adult Mosquito Control: Corrals, feed lots, pens. Apply 8 to 10 lbs. per acre. Aerial or ground application. Repeat as needed. Avoid farm buildings including dairy barns. Do not pasture animals on treated areas.