

*U.S. Patent No 2.873.228

EPA Reg No 7001-223-AA EPA Est No 7001-CA-1

CAUTION

KEEP OUT OF REACH OF CHILDREN

DO NOT BREATHE DUST

DO NOT GET ON SKIN, IN EYES, OR ON CLOTHING
MAY BE ABSORBED THROUGH SKIN
WASH THOROUGHLY AFTER HANDLING

DO NOT STORE NEAR FOOD AND FEED PRODUCTS

This product is toxic to fish shrimp, crabs and other wildlife. Keep out of lakes streams ponds tidal marshes and estuaries. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where these are important resources. Apply this product only as specified on this label.

This product is toxic to bees and should not be applied when bees are actively visiting the area. Do not reuse empty container. Destroy it by burying with waste

DIRECTIONS FOR USE

400 To be applied by ground application or aircraft.

Almonds: Brown Mite, European Red Mite, Two-spotted Spider Mite. Apply 30 to 50 lbs. per acre. Do not apply more than once during fruiting period. Do not apply within 30 days of harvest.

Apples: Apple Rust Mite. Brown Mite. Codling Moth. European Red Mite. Two-spotted Spider Mite. Apply 50 to 90 lbs. per acre. For Codling Moth. apply at petal fall and every 10 to 14 days thereafter until control is achieved. Do not apply within 40 tays of harvest if no more than 2 applications are made. Do not use on early McIntosh variety, or varieties ripening before or at the same time as early McIntosh.

Beans: Bean Leaf Beetle, Mexican Bean Beetle, Two-spotted Spider Mite. Apply 15

** 20 lbs per acre. Do not apply within 2 days of harvest. Do not feed treated foliage to vestock. Lygus Bugs. Apply 30 to 40 lbs. per acre. Do not apply within 4 days of harvest. Do not feed treated foliage to livestock.

Cotton: Spider Mites (other than Western United States). Apply 12 lbs per acre Aphids (other than Western United States). Apply 12 to 25 lbs per acre. Plant Bugs including Lygus Bugs and Spider Mites (Western United States only). Apply 25 to 40 lbs per acre. Do not apply after cotton bolls open. Do not graze meat or dairy animals in treated fields. Workers entering treated fields within 24 hours following application should wear protective clothing.

Eggplant: Two-spotted Spider Mite. Apply 6 to 12 lbs per acre May be applied up to day of harvest.

Grapefruit, Oranges, Tangelos, Tangerines: Citrus Rea Mite. Citrus Rust Mite. Texas Citrus Mite. Apply 50 to 100 lbs per acre Citrus Thrips (Arizona & California only) Apply 35 to 100 lbs per acre May be applied up to day of harvest. Do not repeat applications within 90 days. On Tangelos and Tangerines, do not apply more than 2 times per season.


Grapes: Grape Leafhopper. Two-spetted Spider Mite. Williamette Mite (Western United States only) Apply 15 to 30 lbs per acre. Do not apply within 15 days of harvest. Do not make more than 2 applications during fruiting season. Do not apply to Febr. Szago. Muscat or Salvador varieties.

Lemons, Limes: Citrus Red Mite. Citrus per acre Citrus Thrips (£ rizona ai Apply only 1 time per season. Do not Melons: Serpentine Leafminer Compleafhopper. Apply 30 to 40 lbs. per Nectarines: Brown Mite. European apply more than 2 times during the harvest.

Peaches: Brown Mite Europear, Red lbs per acre. Do not suply more than within 30 days of harvest

Peppers, Pimentos: Serpentine Lea lbs. per acre. May be applied up to Plums, Prunes: An Mite. Europea to 60 lbs per acre. Do not apply mor apply within 45 days of harvest

Strawberries: Strawberry Leafroller. Tacre Do not apply within 2 days of Interest Tomatoes: Leafnmers. Tropical Spid per acre. Do not apply within 2 days



Occidental Ch

CONDITIONS OF SALE: I Seiler warrants that the easonably fit for one as directed on the label. No make any other warranty guarantee or direction column to of application and other conditions of use are handing, storage and use of this product is limited to

JIY 4 DU3 I

· BY WT.

DIRECTIONS FOR USE

To be applied by ground application or aircraft

Almonds: Brown Mite. European Red Mite, Two-spotted Spider Mite. Apply 30 to 50 lbs. per acre. Do not apply more than once during fruiting period. Do not apply within 30 days of harvest.

Apples: Apple Rust Mite. Brown Mite. Codling Moth. European Red Mite. Two-spotted Spider Mite. Apply 50 to 90 lbs. per acre. For Codling Moth. apply at petal fall and every 10 to 14 days thereafter until control is achieved. Do not apply within 40 days of harvest if no more than 2 applications are made. Do not use on early Molntosh variety, or varieties repening before or at the same time as early Molntosh. Beans: Bean Leaf Beetle. Mexican Bean Beetle. Two-spotted Spider Mite. Apply 15 to 20 lbs. per acre. Do not apply within 2 days of harvest. Do not feed treated foliage to vestock. Lygus Bugs. Apply 30 to 40 lbs. per acre. Do not apply within 4 days of harvest. Do not feed treated foliage to livestock.

Cotton: Spider Mites (other than Western United States). Apply 12 lbs. per acre. Aphids (other than Western United States). Apply 12 to 25 lbs. per acre. Plant Bugs including Lygus Bugs and Spider Mites (Western United States only). Apply 25 to 40 lbs per acre. Do not apply after cotton bolls open. Do not graze meat or dairy animals in treated fields. Workers entering treated fields within 24 hours following application should wear protective clothing

Eggplant: Two-spotted Spider Mite. Apply 6 to 12 lbs. per acre. May be applied up to day of harvest

Grapefruit, Oranges, Tangelos, Tangerines: Citrus Fied Mite. Citrus Rust Mite. Texas Citrus Mite. Apply 50 to 100 lbs. per acre Citrus Thrips (Arizona & California only) Apply 35 to 100 lbs per acre May be applied up to day of harvest. Do not repeat applications within 90 days. On Tangelos and Tangerines, do not apply more than 2 times per season.

Grapes: Grape Leafhopper. Two-spotted Spider Mire. Willamette Mite (Western United States only) Apply 15 to 30 lbs per acre. Do not apply within 15 days of harvest. Do not make more than 2 applications during truiting season. Do not apply to Fehr-Szago. Muscat or Salvador varieties

LDREN

N CLOTHING

SKIN IDLING

) PRODUCTS

vildlife. Keep out of lakes. oly where runoff is likely to from areas treated. Do not of wastes. Shrimp and crabiliabel. Do not apply where as specified on this label.

d when bees are actively y it by burying with waste

Lemons, Limes: Citrus Red Mite. Citrus Rust Mite. Texas Citrus Mite. Appl., 50 to 100 lbs. per acre. Citrus Thrips (Arizona and California only). Apply 35 to 100 lbs. per acre. Apply only 1 time per season. Do not apply within 21 days of harvest.

Melons: Serpentine Leafminer Complex, Two-spotted Spider Mite. Western Polato Leafhopper. Apply 30 to 40 lbs. per acre. May be applied up to day of harvest

Nectarines: Brown Mite. European Red Mite. Apply 40 to 60 lbs per acre. Do not apply more than 2 times during the fruiting period. Do not apply within 30 days of harvest.


Peaches: Brown Mite. European Red Mite. Two-spotted Spider Mite. Apr., 40 to 60 lbs. per acre. Do not apply more than 2 times during the fruiting period. Do not apply within 30 days of harvest.

Peppers, Pimentos: Serpentine Leafminer, Two-spotted Spider Mite Apply 6 to 12 lbs. per acre. May be applied up to day of harvest

Plums, Prunes: Brown Mite. European Red Mite. Two-spotted Spider 'the Apply 40 to 60 lbs. per acre. Do not apply more than 2 times during the fruiting proof Do not apply within 45 days of harvest.

Strawberries: *Strawberry Leafroller. Two-spotted Spider Mite* Apply 20 to 30 lbs per acre. Do not apply within 2 days of harvest

Tomatoes: Leafminers. Tropical Spider Mite. Two-spotted Spider Mite. Apply 20 lbs per acre. Do not apply within 2 days of harvest.



Occidental Chemical Company

rong tions of sale: 1. Seller warrants that this product consists of the ingredients specified and is reasonably fit for use as directed on the label. No one other than an officer of Seller, is authorized to make any other warranty, guarantee or direction concerning this product. 2. Because the time, place, rate of application and other conditions of use are beyond Seller's control. Seller's liability from the handing, storage and use of this product is limited to replacement of product or refund of purchase price.



ETHION*4 DUST

ACTIVE INGREDIENT:

Ethion (IO.O.O.O.O. detraethy! SIS -methylene bisphosphorodithioate

INERT INGREDIENTS:

*U.S. Patent No. 2,873,228

EPA Reg No. 7001-223-AA EPA Est No 7001-CA-1



HARMFUL IF SWALLOWED **DO NOT BREATHE DUST** DO NOT GET ON SKIN, IN EYES, OR ON CLOTHING MAY BE ABSORBED THROUGH SKIN **WASH THOROUGHLY AFTER HANDLING**

DO NOT STORE NEAR FOOD AND FEED PRODUCTS

streams, ponds, tidal marshes and estuaries. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not apply when weather conditions favor drift from areas treated. Do not apply when weather conditions favor drift from areas treated. Do not apply when weather conditions favor drift from areas treated. Do not apply when weather conditions favor drift from areas treated. Do not apply when weather conditions favor drift from areas treated. Do not apply when weather conditions favor drift from areas treated. Do not apply when weather conditions favor drift from areas treated. Do not apply when weather conditions favor drift from areas treated. Do not apply when weather conditions favor drift from areas treated. Do not apply when weather conditions favor drift from areas treated. Do not apply when weather conditions favor drift from areas treated. Do not apply when weather conditions favor drift from areas treated. Do not apply when weather conditions favor drift from areas treated. Do not apply when weather conditions favor drift from areas treated. Do not apply when weather conditions favor drift from areas treated. Do not apply when weather conditions favor drift from areas treated. Do not apply when weather conditions favor drift from areas treated. contaminate water by cleaning of equipment or disposal of wastes Shrimp and crab United States only) Apply 15 to 30 lbs per acre. Do not apply within 15 days of may be killed at application rates recommended on this label. Do not apply where harvest. Do not make more than 2 applications during fruiting season. Do not apply to these are important resources. Apply this product only as specified on this label.

This product is toxic to bees and should not be applied when bees are actively visiting the area. Do not reuse empty container. Destroy it by burying with waste

DIRECTIONS FOR USE

4% To be applied by ground application or aircraft

Almonds: Brown Mite, European Red Mite. Two-spotted Spider Mite. Apply 30 to 50 this per acre. Do not apply more than once during fruiting period. Do not apply within 30 days of harvest

Apples: Apple Rust Mite Brown Mite, Codling Moth, European Red Mite, Twospotted Spider Mite. Apply 50 to 90 lbs per acre. For Codling Moth, apply at petal fall Peacher and every 10 to 14 days thereafter until control is achieved. Do not apply within 40 days of harvest if no more than 2 applications are made. Do not use on early within 1.1. Intosh variety, or varieties ripening before or at the same time as early McIntosh.

Pepper Beans: Bean Leaf Beetle, Mexican Bean Beetle, Two-spotted Spider Mite. Apply 15 to 20 lbs per acre. Do not apply within 2 days of harvest. Do not feed treated to liage to Plums, vestock Lygus Bugs. Apply 30 to 40 lbs. per acre. Do not apply within 4 days of harvest. Do not feed treated foliage to livestock.

Cotton: Spider Mites (other than Western United States). Apply 12 lbs. per acre. Strawbe Aphids (other than Western United States). Apply 12 to 25 lbs per acre. Plant Bugs including Lygus Bugs and Spider Mites (Western United States only). Apply 25 to 40 Tomato lbs. per acre. Do not apply after cotton bolls open. Do not graze meat or dairy animals in freated fields. Workers entering treated fields within 24 hours following application per across should wear protective clothing.

Eggplant: Two spotted Spider Mite. Apply 6 to 12 lbs per acre. May be applied up to day of harvest

Grapetruit, Oranges, Tangelos, Tangerines: Citrus Red Mite, Citrus Rust Mite, Texas Citrus Mite Apply 50 to 100 lbs per acre Citrus Thrips (Arizona & California only). Apply 35 to 100 lbs per acre. May be applied up to day of harvest. Do not repeat This product is toxic to fish, shrimp, crabs and other wildlife. Keep out of lakes, applications within 90 days. On Tangelos and Tangerines, do not apply more than 2

Fehr-Szago Muscat or Salvador varieties

Lemon Melons:

DUST

DIRECTIONS FOR USE

to be applied by ground application or aircraft...

Almonds: Brown Mite, European Red Mite, Two-spotted Spider Mite. Apply 30 to 50 bs. per acre. Do not apply more than once during fruiting period. Do not apply within 40 days of harvest.

Apples: Apple Rust Mite. Brown Mite, Codling Moth, European Red Mite. Two-potted Spider Mite. Apply 50 to 90 lbs per acre For Codling Moth, apply at petal fall and every 10 to 14 days thereafter until control is achieved. Do not apply within 40 days of harvest if no more than 2 applications are made. Do not use on early Volatiosh variety, or varieties repening before or at the same time as early Mothtosh.

Beans: Bean Leaf Beetle, Mexican Bean Beetle, Two-spotted Spider Mite. Apply 15 to 20 lbs. per acre. Do not apply within 2 days of harvest. Do not feed treated foliage to vestock. Lygus Bugs. Apply 30 to 40 lbs. per acre. Do not apply within 4 days of narvest. Do not feed treated foliage to livestock.

Cotton: Spider Mites (other than Western United States) Apply 12 lbs per acre Aphids (other than Western United States) Apply 12 to 25 lbs per acre. Plant Bugs including Lygus Bugs and Spider Mites (Western United States only) Apply 25 to 40 lbs. per acre. Do not apply after cotton bolls open. Do not graze meat or dairy animals in treated fields. Workers entering treated fields within 24 hours following application should wear protective clothing.

Eggplant: Two-spotted Spider Mite. Apply 6 to 12 lbs. per acre. May be applied up to day of harvest.

Grapetruit, Oranges, Tangelos, Tangerines: Citrus Red Mite. Citrus Rust Mite. Texas Citrus Mite. Apply 50 to 100 lbs per acre. Citrus Thrips (Arizona & California only) Apply 35 to 100 lbs per acre. May be applied up to day of harvest. Do not repeat applications within 90 days. On Tangelos and Tangerines, do not apply more than 2 times, per season.

Grapes: Grape Leafhopper Two-spotted Spider Mite, Willamette Mite (Western United States only) Apply 15 to 30 lbs per acre. Do not apply within 15 days of harvest. Do not make more than 2 applications during fruiting season. Do not apply to Fehr-Szago. Muscat or Salvador varieties.

Lemons, Limes: Citrus Red Mite, Citrus Rust Mite, Texas Citrus Mite, Apply 50 to 100 lbs per acre. Citrus Thrips (Arizona and California only) Apply 35 to 100 lbs per acre. At ply only 1 time per season. Do not apply within 21 days of harvest

Melons: Serpentine Leafminer Complex Two-spotted Spider Mite. Western Potato Leafminer Apply 30 to 40 lbs per acre. May be applied up to day of narvest Nectarines: Brown Mite. European Red Mite. Apply 40 to 60 lbs per acre. Do not apply more than 2 times during the fruiting period. Do not apply within 30 days of bary end.


Peaches: Brown Mite: European Red Mite: Two-spotted Spider Mite. Apply 40 to 60 lbs: per acre. Do not apply more than 2 times during the fruiting period. Do not apply within 30 days of a givest.

Peppers, Pimentos: Serpentine Lealminer Two-spotted Spider Mite Apply 6 to 12 lbs per acre. May be applied up to day of harvest

Plums, Prunes: Brown Mite European Red Mite Two-spotted Spider Mite Apply 40 to 60 bs per acre. Do not apply more than 2 times during the fruiting period. Do not apply within 45 days of harvest.

Strawberries: Strawberry Leatroller Two-spotted Spider Mite. Apply 20 to 30 lbs. per acros. (To not app.), within 2 days of harvest.

Tomatoes: Leatminers Tropical Spider Mite. Two-spotted Spider Mite. Apply 20 lbs per acre. Do not apply a transfer of the content of the cont



Common vata

Occidental Chemical Company

CONDITIONS OF SALE: 1 Seller warrants that this product consists of the ingredients specified and is sensimably it for vie as directed on the label. No creative than an officer of Seller, is authorised to make any other warrants, gubantee as direction common goths trudget. 2 Sections the time, place, in form any or and other world the country of the law of the control of product products of the country form the control of products of products price.