



GAVICIDE 2,4-D L.V. 4

WEED KILLER

INGREDIENTS:

Diethyl ester of 2,4-Dichlorophenoxyacetic acid	69.7%
2,4-Dichlorophenoxyacetic acid	30.3%
	100.0%

4.0 pounds 2,4-Dichlorophenoxyacetic acid per gallon.

CAUTION

KEEP OUT OF REACH OF CHILDREN

May cause skin irritation. Avoid contact with eyes, skin, or clothing. Do not store near food, fertilizer, seeds, insecticides, or fungicides. Do not swallow.

P. No. 1202-113 NET CONTENTS GALLON



SMALL GRAINS (Wheat, Barley, Rye, Grasses): Use 1/2 pint per acre in grain for controlling annual and biennial weeds such as ~~Redweed~~, Mustard, Lambsquarter, Pigweed, and Wild Radish. Apply 3/4 pint per acre to control perennial weeds such as Bindweed, Knotweed, Canada Thistle, and Wild Morning Glory. Treat winter wheat only in spring when weeds are small and before the crop begins heading. Grain interplanted with legumes should not be sprayed unless injury to the legumes can be tolerated. Do not forage or graze treated grain fields within 2 weeks after treatment 2,4-D. Do not feed treated straw to livestock.

Treat grain after tillering has begun and before it emerges from the boot or before jointing. Application at other times may cause abnormal heading.

CORN: For controlling susceptible annual or perennial weeds, apply not more than 1/2 pint per acre. If crop is 8 inches or taller, avoid application of 2,4-D to the growing point. Use drop nozzles and direct spray near base of the crop. Do not use after tassels appear. Different varieties may vary in susceptibility to 2,4-D. Do not use 2,4-D in corn grown for the production of hybrid seed. Check with state agricultural experimental stations or extension service weed specialist for specific recommendations. Avoid cultivating soon after treatment, when corn may be brittle.

For pre-emergence treatment of weeds in corn, use 1-1/2 to 3 pints per acre, applied as a broadcast spray one to three days before crop emergence. Do not use pre-emergence sprays on light, sandy soils.

SORGHUM: Apply 1/2 pint per acre. Crop is most tolerant to 2,4-D when 4 to 12 inches tall. Severe injury may result if 2,4-D is applied to plants less than 3 inches tall, or after 12 inches tall through bloom period. Check with state agricultural experimental stations or extension service weed specialists for specific recommendations.

PASTURES, LAWNS, GOLF COURSES AND SIMILAR TURF: Apply 2 pints GAVICIDE 2,4-D L.V. 4 Weed Killer per acre in sufficient water to provide good coverage. For small areas, use 1 tablespoonful per gallon of water. Do not apply to newly seeded lawns until grass has become well established. Do not use on bent or other creeping grasses. Clover or other legumes may be injured by 2,4-D, especially during hot summer weather. Avoid spraying grains or seed grasses when in the blossoming or pollinating stage.

NON-CROP AREAS: For controlling perennial weeds in fence rows, ditch-banks, roadsides, and similar areas, use 2 to 3 pints in 100 gallons of water and wet foliage thoroughly. Repeat applications may be necessary on certain difficult-to-kill perennials, such as Canada Thistle, Bindweed, Wild Morning Glory, and White Top. Re-treat when regrowth is visible. Use only where drift will not injure susceptible vegetation. **NOTE:** For spot treatment of weeds in crops listed above, use 1 1/2 tablespoonfuls per gallon of water.

WOODY PLANTS: For susceptible woody species such as Willow, Sumac, Buckbrush, or Grape, use 3 1/2 to 6 pints per hundred gallons of water and wet all foliage thoroughly. For small areas, use 2 tablespoonfuls per gallon of water. Best results are obtained when brush is growing rapidly, in late spring or early summer. Brush may require several applications to control resistant species. Do not use where drift may injure desirable plants.

For spraying stumps to prevent resprouting, dilute one gallon in 15 to 20 gallons of No. 2 fuel oil or diesel oil and wet thoroughly. For small areas, use 1/3 pint per gallon of oil. Application may be made at any time of year.

CONDITIONS OF SALE: 1. PureGro Company (manufacturer) warrants that this material conforms to the chemical description on the label and is reasonably fit for use as directed hereon. Manufacturer neither makes, nor authorizes any agent or representative to make, any other warranty of FITNESS or of MERCHANTABILITY, guarantee or representation, express or implied, concerning this material.

2. Critical and unforeseeable factors beyond the manufacturer's control prevent it from eliminating all risks in connection with the use of chemicals. Such risks include, but are not limited to, damage to plants and crops to which the material is applied, lack of complete control, and damage caused by drift to other plants or crops. Such risks occur even though the product is reasonably fit for the uses stated hereon and even though label directions are followed. Buyer and user acknowledge and assume all risks and liability (except those assumed by the manufacturer under 1 above) resulting from handling, storage, and use of this material.

LOT NO. 27767RJW



~~GAVIGIDE~~ 2,4-D

BUTYL 6

ACTIVE INGREDIENTS:

*Butyl ester of
2,4-Dichlorophenoxyacetic acid 79%

INERT INGREDIENTS 21%
100%

*Equivalent to 61.5% 2,4-Dichlorophenoxyacetic acid or 6.0 pounds
per gallon.

CAUTION

KEEP OUT OF REACH OF CHILDREN

CAUTION: May cause skin irritation. Avoid contact with eyes, skin, or clothing. Do not store near food, fertilizers, seeds, insecticides, or fungicides. Do not contaminate water used for domestic or spray purposes. Harmful if swallowed.

