

SIMPLON LATHROP CALIFORNIA

7001-259

2-3-87

P.04

New Version 1/2

Not to be sold, offered for sale, held for sale, shipped, delivered for shipment, offered for delivery, or received after December 31, 1988. Not to be used after March 31, 1989. Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinseate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your state pesticide or environmental control agency, or the hazardous waste representative at the nearest EPA regional office for guidance.

# Dicofol



| ACTIVE INGREDIENTS:                          | BY WT.        |
|--|---------------|
| 1,1-Bis(chlorophenyl)-2,2,2-trichloroethanol | 41.1%         |
| INERT INGREDIENTS:                           | 58.9%         |
| <b>TOTAL</b>                                 | <b>100.0%</b> |

EPA Reg. No. 7001-259  
EPA Est. No. 7001-CA-1 Contains 4 lb Dicofol per gallon

**KEEP OUT OF REACH OF CHILDREN**

## CAUTION

### PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Avoid skin contact. Wash thoroughly after using. Avoid inhalation of mist. Harmful if swallowed. If taken internally get medical attention. Do not store near feed or foodstuffs.

### STATEMENT OF PRACTICAL TREATMENT

**IF SWALLOWED** call a physician immediately. If no other treatment is available induce vomiting by giving victim one or two glasses of water and insert finger in back of throat. Repeat until vomit fluid is clear. Do not induce vomiting or give anything by mouth to an unconscious person.

**IF INHALED** remove victim to fresh air. Apply respiration if breathing is irregular or has stopped. Call a physician immediately.

**IF ON SKIN** wash immediately with soap and water.

**IF IN EYES** flush eyes for at least 15 minutes with water. Call a physician immediately.

### ENVIRONMENTAL HAZARDS

This product is toxic to fish. Keep out of any body of water.

# 4 e c

SIMPLOT LATHROP CALIFORNIA

P. 05

2/2

**CHEMICAL HAZARDS**

**WARNING** - This product must be stored at temperatures above 40° F. Storage below may affect the quality of the product.

**RE-ENTRY STATEMENT**

Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons. Do not enter treated areas with protective clothing until sprays have dried.

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Written and oral warnings must include the following information: **CAUTION.** Area treated with Dicofof 4EC on (date of application). Do not enter without appropriate protective clothing until sprays have dried. **REQUIRED PROTECTIVE CLOTHING:** Long sleeved shirt, long legged pants, rubber gloves and a respirator approved by the Mining Enforcement and Safety Administration and by the National Institute for Occupational Safety and Health.

**DIRECTIONS FOR USE**

It is a violation of federal law to use this product in a manner inconsistent with its labeling. Skin contact with this pesticide may be hazardous; wear chemical resistant gloves when mixing, loading, or applying this product.

**STORAGE AND DISPOSAL**

Do not contaminate water, food or feed by storage or disposal. Wastes resulting from the use of this product may be disposed of on-site or at an approved waste disposal facility.

**STORAGE:** Store at temperatures above 40° F. Storage below may affect the quality of the product.

**DISPOSAL:** Triple rinse (or equivalent). Then offer for recycling, or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities.

**DIRECTIONS FOR DILUTION**

**DILUTE APPLICATION:** *Ground Application:* Apply specified rate in 20 to 80 gallons of water per acre. *Orchard Application:* Apply specified rate in 100 to 800 gallons of water per acre.

**CONCENTRATE APPLICATION:** *Ground Application:* Apply specified rate in not less than 5 gallons of water per acre. *Orchard Application:* Apply specified rate in 20 to 100 gallons of water per acre. Special concentrate equipment is necessary for these uses.

**AIR APPLICATION:** Apply specified rate in 5 to 20 gallons of water per acre. Orchard rates should not be applied in less than 10 gallons of water per acre.

**Alfalfa, Clover (Both grown for seed only—use restricted to California, Idaho, Nevada, Oregon and Washington):** *Carmine, Pacific, Strawberry and Two-Spotted Mites.* Apply 1/4 to 1 1/2 qts. per acre. Do not graze livestock in treated fields, or feed crop residues or cuttings to livestock or bale it for feed. Field must be cleared by burning crop residues or by discing them into the soil.

**Apples, Pears (Western states only):** *Apple Rust, Brown, European Red, McDaniel Spider, Pacific, Schoene Spider, Willamette and Yellow Spider Mites.* Apply 1/2 to 4 qts. per acre. Do not apply within 7 days of harvest. Dicofof should not be used in late cover sprays with glyodin as plant injury may result. Dicofof may cause fruit spotting under certain conditions and only the lower dosages should be used. Control for *Brown Mite* and *European Red Mite* on pears are for cleanup of late spray population. Repeat as needed, but allow 30 days between treatments.

**Apricots:** *Brown, European Red, Pacific Spider, Peach Silver, Plum Rust, Two-Spotted Spider and Willamette Mites.* Apply 1/2 to 2 1/2 qts. per acre. Allow 30 days between treatments. Do not apply within 14 days of harvest.

**Beans:** *Carmine Spider, Strawberry Spider, Pacific Spider, Two-Spotted Mites.* Apply 1/3 to 1 1/2 qts. per acre other than California. Apply 7 days before harvest through 1 qt. per acre. Apply 1/2 to 1 1/2 qts. 21 days before harvest. *California Only:* Apply 1/4 to 1 1/2 qts. 45 days before harvest. Do not feed treated fresh hay or spent hay to dairy or meat animals.

**Bleackberries:** *Brown, European Red, McDaniel Spider, Pacific Spider, Two-Spotted Spider, Willamette and Yellow Spider Mites.* Apply 1/2 to 1 qt. per acre. Do not apply within 2 days of harvest.

**Cherries:** *Brown, European Red, McDaniel Spider, Pacific Spider, Peach Silver, Plum Rust, Two-Spotted Spider, Willamette and Yellow Spider Mites.* Apply 1/2 to 2 1/2 qts. per acre. Do not apply within 7 days of harvest. Repeat if needed, but allow 30 days between treatments.

**Citrus (Grapefruit, Kumquats, Lemons, Limes, Oranges, Tangelos, Tangerines):** *Citrus Flat, Citrus Red, Citrus Rust, Six-Spotted, Texas Citrus, Yuma Spider, Pacific Spider, Two-Spotted Spider Mites.* Apply 1/2 to 4 qts. per acre. Do not apply within 7 days of harvest.

**Cotton:** *Desert Spider, Pacific Spider, Two-Spotted Spider, Strawberry Spider Mites.* Apply 1/4 to 1 1/2 qts. per acre. Do not apply within 14 days of harvest. Do not graze dairy or meat animals in treated fields or feed gin trash to them.

**Cucumbers, Melons (Including Watermelons), Pumpkins, Squash:** *Pacific Six Mite, Two-Spotted Spider Mite.* Apply 1/3 to 2/3 pt. per acre. Do not apply within 2 days of harvest.

**Crabapples:** *Apple Rust Mite, Brown Mite, European Red Mite, McDaniel Spider Mites, Pacific Spider Mite, Schoene Spider Mite, Yellow Spider Mite, Two-Spotted Spider Mite, Willamette Mite.* Apply 1/4 to 4 qts. per acre. Claims for *Brown Mite* and *European Red Mite* are for clean-up of late spring populations. Re as needed, but allow 30 days between treatments. As preventive schedule a 1/2 pint per acre. Make two successive applications at 10 to 14 day intervals.

**Deerberries (Including Boysenberries, Loganberries, Oatlike Berries):** *Brown Mite, European Red, McDaniel Spider, Pacific Spider, Willamette, Yellow Spider Mite.* Apply 1/2 to 1 qt. per acre. Do not apply within 2 days of harvest.

**Figs:** *European Red Mite, Pacific Spider Mite, Two-Spotted Spider Mite.* Apply 1/2 to 2 qts. per acre. Do not apply within 7 days of harvest.

**Filberts, Walnuts, Hickory Nuts, Pecans, Chestnuts:** *Brown Mite, European Red Mite, Two-Spotted Spider, Pacific Spider Mites, Yellow Spider Mite.* Apply 1/4 to 4 qts. per acre. Do not apply within 14 days of harvest. Do not feed treated husk livestock.

**Grapes:** *European Red, McDaniel Spider, Pacific Spider, Two-Spotted Spider, Willamette Mites.* Apply 1/2 to 1 qt. per acre. Do not apply within 7 days of harvest.

**Hops:** *Pacific Spider, Two-Spotted Spider Mite.* Apply 1/4 to 1 qt. per acre. Do not apply within 7 days of harvest.

**Nectarines, Peaches:** *Brown Mite, European Red, McDaniel Spider, Peach Silver, Peach Silver, Schoene Spider, Two-Spotted Spider, Yellow Spider Mite.* Apply 1/4 to 2 1/2 qts. per acre. Do not apply within 14 days of harvest.

**Pears:** *Apple Rust Mite, Brown Mite, European Red Mite, McDaniel Spider, Pacific Spider, Schoene Spider, Two-Spotted Spider, Willamette, Yellow Spider Mite.* Apply 1/2 to 4 qts. per acre. Claims for *Brown Mite* and *European Red Mite* are for cleanup of late spring populations. Repeat as needed but allow 30 days between treatments. Dicofof should not be used in late cover sprays with glycin as plant injury may result.

**Peppers, Tomatoes:** *Pacific Spider, Tomato Russet, Tropical Spider, Two-Spotted Spider Mites.* Apply 1/2 to 1/4 qt. per acre. Do not apply within 2 days of harvest.

**Plums, Prunes:** *Brown Mite, European Red, McDaniel Spider, Pacific Spider, Peach Silver, Plum Rust, Two-Spotted Spider Mites.* Apply 1/2 to 2 qts. per acre. Do not apply within 7 days of harvest. Repeat if needed, but allow 30 days between treatments.

**Quince:** *Apple Rust, Brown, European Red, McDaniel Spider, Pacific Spider, Schoene Spider, Two-Spotted Spider, Willamette, Yellow Spider Mites.* Apply 1/2 to 4 qts. per acre. Do not apply within 7 days of harvest.

**Raspberries:** *European Red, McDaniel Spider, Pacific Spider, Two-Spotted Spider, Willamette, Yellow Spider Mites.* Apply 1/2 to 1 qt. per acre. Do not apply within 2 days of harvest.

**Strawberries:** *Strawberry Spider, Two-Spotted Spider Mites.* Apply 1/2 to 2 qts. per acre. *Cyclamen Mites.* Apply 2 qts. per acre. Do not apply within 2 days of harvest; to control *Cyclamen Mites*, make 2 or 3 applications at 10 to 20 day intervals starting when new spring growth begins. Sprays should be thorough dranching, high pressure applications. Small areas of infestation may also be treated during the production season.

**ORNAMENTALS**

**Lawns, Turf:** *Bermudagrass - Clover Mites.* Apply 1 oz. in 2.5 gallons of spray per 1,000 sq. ft. Repeat in 10 to 14 days if needed. For *Clover Mite*, be sure to treat areas at base of buildings.

**Flowering Plants (Including African Violets, Chrysanthemums, Sweet Pea Orchids, Shrubs, Trees, Vines, Nursery Stock):** *Mites.* Apply 1 oz. in 2.5 gallons of water. Thoroughly cover upper and lower leaf surfaces. Wet to the dripping point.

AGC 585 :2

**J. R. SIMPLOT COMPANY**

Agricultural Chemicals Department • P. O. Box 198, Lathrop, California 95330

**DISCLAIMER OF WARRANTIES.** Seller warrants that the chemical composition of this product conforms to the chemical description given on this label. THIS WARRANTY IS EXPRESSLY LIMITED TO ALL OTHER WARRANTIES AND REPRESENTATIONS, EXPRESSED, IMPLIED OR STATUTORY INCLUDING WARRANTIES OF MERCHANTABILITY AND FITNESS FOR A PARTICULAR USE. Timing, rate and method of application, weather and crop conditions, mixtures of chemicals not specifically recommended on this label or an accompanying written recommendation are beyond the control of seller. Buyer assumes all risks of use, storage and handling of this material in strict accordance with directions given hereon. Buyer further agrees in the event of damage arising from the use of this product to accept a replacement of this product or a refund of the purchase price of the product, at buyer's option, as full discharge of seller's liability. No one is authorized to make any other warranty, guarantee or directions concerning this product, and no such warranties, guarantees or directions shall be valid or binding upon seller.

Simplot

NET CONTENTS 5 GALLONS

**WARNING—THIS PRODUCT MUST BE STORED AT TEMPERATURES ABOVE 40° F.  
STORAGE BELOW MAY AFFECT THE QUALITY OF THE PRODUCT.**

Fast Pass