

Not to be sold, offered for sale, held for sale, shipped, delivered, for shipment, offered for delivery, or received after December 31, 1988. Not to be used after March 31, 1989. Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your state pesticide or environmental control agency, or the hazardous waste representative at the nearest EPA regional office for guidance.

Dicofol



ACTIVE INGREDIENTS:	BY WT.
1,1-Bis(chlorophenyl)-2,2,2-trichloroethanol	41.1%
INERT INGREDIENTS:	58.9%
TOTAL	100.0%

EPA Reg. No. 7001-259
 EPA Est. No. 7001-CA-1 Contains 4 lb Dicofol per gallon

KEEP OUT OF REACH OF CHILDREN

CAUTION

PRECAUTIONARY STATEMENTS
 HAZARDS TO HUMANS
 AND DOMESTIC ANIMALS

Avoid contact. Wash thoroughly after using. Avoid inhalation of mist. Harmful if swallowed. If taken internally get medical attention. Do not store near feed or foodstuffs.

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED call a physician immediately. If no other treatment is available induce vomiting by giving victim one or two glasses of water and insert finger in back of throat. Repeat until vomit fluid is clear. Do not induce vomiting or give anything by mouth to an unconscious person.

IF INHALED remove victim to fresh air. Apply respiration if breathing is irregular or has stopped. Call a physician immediately.

IF ON SKIN wash immediately with soap and water.

IF IN EYES flush eyes for at least 15 minutes with water. Call a physician immediately.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Keep out of any body of water.

ACCEPTED

OCT 23 1987

Under Fungicide, Insecticide, and Plant Growth Regulator Act, as amended, the pesticide registered under EPA Reg. No. 7001-259

499

CHEMICAL HAZARDS

WARNING—This product must be stored at temperatures above 40°F. Storage below may affect the quality of the product.

RE-ENTRY STATEMENT

Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons. Do not enter treated areas without protective clothing until sprays have dried.

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Written and oral warnings must include the following information: CAUTION. Area treated with Dicolol 4EC on (date of application). Do not enter without appropriate protective clothing until sprays have dried. REQUIRED PROTECTIVE CLOTHING: Long sleeved shirt, long legged pants, rubber gloves and a respirator approved by the Mining Enforcement and Safety Administration and by the National Institute for Occupational Safety and Health.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling. Skin contact with this pesticide may be hazardous; wear chemical resistant gloves when mixing, loading, or applying this product.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

STORAGE: Store at temperatures above 40°F. Storage below may affect the quality of the product.

DISPOSAL: Triple rinse (or equivalent). Then offer for recycling, or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities.

DIRECTIONS FOR DILUTION

DILUTE APPLICATION: *Ground Application:* Apply specified rate in 20 to 60 gallons of water per acre. *Orchard Application:* Apply specified rate in 100 to 800 gallons of water per acre.

CONCENTRATE APPLICATION: *Ground Application:* Apply specified rate in not less than 5 gallons of water per acre. *Orchard Application:* Apply specified rate in 20 to 100 gallons of water per acre. Special concentrate equipment is necessary for these uses.

AIR APPLICATION: Apply specified rate in 5 to 20 gallons of water per acre. Orchard rates should not be applied in less than 10 gallons of water per acre.

Apples, Pears (Western states only): *Apple Rust, Brown, European Red, McDaniel Spider, Pacific, Schoone Spider, Willamette and Yellow Spider Mites.* Apply 1½ to 4 qts. per acre. Do not apply within 7 days of harvest. Dicolol should not be used in late cover sprays with glyodin as plant injury may result. Dicolol may cause fruit spotting under certain conditions and only the lower dosages should be used. Control for *Brown Mite* and *European Red Mite* on pears are for cleanup of late spring population. Repeat as needed, but allow 30 days between treatments.

Beans: *Carmina Spider, Strawberry Spider, Pacific Spider, Two-Spotted Mites.* Apply 1/3 to 1½ qts. per acre other than California. Apply 7 days before harvest through ½ qt. per acre. Apply ½ to 1½ qts. 21 days before harvest. *California Only:* Apply ¾ to 1½ qts. 45 days before harvest. Do not feed treated fresh hay or spent hay to dairy or meat animals.

Citrus (Grapefruit, Kumquats, Lemons, Limes, Oranges, Tangelos, Tangerines): *Citrus Flat, Citrus Red, Citrus Rust, Six-Spotted, Texas Citrus, Yuma Spider, Pacific Spider, Two-Spotted Spider Mites.* Apply 1½ to 4 qts. per acre. Do not apply within 7 days of harvest.

CORN (Field): For the control of *Two-Spotted, Pacific and Grass Mites.* EARLY SEASON CONTROL ONLY. Use three pints Dicolol 4 EC per acre either by ground sprayer or aircraft in sufficient water to provide thorough coverage, 40 to 60 gallons per acre for ground equipment and 10 to 15 gallons per acre for aircraft. Apply when mites first appear, and repeat if necessary. DO NOT apply after ears begin to form. DO NOT use treated corn plant for ensilage or as forage for livestock. The field must be cleared by burning residues or discing into the soil.

Cotton: *Desert Spider, Pacific Spider, Two-Spotted Spider, Strawberry Spider Mites.* Apply ¾ to 1½ qts. per acre. Do not apply within 14 days of harvest. Do not graze dairy or meat animals in treated fields or feed gin trash to them.

Cucumbers, Melons (Including Watermelons), Pumpkins, Squash: *Pacific Spider Mite, Two-Spotted Spider Mite.* Apply 1/3 to 2/3 pt. per acre. Do not apply within 2 days of harvest.

Crabapples; *Apple Rust Mite, Brown Mite, European Red Mite, McDaniel Spider Mites, Pacific Spider Mite, Schoone Spider Mite, Yellow Spider Mite, Two-Spotted Spider Mite, Willamette Mite.* Apply 1½ to 4 qts. per acre. Claims for *Brown Mite* and *European Red Mite* are for cleanup of late spring populations. Repeat as needed, but allow 30 days between treatments. As a preventive schedule apply ½ pint per acre. Make two successive applications at 10 to 14 day intervals.

Filberts, Walnuts, Hickory Nuts, Pecans, Chestnuts: *Brown Mite, European Red, Two-Spotted Spider, Pacific Spider Mites, Yellow Spider Mite.* Apply 1½ to 4 qts. per acre. Do not apply within 14 days of harvest. Do not feed treated husks to livestock.

Grapes: *European Red, McDaniel Spider, Pacific Spider, Two-Spotted Spider, Willamette Mites.* Apply ¾ to 1 qt. per acre. Do not apply within 7 days of harvest.

Hops: *Pacific Spider, Two-Spotted Spider Mite.* Apply ½ to 1 qt. per acre. Do not apply within 7 days of harvest.

Mint (peppermint and spearmint for oil production): *Two-spotted Spider Mite.* Apply ¾ to 1 qt. per acre. Do not apply within 30 days of harvest. Do not make more than one application per crop season. Do not feed treated fresh hay or spent hay to livestock.

Pears: *Apply Rust Mite, Brown Mite, European Red Mite, McDaniel Spider, Pacific Spider, Schoone Spider, Two-Spotted Spider, Willamette, Yellow Spider Mite.* Apply 1½ to 4 qts. per acre. Claims for *Brown Mite* and *European Red Mite* are for cleanup of late spring populations. Repeat as needed but allow 30 days between treatments. Dicolol should not be used in late cover sprays with glyodin as plant injury may result.

Peppers, Tomatoes: *Pacific Spider, Tomato Russet, Tropical Spider, Two-Spotted Spider Mites.* Apply ¾ to 1 qt. per acre. Do not apply within 2 days of harvest.

Quince: *Apply Rust, Brown, European Red, McDaniel Spider, Pacific Spider, Schoone Spider, Two-Spotted Spider, Willamette, Yellow Spider Mite.* Apply 1½ to 4 qts. per acre. Do not apply within 7 days of harvest.

Strawberries: *Strawberry Spider, Two-Spotted Spider Mites.* Apply ¾ to 2 qts. per acre. *Cyclamen Mites.* Apply 2 qts. per acre. Do not apply within 2 days of harvest; to control *Cyclamen Mites*, make 2 or 3 applications at 10 to 20 day intervals starting when new spring growth begins. Small areas of infestation may also be treated during the production season.

ORNAMENTALS

Lawns, Turf: *Bermudagrass—Clover Mites.* Apply 1 oz. in 2.5 gallon spray per 1,000 sq. ft. Repeat in 10 to 14 days if needed. For *Clover Mites*, be sure to treat areas at base of buildings.

Flowering Plants (Including African Violets, Chrysanthemums, Peas), *Orchids, Shrubs, Trees, Vines, Nursery Stock:* *Mites.* Apply ½ to 2.5 gallons of water. Thoroughly cover upper and lower leaf surfaces. Wet to the dripping point.

AGC 555 (5-77)

NET CONTENTS 5 GALLONS

Simplot

J. R. SIMPLOT COMPANY

Agricultural Chemicals Department

P.O. Box 198

Lathrop, CA 95330

DISCLAIMER OF WARRANTIES: Seller warrants that the chemical composition of this product conforms to the chemical description given on this label. THIS WARRANTY IS EXPRESSLY IN LIEU OF ALL OTHER WARRANTIES AND REPRESENTATIONS, EXPRESSED, IMPLIED, OR STATUTORY, INCLUDING WARRANTIES OF MERCHANTABILITY AND FITNESS FOR A PARTICULAR USE. Timing, rate and method of application, weather and crop conditions, mixtures with chemicals not specifically recommended on this label or an accompanying written recommendation, are beyond the control of seller. Buyer assumes all risks of use, storage and handling of this material not in strict accordance with recommendations given herewith. Buyer further agrees in the event of damages arising from the use of this product to accept a replacement of this product or a refund of the purchase price of the product, at buyer's option, as full discharge of seller's liability. No one is authorized to make any other warranty, guarantee or directions concerning this product, and no such warranties, guarantees or directions shall be valid or binding upon seller.

WARNING—THIS PRODUCT MUST BE STORED AT TEMPERATURES ABOVE 40° F.
STORAGE BELOW MAY AFFECT THE QUALITY OF THE PRODUCT.