




SULFUR 6 FLOWABLE



ACTIVE INGREDIENT

Sulfur.....


INERT INGREDIENTS

.....

Contains 6 lbs. Sulfur per gallon

EPA REG. NO. 7001-261

EPA EST. NO. 7001-CA-1



BY WT.

52.0%

48.0%

TOTAL 100.0%

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or cleaning of equipment. Do not store near other pesticides or seeds. Open dumping is prohibited.

Pesticide, spray mixture or rinsate that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies.

Triple rinse (or equivalent) and offer for recycling, reconditioning, or disposal in approved landfill or bury in a safe place.

Consult Federal, State or local disposal authorities for approved alternative procedures.

Apples: Apples cedar rust, black rot, frog-eye leaf spot, powdery mildew, quince rust, scab, sooty-blotch. Apply 1 1/2 to 7 pts. in 100 gallons of water per acre from prepink through calyx periods and 1/3 to 5 1/2 pts. in 100 gallons of water per acre in cover sprays. NOTE: McIntosh, Golden Delicious, Jonathan and certain other varieties may be injured by sulfur applications under certain conditions.

Cherries: Brown rot, leaf spot, powdery mildew, rust, scab and stem blight - Apply 1 1/2 to 3 1/2 pts. in 100 gallons of water per acre through bloom stage and 1/2 to 1 1/2 pts. in 100 gallons of water from petal fall through cover sprays for brown rot and leaf spot.

Citrus: Rust mite, clover mite - Apply 1/2 to 4 1/2 pts. in 100 gallons of water per acre in November through May. Thrips (Early spring treatment for the reduction of thrip populations) - Apply 1 to 2 gallons in 100 gallons of water per acre for thorough coverage. Make application after spring flush has made 3 to 4 inch growth. Repeat if necessary. Do not apply within 2 months of an oil application.

Grapes: Powdery mildew, rust (Physopella) - Apply 1 to 7 pts. in 100 gallons of water per acre using up to 8 pts. per acre. Begin when new shoots are 6 to 10 inches long. Repeat before blossoms open and continue at 10 to 14 day intervals as necessary. Note: Concord and other labrusca type grapes may be injured.

Nectarines, Peaches: Brown rot, leaf spot, powdery mildew, peach canker, rust, scab, shothole - Apply 1 1/2 to 3 1/2 pts. in 100 gallons of water per acre through bloom and 1/2 to 1 1/2 pts. in 100 gallons of water per acre in cover sprays. Apply from early petal fall through cover periods as necessary.

Pears: Brown rot, leaf spot, powdery mildew, rust, scab - Apply 1 1/2 to 3 pts. in 100 gallons of water per acre through calyx and 1/2 to 2 pts. in 100 gallons of water per acre in cover sprays. Apply from prebloom through calyx periods as necessary.

Plums, Prunes: Brown rot, leaf spot, powdery mildew, rust, scab, shothole - Apply 1 1/2 to 7 1/2 pts. in 100 gallons of water per acre through bloom and 1/2 to 2 pts. in 100 gallons of water per acre through cover sprays for brown rot and leaf spot.

Strawberries: Powdery mildew - Apply 1 pt. in 100 gallons of water per acre beginning at first sign of disease and repeat as necessary. NOTE: Sulfur may injure strawberries under certain climatic conditions.

KEEP OUT OF REACH OF CHILDREN

CAUTION

Harmful if swallowed. Avoid breathing spray mist. May cause irritation of nose, throat or skin. Avoid contact with eyes, skin and clothing.

FIRST AID

IF SWALLOWED, drink 1 or 2 glasses of water and induce vomiting by sticking finger or blunt object down throat.

IF IN EYES, wash immediately with water for 15 minutes

IF ON SKIN, wash with soap and water.

ENVIRONMENTAL HAZARDS

Keep out of lakes, streams or ponds. Do not contaminate water by cleaning equipment or disposal of waste

CAUTION: Some plants are susceptible to injury from sulfur under certain climatic conditions. The user is advised not to use sulfur on any crop unless local experience has proven that sulfur is safe in that locality.

During periods of high temperature sulfur may burn foliage and fruit. Do not make sulfur applications at excessively high temperatures. Do not use with oil or within 4 weeks of an oil application.

DIRECTIONS FOR DILUTION

Stir until smooth before using. Wash recommended amount through screen into partially filled spray tank or pre-mix in a small amount of water before pouring into tank. Do not use a screen finer than 50 mesh in entire system. Keep agitator running during filling and spraying operations. Do not allow mixture to stand. Failure to maintain agitation will cause sulfur to settle and necessitate manual removal of the deposite.

This product should be applied by ground or aircraft, in sufficient water to provide thorough coverage.

MANUFACTURED BY

Occidental Chemical Company

P.O. Box 198, Lathrop, California 95330 P.O. Box 1185, Houston, Texas 77001

DISCLAIMER OF WARRANTIES: Seller warrants that the chemical composition of the product conforms to the chemical description given on this label. THIS WARRANTY IS EXPRESSLY IN LIEU OF ALL OTHER WARRANTIES AND REPRESENTATIONS, EXPRESSED, IMPLIED, OR STATUTORY, INCLUDING WARRANTIES OF MERCHANTABILITY AND FITNESS FOR A PARTICULAR USE. Timing, rate and method of application, weather and crop conditions, mixtures with chemicals not specifically recommended on this label or an accompanying written recommendation, are beyond the control of seller. Buyer assumes all risks of use, storage and handling of this material not in strict accordance with directions given here-with. Buyer further agrees in the event of damages arising from the use of this product to accept a replacement of the product or a refund of the purchase price of the product, at buyer's option, as full discharge of seller's liability. No one is authorized to make other warranty, guarantee or directions concerning this product, and no such warranties, guarantees or directions shall be valid or binding upon seller.