

7001-26 Pm 21 10F2

US ENVIRONMENTAL PROTECTION AGENCY OFFICE OF PESTICIDES PROGRAMS REGISTRATION DIVISION (TS-767) WASHINGTON, DC 20460	EPA REGISTRATION NO. 7001-261	DATE OF ISSUANCE
	TERM OF ISSUANCE	
	NAME OF PESTICIDE/PRODUCT Sulfur 6 Flowable	

FEB 2 198

NOTICE OF PESTICIDE: REGISTRATION
 REREGISTRATION
 (Under the Federal Insecticide, Fungicide,
 and Rodenticide Act, as amended)

NAME AND ADDRESS OF REGISTRANT (Include ZIP code)

J.R. Singlot Company
 16777 South Oakland Avenue
 P.O. Box 198
 Lathrop, CA 95330

182687
 2
 656
 17

NOTE: Changes in labeling formula differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above U.S. EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby Registered/Reregistered under the Federal Insecticide, Fungicide, and Rodenticide Act.

A copy of the labeling accepted in connection with this Registration/Reregistration is returned herewith.

Registration is in no way to be construed as an indorsement or approval of this product by this Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

The labeling referred to above, submitted in connection with registration under FIFRA, is acceptable subject to the comment listed below. Five copies of the finished labeling must be submitted.

- Change "Caution" under the "Directions for Use" section to "Note to User."

Henry H. Jacoby
 Product Manager (21)
 Fungicide-Herbicide Branch
 Registration Division (TS-767C)

BEST AVAILABLE COPY

ATTACHMENT IS APPLICABLE

SIGNATURE OF APPROVING OFFICIAL

DATE

ACTIVE INGREDIENTS	52.0%
Sulfur	52.0%
INERT INGREDIENTS	48.0%
Contains 6 lbs. Sulfur per gallon	TOTAL 100.0%
EPA Reg. No. 7001-261	
EPA Est. No. 7001-CA-1	

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS & DOMESTIC ANIMALS**

Avoid breathing spray mist. Avoid contact with eyes, skin and clothing. Harmful if swallowed, inhaled, or absorbed through skin. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

ENVIRONMENTAL HAZARDS

Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply when weather conditions favor drift from treated areas.

PHYSICAL HAZARDS

Sulfur dust suspended in air ignites easily. Keep away from heat, sparks, or flame. Do not smoke while applying this product.

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED, drink 1 or 2 glasses of water and induce vomiting by sticking finger or blunt object down throat. Do not induce vomiting if victim is unconscious.

IF IN EYES, wash immediately with water for 15 minutes.

IF ON SKIN, wash with soap and water.

RE-ENTRY STATEMENT

Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons. Do not enter treated areas without protective clothing for 24 hours after application. Certain states may require more restrictive re-entry intervals for various crops treated with this product. Consult your State Department of Agriculture for further information.

Written or oral warnings must be given to workers who are expected to enter a treated area or in an area about to be treated with this product. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Written and oral warnings must include the following information: "CAUTION. Area treated with sulfur on (date of application). Do not enter treated areas without appropriate protective clothing for 24 hours following application." Required Protective Clothing: long-sleeved shirt, long-legged pants and gloves.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

STORAGE: Store in a cool, dry area.

DISPOSAL: Completely empty bag into application equipment. Dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

DIRECTIONS FOR DILUTION

Stir until smooth before using. Wash recommended amount through screen into partially filled spray tank or pre-mix in a small amount of water before pouring into tank. Do not use a screen finer than 50 mesh in entire system. Keep agitator running during filling and spraying operations. Do not allow mixture to stand. Failure to maintain agitation will cause sulfur to settle and necessitate manual removal of the deposit. This product should be applied by ground equipment or aircraft in

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

CAUTION: Some plants are susceptible to injury from sulfur under certain climatic conditions. The user is advised not to use sulfur on any crop unless local use has proven that sulfur is safe in that locality.

During periods of high temperature sulfur may burn foliage and fruit. Do not make sulfur applications at excessively high temperatures. Do not use with oil or within 4 weeks of an oil application. When growing crops for processing, consult the processor before applying sulfur.

Tree Crops: Unless otherwise stated for specific crop, dosage rates are given as pints of SULFUR 6 FLOWABLE per 100 gal. water for use in a thorough coverage spray. The total amount of SULFUR 6 FLOWABLE per acre should be 2 to 4 gallons depending on size of trees. Because of spray equipment variations and type used, a range of low and high rates are listed. For high volume sprayers (output 800-1000 gals. spray per acre) use the low rate. For low volume sprayers (output 20-300 gals. spray per acre) use high rate.

Apples: *Apple cedar rust, black rot, frog-eye leaf spot, powdery mildew, quince rust, scab, sooty blotch*—Apply 1½ to 3½ pts. from prepink through calyx periods and ¾ to 2 pts. in cover sprays. **NOTE:** McIntosh, Golden Delicious, Jonathan and certain other varieties may be injured by sulfur application under certain conditions.

Cherries: *Brown rot, leaf spot, powdery mildew, rust, scab and stem blight*—Apply 1½ to 3½ pts. through bloom stage and 2/3 to 1½ pts. from petal fall through cover sprays for brown rot and leaf spot.

Citrus: *Rust mite, clover mite*—Apply ¾ to 4¼ pts. in November through May. *Thrips* (Early coring treatment for the reduction of thrip population)—Apply 1 to 2 gallons per acre in sufficient water for thorough coverage. Make application after spring flush has made 3 to 4 inch growth. Repeat if necessary. Do not apply within 21 days of an oil application.

Grapes: *Powdery mildew, rust (physopella)*—Apply 1 to 2 pts per 100 gal. water using up to 8 pints Super 6 Flowable per acre. Begin when new shoots are 6 to 10 inches long. Repeat before blossoms open and continue at 10 to 14 day intervals as necessary. **NOTE:** Concord and other labrusca type grapes may be injured.

Nectarines, Peaches: *Brown rot, leaf spot, powdery mildew, peach canker, rust, scab, shothole*—Apply 1½ to 3½ pts. through bloom and 2/3 to 1½ pts. in cover sprays. Apply from early petal fall through cover periods as necessary.

Pears: *Brown rot, leaf spot, powdery mildew, rust, scab*—Apply 1½ to 3½ pts. through calyx and ¾ to 2 pts. in cover sprays. Apply from prebloom through calyx periods as necessary.

Plums, Prunes: *Brown rot, leaf spot, powdery mildew, rust, scab, shothole*—Apply 1½ to 7¼ pts. through bloom and ¾ to 2 pts. through cover sprays for brown rot and leaf spot.

Strawberries: *Powdery mildew*—Apply 2 pts. per acre beginning at first sign of disease and repeat as necessary. **NOTE:** Sulfur may injure strawberries under certain climatic conditions.

AGC 479 (4-63)

DISCLAIMER OF WARRANTIES. Seller warrants that the chemical composition of this product conforms to the chemical description given on this label. THIS WARRANTY IS EXPRESSLY IN LIEU OF ALL OTHER WARRANTIES AND REPRESENTATIONS, EXPRESSED, IMPLIED, OR STATUTORY, INCLUDING WARRANTIES OF MERCHANTABILITY AND FITNESS FOR A PARTICULAR USE. Timing, rate and method of application, weather and crop conditions, mixtures with chemicals not specifically recommended on this label or accompanying written recommendation, are beyond the control of seller. Buyer assumes all risks of use, storage and handling of this material not in strict accordance with directions given hereon. Buyer further agrees in the event of damages arising from the use of this product to accept a replacement of the product or a refund of the purchase price of the product, at buyer's option, as full discharge of seller's liability. No one is authorized to make any other warranty, guarantee or directions concerning this product, and no such warranties, guarantees or directions shall