

GUARANTEED ANALYSIS

Total Nitrogen (N)	20.00%
8.00% Ammoniacal Nitrogen derived from Ammonium Sulfate	
12.00% Water Soluble Nitrogen derived from Urea	
Available Phosphoric Acid (P ₂ O ₅) derived from	
Ammonium Phosphate	4.00%
Soluble Potash (K ₂ O) derived from Muriate of Potash	4.00%
Sulfur (S) expressed as elemental derived from	
Ammonium Sulfate	9.00%

Potential Acidity 1250 lbs. Calcium Carbonate Equivalent per ton.

Total Primary Plant Food 28%

BEST DOCUMENT AVAILABLE

DESCRIPTION

1. Best Turf Food Plus Weed Control is especially designed to feed grass lawns and to control broadleaf weeds.
2. Nitrogen, phosphate and potassium are combined in a ratio recommended by turf authorities.
3. Added sulfur helps to overcome deficiencies common in many western soils.
4. Dicamba and 2,4-D effectively control most common lawn weeds, annual and perennial, including:

Black Medic	Honbit	Plantain, Buckhorn
Chickweed, Common	Ground Ivy	Plantain, Common
Chickweed, Mouseear	Knotweed	Sheep Sorrel
Clover	Lambsquarter	Speedwell (Veronica)
Dandelion	Mallow	Spotted Spurge
Dock, Curly	Morningglory	Wild Lettuce
Heal-All	Pigweed	Yarrow

WHEN TO USE

Best Turf Food Plus Weed Control should be used anytime weeds are actively growing. Hormone weed killers are most effective when daytime temperatures are above 60° F. Do not use on newly planted lawns until after the third mowing.

DIRECTIONS FOR USE

1. Use only on grass lawns including Kentucky Bluegrass, Bermudagrass, Fescue, & Ryegrass.
2. Water lawn thoroughly before treatment. Do not water or mow lawn for at least two days after treatment - then resume normal watering.
3. Apply to lawn at recommended rate when lawn is damp (after rain, sprinkling, or dew) so pellets will stick to weeds.
4. Avoid drift to or contact with desirable plants. Do not apply within dripline of trees and shrubs.
5. Do not exceed recommended rate of application which is 5 lbs. per 1000 square feet (20' x 50'), or 217.8 lbs. per acre.

HORTICULTURAL CAUTION: Do not use on Dichondra, St. Augustine, or ground covers. Delay planting until four weeks after treatment.

KEEP OUT OF REACH OF CHILDREN CAUTION

PRECAUTIONARY STATEMENT HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Use only in accordance with label directions. May be fatal if swallowed or inhaled. If swallowed, induce vomiting. Call physician immediately. Avoid contact with skin, eyes, and clothing. Keep pets and children off treated areas until material has been washed into soil and lawn is dry. In case of contact with skin or eyes, flush with plenty of water. For eyes get medical attention. Wash hands with soap and water after handling and before eating or smoking. Wash clothing. Do not apply when weather conditions favor drift from areas treated. Do not apply directly on vegetables, fruits or crops. Do not store near fertilizers, seeds, insecticides or fungicides.

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED - Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. Get medical attention. **IF INHALED** remove person to fresh air. **IF ON SKIN** wash with soap and water. If irritation persists

contact a physician. **IF IN EYES** flush with water for at least 15 minutes. If irritation persists contact a physician.

ENVIRONMENTAL HAZARDS

KEEP OUT OF ANY BODY OF WATER. DO NOT USE WHERE RUNOFF WILL CONTAMINATE STREAMS, LAKES, PONDS, IRRIGATION OR DOMESTIC WATER.

STORAGE AND DISPOSAL

STORAGE: Do not store near fertilizer, seed, feed, food, insecticides or fungicides. Store in a cool, dry, well ventilated area away from heat and open flame.

DISPOSAL: Securely wrap original container in several layers of newspaper and discard in trash.

Read entire label. Use strictly in accordance with label cautions, warnings and directions.

NOTICE: Buyer assumes all responsibility for safety and use not in accordance with directions.



J. R. SIMPLOT COMPANY

MINERALS & CHEMICAL DIVISION

WESTERN REGION

P. O. Box 198 • Lathrop, CA 95330

Lisa Strong
J.R. Simplot Company
16777 South Howland Ave.
P.O. Box 198
Lathrop, CA 95330

JUL 21 1984

Dear Ms. Strong:

Subject: Amendment - Label Revisions
Turf Food Plus Weed Con...
EPA Registration No. 7001-2/2
Your Submission of July 20, 1984

The amendment referred to above submitted in connection with registration under FIFRA is acceptable provided that you:

1. Make the labeling changes listed below before you release the product for shipment:
 - a. Add the human hazard signal word and child hazard statement to front panel.
 - b. Under Hazard to Humans, remove instructions dealing with use restrictions. These are more appropriately placed in the Directions for Use section.
 - c. Under Environmental Hazards, the following language is required: "Do not apply to water. Do not contaminate water by cleaning of equipment or disposal of wastes."
 - d. Under Storage and Disposal, add the following container disposal statement: "Do not reuse bag."
 - e. Under Directions for Use, add: "It is a violation of Federal law to use this product in a manner inconsistent with its labeling."
2. Submit one (1) copy of the final printed labeling before you release the product for shipment. A stamped copy of the labeling is enclosed for your records.

Sincerely yours,

BEST DOCUMENT AVAILABLE

Richard F. Mountfort *RFM*
Product Manager (23)
Fungicide-Herbicide Branch
Registration Division (TS-767)

RD/FHB:Mountfor:DCR-32377:WANG-U073L:eg:Raven:479-2013:7/27/84:Del.6/13/84

CONCURRENCES							
SYMBOL ▶							
SURNAME ▶							
DATE ▶							



TURF FOOD

plus

WEED

CONTROL

20-4-4

BEST DOCUMENT AVAILABLE

- FEEDS GRASS LAWNS
- CONTROLS MOST COMMON BROADLEAF WEEDS
- CONTAINS DICAMBA AND 2,4-D
- 9% SULFUR

TURF FOOD

plus


WEED

CONTROL

20-4-4

BEST DOCUMENT AVAILABLE

NOTE:
Change
7/8/50



- FEEDS GRASS LAWNS
- CONTROLS MOST COMMON BROADLEAF WEEDS
- CONTAINS DICAMBA AND 2,4-D
- 9% SULFUR

ACTIVE INGREDIENTS:

*Dimethylamine Salt of Dicamba (3,6-Dichloro-O-Anisic Acid	0.198%
Dimethylamine Salt of Related Acids	0.022%
**Dimethylamine Salt of 2,4-Dichlorophenoxyacetic Acid	0.687%
INERT INGREDIENTS:	99.153%
TOTAL	100.000%

*Equivalent to 0.11% 3,6-Dichloro-O-Anisic Acid
 **Equivalent to 0.57% 2,4-Dichlorophenoxyacetic Acid

EPA Reg. No. 7001-372
 EPA Est. No. 10835-CA-1

NET WEIGHT 40 LBS.

BEST.

TURF FOOD plus WEED CONTROL 20-4-4

GUARANTEED ANALYSIS

Total Nitrogen (N)	20.00%
8.00% Ammoniacal Nitrogen derived from Ammonium Sulfate	
12.00% Water Soluble Nitrogen derived from Urea	
Available Phosphoric Acid (P ₂ O ₅) derived from	
Ammonium Phosphate	4.00%
Soluble Potash (K ₂ O) derived from Muriate of Potash	4.00%
Sulfur (S) expressed as elemental derived from	
Ammonium Sulfate	9.00%

Potential Acidity 1250 lbs. Calcium Carbonate Equivalent per ton.

Total Primary Plant Food 28%

BEST DOCUMENT AVAILABLE

DESCRIPTION

1. Best Turf Food Plus Weed Control is especially designed to feed grass lawns and to control broadleaf weeds.
2. Nitrogen, phosphate and potassium are combined in a ratio recommended by turf authorities.
3. Added sulfur helps to overcome deficiencies common in many western soils.
4. Dicamba and 2,4-D effectively control most common lawn weeds, annual and perennial, including:

Black Medic	Honbit	Plantain, Buckhorn
Chickweed, Common	Ground Ivy	Plantain, Common
Chickweed, Mouseear	Knotweed	Sheep Sorrel
Clover	Lambsquarter	Speedwell (Veronica)
Dandelion	Mallow	Spotted Spurge
Dock, Curly	Morningglory	Wild Lettuce
Heal-All	Pigweed	Yarrow

WHEN TO USE

Best Turf Food Plus Weed Control should be used anytime weeds are actively growing. Hormone weed killers are most effective when daytime temperatures are above 60° F. Do not use on newly planted lawns until after the third mowing.

DIRECTIONS FOR USE

1. Use only on grass lawns including Kentucky Bluegrass, Bermudagrass, Fescue, & Ryegrass.
2. Water lawn thoroughly before treatment. Do not water or mow lawn for at least two days after treatment - then resume normal watering.
3. Apply to lawn at recommended rate when lawn is damp (after rain, sprinkling, or dew) so pellets will stick to weeds.
4. Avoid drift to or contact with desirable plants. Do not apply within dripline of trees and shrubs.
5. Do not exceed recommended rate of application which is 5 lbs. per 1000 square feet (20' x 50'), or 217.8 lbs. per acre.

HORTICULTURAL CAUTION: Do not use on Dichondra, St. Augustine, or ground covers.
Delay planting until four weeks after treatment.

KEEP OUT OF REACH OF CHILDREN