

ACCEPTED

APR 1 1976

Under the Federal Insecticide, Fungicide, and Rodenticide Act, registered...

7001-276

Hi-Yield Livestock Spray

For Use Against Lice, Ticks, Horn Flies, Mosquitoes, Sarcopitic Mange, and other pests on Cattle, Sheep, and Goats.

ACTIVE INGREDIENTS:
 Cypermethrin (0.05%)
 Deltamethrin (0.05%)
 Malathion (0.05%)
 Permethrin (0.05%)
 Pyrethrin (0.05%)
 Pyrethrum (0.05%)
 Resmethrin (0.05%)
 Tetrahydrocannabinol (0.05%)
 Zeta-cypermethrin (0.05%)

OTHER INGREDIENTS:
 Water, Perfume, Propellant, and other inert ingredients.

WARNING:
 KEEP OUT OF REACH OF CHILDREN
 See Side Panel for Additional Warnings

NET CONTENTS ONE GALLON

PRECAUTIONS IN USING THIS PRODUCT: Do not treat calves under 1 month of age. Do not dip or spray animals within 28 days of slaughter. Do not dip or spray within 2 weeks of freshening. Do not dip or spray animals when weather is cold or stormy as injury may result from chilling. Do not dip or spray overheated, thirsty or sick animals. DO NOT USE ON DAIRY ANIMALS IN DAIRY BARN, MILK ROOMS OR POULTRY HOUSES. DO NOT USE ON LACTATING DAIRY ANIMALS OR GOATS. DO NOT SPRAY DAIRY ANIMALS UNDER 3 MONTHS OF AGE.

AS A SPRAY:

CATTLE: Lice, Ticks, Horn Flies, Mosquitoes, Sarcopitic Mange — Mix 1 cup per 5 gal. of water (or 2 1/4 pints per 25 gal. of water). Spray animal thoroughly; do not overspray. A spray of about 2 quarts of dilute spray per adult animal affords protection for 2-3 weeks. Repeat as necessary. For lice one treatment will usually be sufficient for a season. Do not use on dairy cattle.

HOGS: Lice — Mix 1 cup per 5 gal. of water (or 2 1/4 pints per 25 gal. of water). Spray animal thoroughly. One treatment is usually sufficient for a season.

SHEEP AND GOATS: Lice and Sheep Keds (ticks) — Mix 1 1/3 tablespoons per gal. of water (or 3/8 cup for 5 gal. of water). Spray animal thoroughly using high pressure in order to penetrate fleeces. Ticks: Mix 1 3/5 tablespoons per gal of water (or 5 tablespoons per 5 gal. of water). Spray thoroughly as above. Repeat at 2-3 weeks or as necessary.

AS A DIP:

For Control of Listed Pests on Cattle: Use 1 gal. HI-Yield LIVESTOCK SPRAY NO. 2 per 100 gals. water. Fill vat first with water, then add required amount of HI-Yield LIVESTOCK SPRAY NO. 2. Mix thoroughly for 3 minutes or longer. Any time dipping operation is stopped for 30 minutes, agitate thoroughly before resuming dipping. Completely submerge animal's body once or twice. Duck head at least twice. Do not use on calves or other young animals under three months of age. Do not use on dogs, cats or other household pets or poultry. Do not dip dairy cattle.

7001-276

4/11/1976

Page 1 of 1