


DURSBAN®* 2.5% G GRANULAR INSECTICIDE

Controls chinch bugs, sod webworms (lawn moths), ants, brown dog ticks, crickets, cutworms, earwigs, sowbugs (pillbugs), and hyperodes (turfgrass) weevils infesting home lawns and other ornamental turf areas.

| | |
|---|---------------|
| ACTIVE INGREDIENT: | BY WT. |
| Chlorpyrifos (O,O-diethyl O-[3,5,6-Trichloro-2-pyridyl] phosphorothioate)** | 2.5% |
| INERT INGREDIENTS: | 97.5% |
| TOTAL | 100.0% |

*Dursban Trademark of the Dow Chemical Company
**U.S. Patent No. 3,244,586

EPA Reg. No. 7001-170-AA
EPA Est. No. 7001-CA-1



DIRECTIONS FOR USE

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS, WARNINGS AND DIRECTIONS.

Apply Dursban 2.5% Granular Insecticide uniformly to home lawns and other ornamental and recreational turf grass areas using any hand or power operated gravity or rotary type spreader designed for application of granular materials. Treat when insects first appear using Dursban 2.5% Granular at the rate of 40 lbs. per acre. To help obtain uniform distribution, apply one-half of the required amount of product in one direction; then apply the balance at a right angle. Immediately after application, sprinkle the turf with 1300 gallons of water per acre so as to wash the granules into the zone of insect infestation. Consult the following table to determine the amount of Dursban 2.5% Granular and water needed for smaller sized areas. Re-treat as necessary.

| AREAS TO BE TREATED | RATES TO USE LBS. OF DURSBAN 2.5% G | GALLONS OF WATER TO SPRINKLE ON TURF AFTER TREATMENT |
|---------------------|--|--|
| 1,000 sq. ft. | 1 | 30 |
| 5,000 sq. ft. | 5 | 150 |
| 10,700 sq. ft. | 10 | 325 |

***To control hyperodes (turfgrass) weevils, use twice the rate of Dursban 2.5% Granular shown in the table. Treat suspected problem areas in mid-April and again in mid-May. Consult your local Agricultural Extension Service specialist for best time to treat in your area.

KEEP OUT OF REACH OF CHILDREN WARNING

MAY BE HARMFUL IF SWALLOWED

Keep children and pets off treated areas until this material is washed into the soil and grass is dry.

This product is toxic to fish, birds, and other wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, streams or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

Dispose of empty container and wastes by burning or by burying in non-crop lands away from domestic water supplies.

Do not take internally. Avoid eye contact, flush contaminated eyes with plenty of water and get medical attention if irritation persists. Wash thoroughly after handling. Wash contaminated clothing before re-use. Keep away from feed, foodstuffs and domestic water supplies.

7001-276

DIRECTIONS FOR USE

Registered under
EPA Reg. No. 71127

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS, WARNINGS AND DIRECTIONS.

Apply Dursban 2.5% Granular Insecticide uniformly to home lawns and other ornamental and recreational turf grass areas using any hand or power operated gravity or rotary type spreader designed for application of granular materials. Treat when insects first appear using Dursban 2.5% Granular at the rate of 40 lbs. per acre. To help obtain uniform distribution, apply one-half of the required amount of product in one direction; then apply the balance at a right angle. Immediately after application, sprinkle the turf with 1300 gallons of water per acre so as to wash the granules into the zone of insect infestation. Consult the following table to determine the amount of Dursban 2.5% Granular and water needed for smaller sized areas. Re-treat as necessary

| AREAS TO BE TREATED | RATES TO USE LBS. OF DURSBAN 2.5% G | GALLONS OF WATER TO SPRINKLE ON TURF AFTER TREATMENT |
|---------------------|--|--|
| 1,000 sq. ft. | 1 | 30 |
| 5,000 sq. ft. | 5 | 150 |
| 10,700 sq. ft. | 10 | 325 |

***To control hyperodes (turfgrass) weevils, use twice the rate of Dursban 2.5% Granular shown in the table. Treat suspected problem areas in mid-April and again in mid-May. Consult your local Agricultural Extension Service specialist for best time to treat in your area.

37

KEEP OUT OF REACH OF CHILDREN WARNING

MAY BE HARMFUL IF SWALLOWED.

Keep children and pets off treated areas until this material is washed into the soil and grass is dry.

This product is toxic to fish, birds, and other wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, streams or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

Dispose of empty container and wastes by burning or by burying in non-crop lands away from domestic water supplies.

Do not take internally. Avoid eye contact, flush contaminated eyes with plenty of water and get medical attention if irritation persists. Wash thoroughly after handling. Wash contaminated clothing before re-use. Keep away from feed, foodstuffs and domestic water supplies.

NOTE TO PHYSICIAN: Active ingredient is cholinesterase inhibitor. Treat symptomatically. Atropine is an antidote.

NET WEIGHT 35 POUNDS

38

38 (16/78)