



# LINDANE BORER SPRAY

CONTAINS LINDANE

Kills Pine Bark Beetles, Peach Tree Borers, Flat Headed Borers, Juniper Twig Girdler, Leaf Miners

## ACTIVE INGREDIENTS:

Lindane (Gamma isomer of benzene hexachloride) ..... 20.0%  
Xylene ..... 67.6%

## INERT INGREDIENTS:

EPA Reg. No. 7001-211-AA  
EPA Est. No. 7001-CA-1

TOTAL

## BY WT.

20.0%  
67.6%  
12.4%

100.0%

Read entire label Use strictly in accordance with label cautions, warnings and directions

Lindane Borer Spray is formulated for the control of boring type insects, as well as certain insects attacking greenhouse and nursery plants. Because timing is important in controlling certain borers, contact your State Agricultural Experiment Station or State Extension Service Specialist for preferred timing in your area. Water solutions may be made with any type of accurate garden sprayer. Oil or kerosene dilutions are best applied with a compressed air tank type sprayer.

Do not use, pour, spill, or store near heat or open flame.

## CAUTION

KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed. Do not breathe vapor or spray mist. Avoid contact with skin. May be absorbed through skin. In case of contact wash immediately with soap and water.

Avoid contamination of feeds and foodstuffs. Do not use on household pets or humans. Keep children and pets away from treated areas until these areas are dry.

Do not apply to foliage of white or sweet potatoes. Do not reuse container. Destroy when empty. Keep pesticide in original container. Never refuel or mix with other pesticides in this container.

## ENVIRONMENTAL CAUTION

This product is toxic to fish, birds, and other wildlife. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

This product is toxic to bees and other beneficial insects. Do not apply directly to blooming plants.

## DIRECTIONS FOR USE

Do not use diluted spray. Make new dilution for each treatment. Do not spray in water unless specifically directed. Do not use in water.

**Peaches, Apricots, Cherries, Plums:** Peach Tree Borer. Apply 1 tablespoon/gal. (3 pts./100 gals.) Spray lower trunk and around base of tree soaking bark and wetting trunk from crotch to ground. Do not apply to fruit or foliage. Apply in late spring or early summer for best results. Do not apply within 60 days of harvest.

**Pines:** Pine Bark Beetles. Apply 8 oz. in 2 1/2 gals. water, kerosene, or No. 2 fuel oil. Spray trunks thoroughly. Wet trunks to runoff.

**Apples, Cherries, Peaches:** Fruit Tree Bark Beetles, Flat Headed Borers, Round Headed Borers. Apply 1 tablespoon/gal. (3 pts./100 gal.) Make applications during appearance of moths in early to late summer. Apply 2 to 3 sprays to trunk from ground to scaffold limbs timed with moth flight. Do not apply to fruit or foliage. Do not apply within 60 days of harvest.

**Junipers:** Twig Girdler. Apply 1 tablespoon/gal. Make 2 applications at 30 day intervals. Northern California: June and mid-July. Southern California: April and May.

**Maple, Aspen, Poplar, Ash, Willows, Elm, Locust:** Bark Beetles, Wood Boring Beetles. Apply 2 tablespoons/gal. (3 pts./50 gal.) Apply in spring when buds begin to break. Repeat in 10 to 14 days. Elms: repeat in fall. Soak trunks to runoff.

**Birch, Elm, Locust:** Leaf Miners. Apply 1 teaspoon/gal. Apply when small miners appear in early and mid-May. For late season early July.

**Rhododendron, Lilac, Dogwood:** Dogwood Borer, Rhododendron Borer, Lilac Borer. Apply 1 tablespoon/gal. (3 pts./100 gal.) Make 2 to 3 applications 7 to 10 days apart in May and early June.

**Taxus, Hemlock:** Taxus Weevil. Apply 1 tablespoon/gal. Apply in May and early June around base of tree, soaking the bark and wetting the soil surface.

**Greenhouse & Nursery Plants (Roses, Camellias, Azaleas, Gardenias, Chrysanthemums, Gladiolus, Camellions, Snap Dragons, Rhododendrons, Primrose):** Aphids, Thrips, Cabbage Loopers, Plant Bug, Whitefly, Psyllid, Leaf Miners, Flea Beetles, Diabrotica Beetles. Apply 1 teaspoon/gal. (1 pt./100 gal.)

**Camellias, Azaleas, Rhododendrons:** Fuller Rose Weevil, Brachymeris, Cucumber Beetle Larvae on roots. Apply 1 teaspoon/gal. (1 pt./100 gal.) Spray to saturate soil around plants in beds. For potted plants spray soil surface lightly comparable to a light rain.

(GL-22:1/76)

DISCLAIMER OF WARRANTIES: Seller warrants that the chemical composition of this product conforms to the chemical description given on this label. THIS WARRANTY IS EXPRESSLY IN LIEU OF ALL OTHER WARRANTIES AND REPRESENTATIONS EXPRESSED OR IMPLIED OR STATUTORY, INCLUDING WARRANTIES OF MERCHANTABILITY AND FITNESS FOR A PARTICULAR USE. Timing, rate and method of application, weather and crop conditions, mixtures with chemicals not specifically recommended on this label or an accompanying written recommendation are beyond the control of seller. Buyer assumes all risks of use, storage and handling of this material not in strict accordance with directions given on this label. Seller assumes no liability for damages arising from the use of this product in any manner not recommended on this label.

## Occidental Chemical Company

P.O. Box 198 Lathrop, California 95330