



ACCEPTED Tomato & Vegetable Dust

DEC 9 1983
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under 7001-310
EPA Reg. No.

BEST DOCUMENT AVAILABLE

CONTAINS
Sevin* • Malathion • Maneb

Controls insects and diseases attacking tomatoes and vegetables
TO APPLY JUST SQUEEZE CONTAINER.

ACTIVE INGREDIENTS:
Malathion (O,O-dimethyl dithiophosphate of diethyl mercaptosuccinate) 5%
Carbaryl (1-naphthyl methylcarbamate) 5%
Maneb (manganese ethylene bisdithiocarbamate) 6%
INERT INGREDIENTS 84%

*Sevin is the registered trade mark of Union Carbide Corporation for carbaryl insecticide

EPA Reg. No. 7001-310
EPA Est. No. 7001-CA-1

KEEP OUT OF REACH OF CHILDREN
CAUTION

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Harmful if inhaled or swallowed. Do not breathe dust. Do not get in eyes or on skin or clothing. Wash thoroughly with soap and water after using. Wear the following protective clothing during application: long pants, long-sleeved shirt, gloves, hat and boots.

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED drink warm water and induce vomiting by touching back of throat with finger or blunt object. Antidote is Atropine Sulfate. Never give fluids or induce vomiting if person is unconscious or having convulsions. Get medical attention.

IF INHALED remove victim to fresh air.

IF IN EYES flush with water for at least 15 minutes. Get medical attention.

IF ON SKIN remove contaminated clothing and flush with water. Wash exposed areas thoroughly with soap and water.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Keep out of lakes, streams and ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

PHYSICAL OR CHEMICAL HAZARDS

Do not use on Boston, Maidenhair and Pteris Ferns, also some species of Crassula. Do not allow this product to drift onto adjoining food or feed crops. Residue may contaminate these crops making them unfit for human consumption.

STORAGE AND DISPOSAL

STORAGE Store in cool, dry place. Do not store near food or feed. Keep container tightly closed.

DISPOSAL Securely wrap original container in several layers of newspaper and discard in trash. Do not reuse container.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Read entire label before using. Use strictly according to directions and cautions. Open protective cap, point opening away from you and squeeze. Apply in short puffs in sweeping motion. Cover foliage lightly but do not over apply. Make certain that upper and lower leaf surfaces are thoroughly covered. Shake container frequently to keep dust free flowing and prevent packing. () circled number indicates days between last use and harvest.

CROP	PEST OR DISEASE	
Tomatoes (5)	Insects: Aphids, Colorado Potato Beetle, Two Spotted Mite, Tomato Pinworm, Drosophila, Tomato Fruit Worm, Tomato Hornworm, European Corn Borer, Sink Bugs, Flea Beetles, Leafhoppers, Armyworm. Diseases: Early Blight, Late Blight, Leaf Spot, Anthracnose.	For disease control begin at first blight cluster and repeat at 7-10 day intervals.
Beans (4)	Insects: Aphids, Leafhoppers, Jugs Bugs, Bean Leaf Beetle, Sink Bugs, Armyworms, Corn Earworms, Flea Beetles, Cucumber Beetles, Two Spotted Mites. Diseases: Downy Mildew, Rust, Anthracnose.	For disease control begin when plants are small and repeat at 5-7 day intervals.
Sweet Corn (3)	Corn Earworm.	Apply when silks first appear. Repeat at 2-3 day intervals during silking period. Do not feed or graze corn forage.
Broccoli, Brussels Sprouts, Cabbage, Cauliflower	Insects: Aphids, Leafhoppers, Sink Bugs, Armyworms, Diamond Back Moth, Corn Earworms, Flea Beetles, Imported Cabbage Worms, Turnip Root Bug, Cabbage Looper. Diseases: Downy Mildew, Alternaria Leaf Spot.	Begin when disease threatens and repeat at 4 day intervals.
Lettuce, Cornichon	Insects: Aphids, Leafhoppers, Jugs Bugs, Flea Beetles, Sink Bugs, Armyworms, Corn Earworms, Cabbage Looper, Turnip Root Bug. Diseases: Downy Mildew.	Remove residues from next lettuce or chard, and remove and burn old lettuce and chard. Wash or blanch before eating. Begin when disease threatens and repeat at 7-10 day intervals.
Cucumbers, Melons, Watermelons, Pumpkins	Insects: Aphids, Leafhoppers, Sink Bugs, Cucumber Beetles, Flea Beetles, Cucumber Beetles, Melonworms, Squash Vine Borer, Two Spotted Mites. Diseases: Downy Mildew, Anthracnose, Leaf Spot, Mosaic, Virus, Bacterial Wilt, Cucumber Mosaic, Watermelon Bacterial Wilt.	Begin when disease threatens and repeat at 4 day intervals.

NET WEIGHT: 10 OUNCES.

J. R. SIMPLOT COMPANY
MINERALS & CHEMICAL DIVISION
WESTERN REGION
P.O. Box 198 • Lathrop, CA 95330