



OXY WEED and GRASS KILLER, GRANULAR

OXY WEED and GRASS KILLER, GRANULAR NONSELECTIVE
U.S. Patents 2700604 and 3464810

ACTIVE INGREDIENTS:

Sodium Chlorate (NaClO_3) 30.00%
Sodium Metaborate Tetrahydrate
 $\text{Na}_2\text{B}_4\text{O}_7 \cdot 4\text{H}_2\text{O}$ 66.50%

Boron Trioxide (B_2O_3) Equivalent 22.7%

Diuron [3-(3,4-Dichlorophenyl)-1,1-dimethylurea] 1.25%
INERT INGREDIENTS: 2.25%

TOTAL 100.00%

EPA Reg. No. 7001-337

EPA Est. No. 10659-MS-1

KEEP OUT OF REACH OF CHILDREN DANGER

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Harmful if swallowed. May irritate eyes, nose, throat and skin. Avoid contact with skin, eyes and clothing. Avoid breathing dust or spray mist. Wash thoroughly after handling. Remove and wash contaminated clothing before reuse. Causes eye damage. Wear goggles or face shield when handling.

Do not apply to areas likely to be grazed by domestic animals. Keep animals off treated areas until rainfall has leached herbicide into soil.

STATEMENT OF PRACTICAL TREATMENT

IF IN EYES flush with water for 15 minutes. Get medical attention. IF ON SKIN wash with plenty of soap and water. Get medical attention if irritation persists.

ENVIRONMENTAL HAZARDS

Keep out of lakes, streams and ponds. Do not apply when weather conditions favor drift from target area. Do not contaminate water by cleaning of equipment or disposal of waste.

STORAGE AND DISPOSAL

Pesticide, spray mixture or rinse water that cannot be used according to label instructions must be disposed of according to Federal or approved state procedures under Subtitle C of the Resource Conservation and Recovery Act.

Completely empty bag by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment. Dispose of bags in a sanitary landfill.

DIRECTIONS FOR USE

GENERAL CLASSIFICATION

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

OXY WEED and GRASS KILLER is specifically designed for quick knock-down and control of general mixed vegetation and such perennial grasses as Johnson grass, Dallis grass and reedgrass (*Phragmites*) on noncropped land industrial sites. OXY WEED and GRASS KILLER may provide season long control of seedlings on treated areas. Soil type, rainfall and length of growing season will influence the residual control of seedlings.

APPLICATION: For all uses uniform application is essential. OXY WEED and GRASS KILLER may be applied either as a dry granular material or as a water spray. Dissolve in water at rates up to 3 pounds per gallon. Both spray and dry applications become effective as soil treatments following rainfall or irrigation.

For DRY application, conventional herbicide spreaders may be used. The hand-operated cyclone type is especially recommended for use where power equipment is not desired.

For SPRAY application use equipment with mechanical agitation. Add as much as 3 pounds OXY WEED and GRASS KILLER per gallon of water, depending on job requirements to spray tank. Continue agitation until the herbicide is completely dissolved and during the spraying operation. To minimize fire hazard due to dry vegetation, treatment while weeds are small is suggested.

GENERAL VEGETATION CONTROL: OXY WEED and GRASS KILLER will give quick knock-down of vegetation and residual control of broadleaved weeds and grasses on parking lots, fence lines, highway and railroad sites, storage yards and other similar areas. For spray applications, add 1/2 to 2 pounds of OXY WEED and GRASS KILLER in sufficient water to cover 100 square feet and apply as a contact spray thoroughly wetting all parts of the plants, particularly the growing tips—1/4 to 1 gallon per 100 square feet is usually adequate for young growth up to 6 inches tall.

The higher rates may be necessary to insure thorough coverage of heavy vegetation or to provide extended residual control. For DRY granular treatment, apply 1/2 to 2 pounds per 100 square feet. The lighter rates should be used only at time of weed emergence on moist soil or when precipitation is anticipated while weeds are still young. Persistent perennial weeds are most effectively treated when nearing maturity or during dormant stages in fall. Spring applications may also be made under moderate rainfall conditions. Summer applications are recommended only where rainfall is dependable for leaching herbicide into soil. Apply 2 to 4 pounds per 100 square feet in the dry granular form or as a spray. On deep-rooted perennials such as bindweed, Canada thistle, Bermuda grass and quackgrass the heavier rates may be required.

JOHNSONGRASS CONTROL: OXY WEED and GRASS KILLER is particularly effective on deep-rooted perennial grasses such as Johnson grass and Dallis grass on industrial sites, roadsides, fence lines and storage yards. Apply dry at 1 to 3 pounds per 100 square feet or add 1 to 3 pounds to 1 to 2 gallons of water to adequately cover 100 square feet. Maximum efficiency on Johnson grass is obtained by treating after heading to late fall. IMPORTANT: This material is not selective in its action, may be toxic to all types of vegetation and may render the entire treated area totally or partially unproductive for one or more years. Care should be taken to confine the use or application to the particular area intended to be treated and to avoid its contact with lawns, trees, shrubs, crops and other desirable plants which are not intended to be destroyed or injured. This includes precautions in treating areas which may be underlaid by roots of adjacent valuable growths. Careless application of this material or washing by water run-offs to areas where desirable plants are growing or which will be used for later planting may result in injury to desirable plants. Do not drain or flush equipment on or near such areas.

Because of the many varying conditions affecting use and application, such as location, seasonal weather, soil composition and type of vegetation, manufacturer warns buyer that these and other conditions may impair or vary the results obtained from the use of this product. In any event complete elimination of weeds is not guaranteed. It is advisable to consult your State Extension Service or Agricultural Experiment Station weed specialists on specific problems.

CONDITIONS OF SALE: 1. Seller warrants that this product consists of the ingredients specified and is reasonably fit for use as directed on the label. No one, other than officer of Seller, is authorized to make any warranty, guarantee or direction concerning this product. 2. Because the time, place, rate of application and other conditions of use are beyond Seller's control, Seller's liability from handling, storage and use of this product is limited to replacement of product or refund of purchase price.

(5/10/82)

MANUFACTURED BY:

ACCIDENTAL CHEMICAL CORPORATION
P.O. Box 198, Lathrop, CA 95330

NET CONTENTS 50 LBS.