


INSECTICIDE

Diazinon W-50 is a wettable powder.

Active Ingredient:

O, O-diethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl)
phosphorothioate 50%
Inert Ingredients: 50%
Total: 100%



WARNING — PELIGRO

KEEP OUT OF REACH OF CHILDREN

PRECAUCION AL USUARIO: Si usted no lee ingles, no use este producto hasta que le entregue haya explicado ampliamente

EPA Reg. No. 7001-349
EPA Est. No. 7001-CA-1

May be fatal if swallowed. May be absorbed through skin. Do not breathe dust or spray mist. Avoid contact with skin, eyes, and clothing. Wash thoroughly after handling. Avoid contamination of food and feed. Keep out of reach of domestic animals. Do not use on humans or household pets. Do not permit children or pets to go on to sprayed grass until spray has completely dried.

Fish and Wildlife Caution:

This product is toxic to fish, and wildlife. Birds feeding on treated areas may be killed. Keep out of any body of water. Do not apply where runoff is like

ly to occur. Do not contaminate water by cleaning of equipment, or disposal of wastes. Do not apply when weather conditions favor drift from areas treated. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where these are important resources. Apply this product only as specified on this label.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

AGC 53319-841

**SEE ATTACHED PACKET FOR ADDITIONAL
PRECAUTIONARY STATEMENTS AND DIRECTIONS FOR USE!**

10/5

Critical F₁
(Graph these)

2000年1月
 2000年1月
 2000年1月

Civil Service
No. 10

Do not a
Grapes

and also not appl

01/04/2004

Do not
Peel the

Sat. 20
Oct 1941

Adm. Serv.
Crim. Div.
Off. Sec.
Training

Oriente
L'Europa

Pouch 1.

continuer

longer

Page

기후변화 대응

Figure 1

Fig. 2. 2.3.4

人 生 之 道

Prüfungsausschuss

Principles

Peach 1

Bryan M

Signature

C. J. C. J.

4000 LTR
4000 LTR

1. **Wahlberechtigte** (eligible voters)
 2. **Wahlberechtigte** (eligible voters)

Figure 4

Figure 1

Figure 2.

20/5

Beans (Pole Beans, Snap Beans, Lima Beans, Black Bean Aphids, Mexican Bean Beetles, Leaf Cucumber Beetles. Apply 1/4 to 1/2 lb in 110 gal. Do not apply within 7 days of bean harvest. Beans: Wait 4 days following treatment if forage.

Lima Beans
Dipterous Leaf Miners, Leafhoppers, Mites. Apply Bean forage or bean hay may be fed to dairy cattle forage is to be cut for hay.

Broccoli, Cabbage, and Cauliflower
Aphids, Diamondback Moths, Imported Cabbageworms. Apply 1/4 to 1/2 lb in 110 gal. Do not apply within 7 days of harvest.

Brussels Sprouts
Aphids, Diamondback Moths, Imported Cabbageworms. Apply 1/4 to 1/2 lb in 110 gal. Do not apply within 7 days of harvest.

Carrots
Aphids. Apply 1 lb per acre. Do not apply within 7 days of harvest.

BEST AVAILABLE COPY

Beans (Pole Beans, Snap Beans, Lima Beans)
Black Bean Aphids, Mexican Bean Beetle, Le
Cucumber Beetles. Apply 1/2-1 lb. in 10 gal.
Do not apply within 7 days of bean harvest. Bee
sheep. Wait 4 days following treatment for forage.

Lima Beans
Dipterous Leaf Miners, Leafhoppers, Mites. For
Bean forage or bean hay may be fed to cattle
forage is to be cut for hay.

Broccoli, Cabbage, and Cauliflower
Aphids, Diamondback Moths. Imported Cabbag
pickworms at 7-10 day intervals until usually be
or to cabbage within 7 days of harvest.

Brussels Sprouts
Aphids, Diamondback Moths. Imported Cabbag
worms and usually be required. Do not apply to b

Carrots
Aphids. Apply 1 lb. per acre. Do not apply with

3

Beans (Pole Beans, Snap Beans, Lima Beans)
Black Bean Aphid, Mexican Bean Beetle, Leaf Miner, Armyworm, Flea Beetle
Cucumber Beetles: Apply 1/4 lb. in 100 gals. of water per acre.
Do not apply within 7 days of bean harvest. Beans forage & beans may be fed to dairy and beef cattle and sheep. Wait 4 days following treatment if forage is to be cut for hay.

Lima Beans
Dipterous Leaf Miners, Leafhoppers, Mites: Apply 1/4 lb. per acre. Do not apply within 7 days of bean harvest. Bean forage or bean hay may be fed to dairy and beef cattle and sheep. Wait 4 days following treatment if forage is to be cut for hay.

Broccoli, Cabbage, and Cauliflower
Aphids, Diamondback Moths, Imported Cabbageworms, Dipterous Leaf Miners: Apply 1/4 lb. per acre. Applications at 7-10 day intervals will usually be required. Do not apply in the field or cut for hay within 10 days or to cabbage within 7 days of harvest.

Brussels Sprouts
Aphids, Diamondback Moths, Imported Cabbageworms: Apply 1/4 lb. per acre. Applications at 7-10 day intervals will usually be required. Do not apply in the field or cut for hay within 7 days of harvest.

Carrots
Aphids: Apply 1 lb. per acre. Do not apply within 10 days of harvest.

Celery
Aphids, Flea Beetles: Apply 1 lb. in 100 gals. of water per acre.
Dipterous Leaf Miners: Apply 1/4 lb. per acre. Sufficient water to achieve thorough coverage when in hot weather is necessary. Repeat as necessary.
Do not apply within 10 days of harvest. Plants should be market trimmed, high rendered before being sold or used. Tops should not be used for feed or food.

Collards, Kale, Swiss Chard, and Turnip Tops
Aphids, Imported Cabbageworms, Diamondback Moths, Harlequin Cabbage Bugs, Cane Earworms: Apply 1/4 lb. per acre. Do not apply to collards, kale or turnip tops within 10 days or to Swiss chard within 12 days of harvest.

Cucumbers
Aphids, Serpentine Leaf Miners, Thrips: Apply 1 lb. per acre. Do not apply within 7 days of harvest.

Lettuces and Endives
Aphids, Dipterous Leaf Miners: Apply 1/4 lb. per acre. Do not apply within 10 days of harvest.

Malons and Squash (Cantaloupes, Casabas, Crenshawes, Honeydew Malons, Muskmelons, Persian Melons, and hybrids of these, Watermelons and their hybrids, Summer and Winter Squash)
Aphids, Leafhoppers, Mites: Apply 1/4 lb. per acre.

Serpentine Leaf Miners, Thrips: Apply 1/4 lb. per acre.
Do not apply to melons or winter squash within 3 days or to summer squash within 7 days of harvest.

Mushroom Houses
Mushroom Flies (Phorids and Scandals): Mix 4-8 lbs. in 50 gals. of water (0.5% to 1.0% actual Daznol) and spray on outside and inside walls and ventilators of mushroom houses after compost in beds has been pasteurized by heating. Do not spray compost in beds or after spawning.
To treat wooden sides of beds, posts, doors and door frames after pasteurizing mix 1 lb. in 6 gals. of water and apply with a brush.
Do not treat compost.

Onions
Onion Thrips: Apply 1 lb. per acre. Do not apply within 10 days of harvest.

Peas
Aphids, Leaf Miners: Apply 1/4 lb. per acre. Peas may be picked immediately. Pea vines and pea vine hay may be fed to dairy and beef cattle and sheep. Wait 4 days following treatment if forage is to be cut for hay.

Peppers
Aphids, Serpentine Leaf Miners: Apply 1/4 lb. per acre. Do not apply within 5 days of harvest.

Potatoes
Aphids, Flea Beetles, Colorado Potato Beetles, Dipterous Leaf Miners: Apply 1/4 lb. per acre.
Banded Cucumber Beetles, Leafhoppers, Southern Armyworms: Apply 1/4 lb. per acre.
Do not apply within 35 days of harvest.

Radishes, Parsnips, and Turnips
Aphids, Flea Beetles, Dipterous Leaf Miners: Apply 1/4 lb. per acre. Do not apply within 10 days of harvest.

Spinach and Beets
Aphids, Dipterous Leaf Miners: Apply 1/4 lb. per acre. Do not apply to spinach within 10 days or to beets within 14 days of harvest.

Squash (Summer and Winter)
See Malons and Squash above.

Sweet Corn
Corn Earworms: Apply 2-2 1/2 lb. per acre when silks first appear. Two or three repeat applications at 2-3 day intervals will usually be necessary. Corn may be picked immediately following last application. Corn forage may be fed to dairy and beef cattle and sheep.
Refer also to Corn under Field and Forage Insects.

Tomatoes
Aphids, Dipterous Leaf Miners: Apply 1/4 lb. per acre. To control leaf miners, weekly applications are usually required.

Banded Cucumber Beetles, Fat Armyworms, Southern Armyworms: Apply 1/4 lb. per acre.
Vinegar Flies (Drosophila spp.): Apply 1/4 lb. per acre. Weekly applications are usually required.
Do not apply within 1 day of harvest.

Watermelons (Harvest Only)
Caterpillar Mites: Apply 1 lb. in 100 gals. of water per acre. Direct spray to crown and growing tops of plants. If infestation is heavy, make a second application later than 10 days after the first application. Repeat applications as necessary. Do not apply within 5 days of harvest.

Vegetable Soil Insects

Daznol W 50 should always be applied as a spray in sufficient water to assure thorough coverage of the soil.

Carrot Root Flies
Carrots: Apply 4 lb. per acre as a drench to the seed furrow at planting time.

Cucumbers (Surface and Subterranean)
Broccoli: Collards, Mustard Greens, Summer Squash
Brussels: Cucumbers, Parsley, Sweet Corn
Sprouts: Endive (escarole), Peas, Swiss Chard
Cabbage: Kale, Peppers, Tomatoes
Cantaloupes: Lettuce, Potatoes, Watermelons
Cauliflower: Lima Beans, Snap Beans, Winter Squash
Celery: Spinach

Broadcast 4-8 lb. per acre just prior to planting. Work into soil 4-8 inches immediately. 2-3 inches for surface cut worms, 3-6 inches for subterranean cutworms.

Garden Symphylans (Centipedes)
Cabbage: Peas, Snap Beans
Carrots: Pole Beans, Tomatoes
Corn: Radishes, Turnips
Lettuce: Red Beets

Broadcast 20 lb. per acre just prior to planting. Work into soil 1-2 inches immediately. This prevent treatment will also control other soil insects such as cutworms, burrows and a few species of mole crickets, mole maggots and wireworms.

Mole Crickets
Broccoli: Collards, Peppers
Brussels: Endive (escarole), Radishes
Sprouts: Kale, Tomatoes
Cabbage: Lettuce, Turnips
Cauliflower: Mustard

Broadcast 2 lb. per acre when insects are present. Make application to plant beds or fields 1-2 days prior to sowing of seed or transplanting and work into soil 1-2 inches immediately.

Onion Maggots
Onions: Follow Application, Apply 2 lbs. per acre in sufficient water to cover the seed furrow at planting time.

Broadleaf Awe: Apply 3-4 lb. per acre just prior to planting. Work into soil 3-4 inches immediately.

Root Maggots
Broccoli: Apply 4-8 lb. per acre just prior to planting. Work into soil 4-8 inches immediately. 2-3 inches for surface cut worms, 3-6 inches for subterranean cutworms.
Cucumbers (Surface and Subterranean): Broadcast 4-8 lb. per acre just prior to planting. Work into soil 4-8 inches immediately. 2-3 inches for surface cut worms, 3-6 inches for subterranean cutworms.
The Sweet Corn Earworm: Apply 2-2 1/2 lb. per acre when silks first appear. Two or three repeat applications at 2-3 day intervals will usually be necessary. Corn may be picked immediately following last application. Corn forage may be fed to dairy and beef cattle and sheep. Wait 4 days following treatment if forage is to be cut for hay.
Watermelons: Apply 2-2 1/2 lb. per acre when silks first appear. Two or three repeat applications at 2-3 day intervals will usually be necessary. Watermelon may be picked immediately following last application. Watermelon forage may be fed to dairy and beef cattle and sheep. Wait 4 days following treatment if forage is to be cut for hay.

Wireworms
Broccoli: Apply 4-8 lb. per acre just prior to planting. Work into soil 4-8 inches immediately. 2-3 inches for surface cut worms, 3-6 inches for subterranean cutworms.
Cucumbers (Surface and Subterranean): Broadcast 4-8 lb. per acre just prior to planting. Work into soil 4-8 inches immediately. 2-3 inches for surface cut worms, 3-6 inches for subterranean cutworms.
The Sweet Corn Earworm: Apply 2-2 1/2 lb. per acre when silks first appear. Two or three repeat applications at 2-3 day intervals will usually be necessary. Corn may be picked immediately following last application. Corn forage may be fed to dairy and beef cattle and sheep. Wait 4 days following treatment if forage is to be cut for hay.
Watermelons: Apply 2-2 1/2 lb. per acre when silks first appear. Two or three repeat applications at 2-3 day intervals will usually be necessary. Watermelon may be picked immediately following last application. Watermelon forage may be fed to dairy and beef cattle and sheep. Wait 4 days following treatment if forage is to be cut for hay.

Southern Potato Wireworms
Potatoes (Southern): Apply 4-8 lb. per acre just prior to planting. Work into soil 4-8 inches immediately. 2-3 inches for surface cut worms, 3-6 inches for subterranean cutworms.

Field and Forage Insects

Daznol W 50 should always be applied as a spray in sufficient water to assure thorough coverage of the foliage or soil, dependent upon the type of application.

Alice and Clover
Spotted Alfalfa Aphids, Green Clover Aphids, Flea Aphids, Leafhoppers, Plant Bug, Hairy Mites: Apply 1/4 lb. per acre.
Grasshoppers: Apply 1 lb. per acre.

Cucumbers (Surface and Subterranean): Broadcast 4-8 lb. per acre just prior to planting. Work into soil 4-8 inches immediately. 2-3 inches for surface cut worms, 3-6 inches for subterranean cutworms.
Daily and bestical care of crops and forage is necessary. If crops are heavily infested, they may be fed green alfalfa or corn immediately after harvest. If crops are heavily infested, they may be fed green alfalfa or corn immediately after harvest. If crops are heavily infested, they may be fed green alfalfa or corn immediately after harvest.

Alfalfa: Apply 1 lb. per acre just prior to planting. Work into soil 1-2 inches immediately. 1-2 inches for surface cut worms, 1-2 inches for subterranean cutworms.
Alfalfa: Apply 1 lb. per acre just prior to planting. Work into soil 1-2 inches immediately. 1-2 inches for surface cut worms, 1-2 inches for subterranean cutworms.

Do not spray livestock directly with the ether and ground equipment except as directed below under Livestock Insects.
Do not treat during bloom to avoid injury to pollinating insects.

Corn
Corn Rootworm Larvae

Postemergence Basal Treatment: Apply 1-2 lb. per acre just prior to planting. Work into soil 1-2 inches immediately. 1-2 inches for surface cut worms, 1-2 inches for subterranean cutworms.
Preemergence Basal Treatment: Apply 1-2 lb. per acre just prior to planting. Work into soil 1-2 inches immediately. 1-2 inches for surface cut worms, 1-2 inches for subterranean cutworms.

Suggested Rates According to Row Spacing For Corn Rootworm Larvae Control

Row Spacing	Linear feet of corn row per acre	Amount Daznol W 50 per acre	
		Linear feet of corn row per acre	Amount Daznol W 50 per acre
46	10,646	14,261	14,261
42	12,455	16,607	16,607
40	13,608	18,160	18,160
38	14,939	20,000	20,000
36	16,250	22,143	22,143
30	17,424	23,896	23,896
24	21,762	29,681	29,681
20	26,138	34,574	34,574
15	34,849	46,465	46,465

Note: For row spacing in feet, calculate number of rows per acre and multiply by the amount of Daznol W 50 per acre.

Seed Corn Maggots: Broadcast 4-8 lb. per acre just prior to planting. Work into soil 4-8 inches immediately.
Cucumbers (Surface and Subterranean): Broadcast 4-8 lb. per acre just prior to planting. Work into soil 4-8 inches immediately. 2-3 inches for surface cut worms, 3-6 inches for subterranean cutworms.
The Sweet Corn Earworm: Apply 2-2 1/2 lb. per acre when silks first appear. Two or three repeat applications at 2-3 day intervals will usually be necessary. Corn may be picked immediately following last application. Corn forage may be fed to dairy and beef cattle and sheep. Wait 4 days following treatment if forage is to be cut for hay.

Corn Rootworm Adults: Apply 1 lb. per acre.

Control Alfalfa Aphids, Green Clover Aphids, Flea Aphids, Leafhoppers, Plant Bug, Hairy Mites: Apply 1/4 lb. per acre.
Grasshoppers: Apply 1 lb. per acre.

Snap Beans: Apply 2-2 1/2 lb. per acre. Apply 2 lbs. per acre just prior to planting. Work into soil 2-3 inches immediately. 2-3 inches for surface cut worms, 3-6 inches for subterranean cutworms.

Corn may be picked immediately following last application. Corn forage may be fed to dairy and beef cattle and sheep.

Cotton
Spider Mites: Apply 1 lb. per acre just prior to planting. Work into soil 1-2 inches immediately. 1-2 inches for surface cut worms, 1-2 inches for subterranean cutworms.

5-25 gals. of water per acre. Repeat as necessary. Do not apply within 10 days of harvest. Do not graze treated cotton in pastures or feed to livestock.

Cowpeas
Cucumbers (Surface and Subterranean): Broadcast 4-8 lb. per acre just prior to planting. Work into soil 4-8 inches immediately. 2-3 inches for surface cut worms, 3-6 inches for subterranean cutworms.

40/3

1. The first step is to identify the problem or question that needs to be answered. This involves understanding the context and the specific information required.

[illegible]