



fuldicide-14

CLEANER • DISINFECTANT
DEODORIZER • FUNGICIDE

This product is a heavy duty cleaner and disinfectant designed for general purpose use where heavy duty cleaning is of prime importance. When used as directed, it will remove most dirt and grime in a single simple application. When used for industrial cleaning, follow the use directions shown below for best results. Not recommended for use on marble.

DANGER: Keep Out of Reach of Children. Can cause severe eye and skin damage. Do not get in eyes, on skin or on clothing. Wear goggles or face shield and protect skin when handling. Harmful or fatal if swallowed. Avoid contamination of food.

FIRST AID: In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. Immediately call a physician. Remove and wash all contaminated clothing before reuse. If swallowed, drink milk, egg whites, gelatin solution, or if these are not available, drink large quantities of water. Call physician.

USE DIRECTIONS

For light duty cleaning, and deodorizing, use two ounces per gallon.

For heavy duty cleaning and disinfecting and fungicidal activity against pathogens fungi, use 4.5 ounces per gallon.

ACTIVE INGREDIENTS:

Allyl (80% C14), 30% C18, 9% C12, 9% C18)	
dimethyl benzyl ammonium chloride	0.8%
Allyl (80% C12, 20% C14) dimethyl	
ethylbenzyl ammonium chloride	0.8%

ACCEPTED

INERT INGREDIENTS:

88.4%

EPA Reg. No 257-202

EPA Est 257-MD-1

MAY 04 1988

Keep Out Of Reach Of Children

DANGER

SEE FIRST AID STATEMENT AND OTHER PRECAUTIONARY STATEMENTS ON SIDE PANEL

NET CONTENTS: GALLONS

CELLUCOR CORP

For best results, use with a cloth or mop and apply to walls, floors and other hard surfaces.

For extra heavy soil or industrial cleaning use second application. Do not mix with soap or other solutions. Is non-corrosive in use dilutions on inanimate surfaces. Rinse empty container thoroughly with water and discard it.

STORAGE AND DISPOSAL

Store only in tightly closed, original container in a secure area inaccessible to children. Do not contaminate water, food, or feed by storage or disposal. Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or residue is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the hazardous waste representative of the nearest EPA Regional Office for guidance.

Do not reuse empty container. Triple rinse empty container with water. Return metal drums to reconditioner or purchaser and dispose of in a sanitary landfill or by other procedures approved by state and local authorities. Plastic containers may be disposed of in a sanitary landfill, incinerated or, if allowed by local authorities, by burning. If burned, stay out of smoke.

Contains, Contains & Packing Product of USA

Made for Industrial & Institutional Use Only

L
J

NOTE: Changes in labeling formula differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above U.S. EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby Registered/Re-registered under the Federal Insecticide, Fungicide, and Rodenticide Act.

A copy of the labeling accepted in connection with this Registration/Re-registration is returned herewith.

Registration in no way to be construed as an endorsement or approval of that product by this Agency. In order to protect health and the environment, the Administrator, in his discretion, may at any time suspend or withdraw the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

BEST AVAILABLE COPY

ATTACHMENT IS APPLICABLE

SIGNATURE OF APPROVING OFFICIAL

DATE 9/2/77

EPA Form 8570-6 (Rev. 5/76)

PREVIOUS COPY ON MAY BE USED UNTIL SUPPLY IS EXHAUSTED

5002; I; Richards; LR-15; RMDCO; 9/01/88; 10/13/88; CB; PBR; VO; CI.

10/3

- e. You may delete the crop restriction statement per PR Notice 87-3).
- f. Update the Environmental Hazards statement by changing
". . . cleaning of equipment and disposing of wastes" to
". . . disposing of equipment washwaters."
- g. Change the Note to Physician to the following:

Note to Physician: Solvent may present aspiration hazard. Gastric lavage may be indicated if product was taken internally. Disulfoton is an organophosphate insecticide. If symptoms of cholinesterase inhibition are present, atropine sulfate by injection is antidotal. 2-PAM is also antidotal and may be administered, but only in conjunction with atropine.

3. Submit a 1-year storage stability study for the formulated product. It must be analyzed for its active ingredient at time zero and during a year of storage. This study should be submitted within 15 months from the date of this Registration Notice.

4. Submit five (5) copies of your final printed labeling before you release the product for shipment. Refer to the A-79 enclosure for a further description of final printed labeling.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

George T. LaRocca
Product Manager (15)
Insecticide-Rodenticide Branch
Registration Division (TS-767C)

Enclosures

COMMERCIAL FERTILIZER

with 1.00% Phostemic-D*

CONTAINS 1.00% DI-SYSTON†

STATEMENT OF INGREDIENTS FOR PEST CONTROL

ACTIVE INGREDIENT:	BY WT.
disulfoton O, O-Dimethyl S-[2-(ethylthio)ethyl] Phosphorodithioate	1.00%
INERT INGREDIENTS	99.00%
TOTAL	100.00%

U.S. Patent No. 2,759,310
DI System Reg. U.S. Pat. Off. by Farnbacher & Bayer, A. G. Chemagro Corporation, Licensee
Phostemic D (DI System) is a systemic pesticide which penetrates by absorption and is translocated in the plant. The active ingredient is released slowly, giving long protection during the plant's susceptible early life.

EPA Reg. No. 7001

EPA Est. No. 7001-CA-1

KEEP OUT OF REACH OF CHILDREN WARNING AVISO

Precaution Al Usuario: Si usted no lee ingles, no use este producto hasta que la etiqueta haya sido explicada en espanol.

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. Get medical attention.

IF INHALED remove victim to fresh air. Apply artificial respiration if indicated.

IF ON SKIN remove contaminated clothing. Wash affected areas with soap and water. If irritation persists call a physician.

IF IN EYES flush with plenty of water. Call a physician.

NOTE TO PHYSICIAN: Contains organic phosphate insecticide, inhibits cholinesterase. If symptoms of cholinesterase inhibition are present atropine sulfate is antidotal and may be administered in conjunction with 2-PAM.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING

Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by other persons. Wear protective clothing and cotton or natural rubber gloves. If cotton gloves are used, they must be laundered or discarded after each day's use. Do not wear the same gloves for other work. Wash rubber gloves with soap and water after each use.

Rapidly absorbed through skin. May be fatal if swallowed. Do not get on skin, in eyes, or on clothing. Wash hands, arms and face thoroughly with soap and water before eating or smoking. Wash all contaminated clothing before re-use.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and wildlife. Do not apply directly to water or wetlands. Drift and runoff from treated areas may be hazardous to fish in neighboring areas. Do not contaminate water by cleaning of equipment or disposal of wastes. Cover or incorporate spills.

This pesticide is toxic to bees exposed to direct application. Applications should be timed to coincide with periods of minimum bee activity, usually late evening or early morning.

CROP RESTRICTION

Do not plant any food or feed crop in rotation after a field treatment with Disulfoton unless it is a registered use for Disulfoton.

RE-ENTRY STATEMENT

Do not apply this product in a manner as to directly or through drift expose workers or other persons. Do not enter treated areas without protective clothing until the dust has settled. Certain states require more restrictive re-entry intervals for various crops treated with this product. Consult your State Department of Agriculture for further information.

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. When oral warnings are given, written warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Written and oral warnings must include the following: WARNING: Area treated with Commercial Fertilizer with Phostemic-D on (date of application). Do not enter without appropriate protective clothing for 24 hours. Do not allow re-entry into treated fields within 24 hours of application unless appropriate protective clothing is worn. Protective clothing: long sleeved shirt, long-legged trousers, impermeable apron, gloves and shoes.

BRUSSELS SPROUTS: For control of Aphids, Mites, Thrips, Leafhoppers and Flea Beetles apply 100 lbs. Acre by sidedressing at planting time or after emergence. Do not apply more than twice per season or within 30 days of harvest.

CABBAGE: For control of Aphids and Flea Beetles apply 100 lbs. Acre by sidedressing at planting time or after emergence. Do not apply more than once per season or within 42 days of harvest.

CAULIFLOWER: For control of Aphids, Leafhoppers and Flea Beetles apply 100 lbs. Acre by sidedressing at planting time or after emergence. Do not apply more than twice per season or within 40 days of harvest.

COTTON: For control of Aphids, Thrips and Mites apply 100 lbs. Acre by sidedressing at planting time or after emergence. Do not apply more than twice per season or within 28 days of harvest. Allow a minimum of 21 days between applications. Read Important statement.

LETTUCE: For control of Aphids, Thrips, Mites and Leafhoppers on spring or summer seeded lettuce apply 100 to 200 lbs. Acre by sidedressing at planting time only. On fall or winter seeded lettuce apply at planting time or thinning time. Do not apply within 60 days of harvest. Read Important statement.

POTATOES: For control of Aphids, Flea Beetles (Except Tuber Flea Beetles), Leafhoppers, Potato Psyllid and Southern Potato Wireworm apply 200 to 400 lbs. Acre in the furrow at planting, or by sidedressing after plants become established. Higher rate to be used in organic soils. Do not apply within 15 days of harvest. Do not make more than 2 applications per season. Do not tank mix this product with Phosalone.

SNAP OR GREEN LIMA BEANS: For control of Aphids, Thrips, Mites and Leafhoppers apply 100 to 200 lbs. Acre by sidedressing at planting time only. Read Important statement. Do not apply more than once per season.

SORGHUM: For control of Aphids, Green Bug, apply 100 lbs. Acre at planting time in furrow or band application. Do not apply within 30 days of harvest for grain or 14 days of harvest for forage or fodder. Do not apply more than once per crop season, regardless of granular application.

SPINACH: For control of Aphids, Thrips, Mites and Leafhoppers apply 100 lbs. Acre by sidedressing at planting time only.

SUGAR BEETS: For control of Aphids, Thrips, Mites and Leafhoppers apply 100 lbs. Acre by sidedressing at planting time or after emergence. Do not apply more than 3 times per season or within 30 days of harvest. Read Important statement.

TOMATOES: For control of Aphids, Flea Beetles, Leaf Miners, Leafhoppers and Mites up to 300 lbs. Acre may be applied by sidedressing at planting time or with no other application is being made during the season. If 2 applications are required, apply 200 lbs. Acre each application. Apply at planting time or after plants become established. Do not apply within 30 days of harvest.

IMPORTANT: When used on BEANS, COTTON, LETTUCE or SUGAR BEETS at the maximum recommended rate under adverse conditions such as extremely cool and wet or extremely dry weather, Di-Syston may cause some delay in emergence, stunting of seedlings or reduction of stand. Such damage may be more pronounced in light, sandy soils. Plant injury may also occur or may be enhanced when certain pre-emergent herbicides are used in conjunction with Di-Syston.

AGC 597 4 87

ACCEPTED
WITH COMMENTS
in EPA Label Draft

05-0

Up to 300 lbs. Acre
Registered under EPA Reg. No.
7001-355

30/3

Net Weight