

400-312 1045

US ENVIRONMENTAL PROTECTION AGENCY OFFICE OF PESTICIDES PROGRAMS REGISTRATION DIVISION (75-767) WASHINGTON, DC 20460	EPA REGISTRATION NO.	DATE OF ISSUANCE
	7001-3773	6/17/94
	TERM OF ISSUANCE	
NOTICE OF PESTICIDE: <input type="checkbox"/> REGISTRATION <input type="checkbox"/> REREГИSTRATION (Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended)		NAME OF PESTICIDE PRODUCT ... 6

NAME AND ADDRESS OF REGISTRANT (Include ZIP code)

J.P. Sappelt Co.
 34777 Houlston Rd.
 P.O. Box 100
 Isthmian, CA 95320

NOTE: Changes in labeling formula differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above U.S. EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby Registered/Reregistered under the Federal Insecticide, Fungicide, and Rodenticide Act.

A copy of the labeling accepted in connection with this Registration/Reregistration is returned herewith.

Registration is in no way to be construed as an indorsement or approval of this product by this Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered under FIFRA section 3(c)(7)(A) provided that you:

1. Submit all data required for registration of your product under FIFRA section 3(c)(7)(A) and comply with all requirements of FIFRA section 3(c)(7)(A).

2. Make the labeling changes listed below before you release the product for shipment:

- Add the phrase "EPA Registration No. 7001-3773"
- Submit five (5) copies of your final printed labeling before you release the product for shipment.

If these conditions are not complied with, this registration will be subject to cancellation in accordance with FIFRA section 3(c)(7)(A). Your release for shipment of the product under this registration is a condition.

For more information, contact the Registration Division, Office of Pesticides Programs, U.S. Environmental Protection Agency, Washington, DC 20460.

BEST AVAILABLE COPY

ATTACHMENT IS APPLICABLE

SIGNATURE OF APPROVING OFFICIAL	DATE
<i>Robert L. Chandler</i>	6/17/94

SOCHLOR 6

ACTIVE INGREDIENT:

Sodium Chlorate 47.2%


INERT INGREDIENTS: 52.8%

TOTAL 100.0%

Each gallon contains 6.0 pounds of Sodium Chlorate per gallon

EPA Reg. No. 7001-

EPA Est. No.



WARNING

AVISO

Si usted no entiende la etiqueta, busque a alguien para se la explique a usted en detalle. (If you do not understand the labels, find someone to explain it to you in detail.)

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Drink promptly a large quantity of milk, egg whites, gelatin solution, or if these are not available, drink large quantities of water. Avoid alcohol. Get medical attention.

IF IN EYES: Flush with water for 15 minutes. Get medical attention.

IF ON SKIN: Wash with plenty of soap and water. Get medical attention if irritation persists.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage.

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

Harmful if swallowed. May irritate eyes, nose, throat and skin. Avoid contact with skin and eyes. Avoid breathing spray mist.

PERSONAL PROTECTIVE EQUIPMENT

APPLICATORS AND OTHER HANDLERS MUST WEAR:



- . Coveralls over short-sleeved shirt and short pants
- Waterproof gloves
- . Chemical-resistant footwear plus socks
- . Protective eyewear
- . Chemical-resistant headgear for overhead exposure
- . Chemical-resistant apron when cleaning equipment, mixing, or loading
- . Dust/mist filtering respirator (MSHA/NIOSH approval number prefix TC-21C)

USER SAFETY RECOMMENDATIONS

Users should: (1) Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet; (2) Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing; (3) Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into cleaning clothing.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, to areas where surface water is present or to intertidal area below the mean high water mark. Do not contaminate water when disposing of equipment washwaters. Do not apply when weather conditions favor drift away from target area.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

STORAGE: Store in a cool, dry, fire-resistant area; separate from acids, solvents, oils, organic substances, sulfur, powdered metals and away from fire hazards.

DISPOSAL: Triple rinse or equivalent. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

IN CASE OF ACCIDENTAL SPILL: Use absorbent material, sweep up or vacuum. If spilled on concrete or metal floors with water after sweeping or vacuuming.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons either directly or through drift. Only protected handlers may be in the area during application. For any requirement specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

BEST AVAILABLE COPY

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), notification to workers and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Workers Protection Standard.

Do not enter or allow entry into treated areas during the restricted entry interval (REI) of 12 hours. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water, is: Coveralls over short-sleeved shirt and short pants; waterproof gloves; chemical-resistant footwear plus socks; protective eyewear; chemical-resistant headgear for overhead exposure.

GENERAL INFORMATION

DEFOLIATION: Weather conditions and plant conditions affect defoliation. Weather conditions that favor good defoliation are high temperatures, high humidity, low wind velocity, high to adequate soil moisture. Plant conditions which favor good defoliation are ample fertility and moisture, complete insect control, heavy boll load, all bolls mature, vegetative growth ceased, little or no second growth. Conditions that contribute to poor defoliation are those adverse to the above. Nutrients should be ample but not too much nitrogen. Leaves should be green and turgid. Not wilted, toughened or stunted by lack of fertilizers.

DESICCATION: To reduce moisture of the crop and desiccate weeds prior to harvest apply during the conditions listed as favorable under the defoliation section of the label. Incomplete coverage will give erratic results. Especially with vines such as morning glory where the tender stems may be sheltered from the spray by the crop canopy. Complete coverage is essential to good desiccation.

DILUTION RATES:

AIR: 4-10 gallons of water.

GROUND: 20-30 gallons of water

BEST AVAILABLE COPY

COTTON: Apply 1/2 to 3/4 gal. per acre 2-3 weeks prior to picking date. Do not apply within 7 days of harvest. Bolls less than 35 days old may not mature after application of this product. Do not apply during conditions of extreme heat during the middle of the day. Do not graze treated areas or feed gin waste to livestock.

FLAX: As a harvest aid to desiccate weeds and mature the crop, apply 1 gal. per acre, 7-10 days before anticipated harvest date. Activity of the spray will be enhanced by the

addition of a good non-ionic surfactant. Refer to the surfactant label for specific rates. Do not graze treated fields or feed treated flax foliage and fodder for 14 days after application.

GRAIN SORGHUM: To reduce the moisture content while the crop is standing, use 3/4 to 1 gal. per acre, 7-10 days before anticipated harvest date. Use the lower rate when grain sorghum and weather conditions are right for rapid desiccation. Do not graze treated fields or feed treated fodder, forage or residual seeds within 14 days of application.

GUAR BEANS: To reduce bean seed moisture, apply 1 gal. per acre, 7-10 days before anticipated harvest date. Do not apply under conditions of extreme heat during the middle of the day. Do not graze treated fields or feed treated fodder or forage to livestock.

SAFFLOWER: To reduce moisture in the fully mature seed heads, apply 1 gal. per acre, 7 or more days before normal harvest and when safflower heads are fully mature. Do not apply later than 7 days before harvest. Do not graze treated fields or feed treated forage.

SOYBEANS: To desiccate weeds in early maturing soybeans and facilitate the harvest, apply 1 gal. per acre, 7-10 days before anticipated harvest date when beans are mature and ready for harvest. Do not graze treated fields or feed treated bean foliage and fodder.

SUNFLOWERS (Confectionery and Oil Seed varieties): To facilitate harvest and reduce moisture in the seedhead, apply 3/4 to 1 gallon, 7-10 days before anticipated harvest date. Use the lower rate when conditions favor rapid desiccation. Do not graze treated fields or use residual seeds for food or feed within 14 days of harvest.

RICE: To facilitate harvest and reduce moisture in the full mature seed heads and to desiccate immature red rice in mature white rice, apply 3/4 to 1 gal. per acre. Apply 7-10 before anticipated harvest. Use the lower rates when crop conditions and weather are right to enhance rapid drying. To avoid drift, do not apply during windy conditions. Do not graze treated fields or feed fodder, forage or residual seeds within 14 days of harvest.

NET CONTENTS:

DISCLAIMER OF WARRANTIES. Seller warrants that the chemical composition of this product conforms to the chemical description given on this label. THIS WARRANTY IS EXPRESSLY IN LIEU OF ALL OTHER WARRANTIES AND REPRESENTATIONS EXPRESSED, IMPLIED OR STATUTORY, INCLUDING WARRANTIES OF MERCHANTABILITY AND FITNESS FOR A PARTICULAR USE. Timing, rate and method of application, weather and crop conditions, mixtures with chemicals not specifically recommended on this label or an accompanying written recommendation, are beyond the control of seller. Buyer assumes all risks of use, storage and handling of this material not in strict accordance with directions given herewith. Buyer further agrees in the event of damages arising from the use of this product to accept a replacement, one is authorized to make any other warranty, guarantee or directions concerning this product, and no such warranties, guarantees, or directions shall be valid or binding upon seller.

Simplot

J.R. SIMPLOT COMPANY
P.O. Box 198
Lathrop, CA 95330

P.O. Box 15057
Boise, ID 83715

BEST AVAILABLE COPY