



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

DEC 18 2009

Mrs. Lisa J. Strong
Regulatory
J.R. Simplot Company
P.O. Box 198
Lathrop, CA 95330

RE: Notification to add Additional Brand Name: **"EverGreen 18-5-0 Minis + 5% Iron with 0.20% Dimension"**

EPA Registration Number: 7001-374
Date of Submission: November 10, 2009

Dear Mrs. Strong:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 dated, November 10, 2009, for the above mentioned product. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the actions requested fall within the scope of PRN 98-10. The label submitted with the application has been stamped "Notification" and will be placed in our records.

If you have any questions, please me directly at 703-305-6249 or Joyce Edwards of my staff at 703-308-5479.

Sincerely,



Linda Arrington
Notifications & Minor Formulations Team Leader
Registration Division (7505P)
Office of Pesticide Programs

2098

Please read instructions on reverse before completing form.

Form Approved, OMB No. 2070-0060

Print Form



United States
Environmental Protection Agency
Washington, DC 20460

☐ Registration
☐ Amendment
☒ Other

OPP Identifier Number

Application for Pesticide - Section I

1. Company/Product Number 7001-374	2. EPA Product Manager Joanne Miller	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) 0.20% Dimension	PM# 23	
5. Name and Address of Applicant (Include ZIP Code) J.R. Simplot Company P.O. Box 198 Lathrop, CA 95330 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	

Section - II

<input type="checkbox"/> Amendment - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____	NOTIFICATION DEC 18 2009
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.	
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below.	

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

NOTIFICATION OF ADDITIONAL BRAND NAMES PER PR NOTICE 98-10 This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.

Section - III

1. Material This Product Will Be Packaged In:				2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes* <input checked="" type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No		<input checked="" type="checkbox"/> Metal	
* Certification must be submitted				<input type="checkbox"/> Plastic	
	If "Yes" Unit Packaging wgt.	No. per container	If "Yes" Package wgt	<input type="checkbox"/> Glass	
			No. per container	<input type="checkbox"/> Paper	
				<input type="checkbox"/> Other (Specify) _____	
3. Location of Net Contents Information <input checked="" type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container 50 lb. bag		5. Location of Label Directions <input checked="" type="checkbox"/> On Label <input type="checkbox"/> On Labeling accompanying product	
6. Manner in Which Label is Affixed to Product plastic sleeve attached to bag		<input type="checkbox"/> Lithograph <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled		<input type="checkbox"/> Other _____	

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)			
Name Lisa J. Strong		Title Regulatory	
		Telephone No. (Include Area Code) 209-858-2511	
Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment both under applicable law.			6. Date Application Received (Stamped)
2. Signature 		3. Title Regulatory	
4. Typed Name Lisa J. Strong		5. Date November 10, 2009	

3068

J.R. SIMPLOT COMPANY 16777 HOWLAND ROAD P.O. BOX 198
LATHROP, CALIFORNIA 95330 (209) 858-2511 FAX (209) 858-2519

AGRIBUSINESS

November 10, 2009

Ms. Joanne Miller
Product Manager (23)
U.S. Environmental Protection Agency
Document Processing Desk (NOTIF)
Registration Division (7504P)
Crystal Mall 2, 1801 South Bell Street
Arlington, VA 22202

**RE: NOTIFICATION – ADDITIONAL BRAND NAME
0.20% DIMENSION
EPA REG. NO. 7001-374**


Dear Ms. Miller:

J.R. Simplot Company is submitting Notification that the subject product will be marketed under the additional brand name:

EverGreen 18-5-0 Minis + 5% Iron with 0.20% Dimension

This Notification is in compliance with PR-Notice 98-10. An Application form 8570-1 is enclosed with the compliance statement.

Sincerely,
J.R. SIMPLOT COMPANY


Lisa J. Strong
Regulatory

3068

NOTIFICATION

DEC 18 2009

EverGreen 18-5-0 *Minis* + 5% Iron with 0.20% DIMENSION®

EverGreen 18-5-0 Minis + 5% Iron with 0.20% Dimension® Is A Herbicide That Provides Control Of Listed Annual Grasses and Broadleaf Weeds In Ornamental Landscape Areas, Established Lawns And Ornamental Turf, Including Golf Course Fairways, Roughs And Tee Boxes. This product also provides postemergence crabgrass control up to the 4-leaf stage

Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in interior plantscapes, ornamental gardens or parks, or on golf courses or lawns and grounds. In New York State, this product may only be used by Professional Applicators at no more than 250 lbs. per acre (5.74 lbs./1,000 sq.ft.), (0.5 lb. active ingredient per acre) per year and is prohibited from use in Nassau and Suffolk Counties.

ACTIVE INGREDIENT:

Dithiopyr, 3,5-pyridinedicarbothioic acid, 2-(difluoromethyl)-4-(2-methylpropyl)-6-(trifluoromethyl)-S,S-dimethyl ester 0.20%

OTHER INGREDIENTS:

TOTAL 99.80%

Read The Entire Label Before Using This Product. Use Only According To Label Instructions.

Product protected by U.S. Patent No. 4,692,184. Other patents pending.

EPA REG. No. 7001-374

EPA EST. No. 61842-CA-1

NOTICE: Before using this product, read the entire Precautionary Statements, Conditions of Sale and Warranty, Directions for Use, Use Restrictions and Storage and Disposal Instructions. If the Conditions of Sale and Warranty are not acceptable, return the product unopened within thirty days of purchase to the place of purchase.

448

KEEP OUT OF REACH OF CHILDREN CAUTION

FIRST AID: IF IN EYES—Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice. IF INHALED—Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice. **HOT LINE NUMBER**—Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact the National Pesticide Information Center at 1-800-858-7378 for emergency medical treatment information.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION! CAUSES EYE IRRITATION. HARMFUL IF INHALED. Avoid contact with eyes or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Prolonged or frequently repeated skin contact while handling this material may cause allergic reaction in some individuals.

ENVIRONMENTAL HAZARDS

This product is toxic to fish and highly toxic to other aquatic organisms including oysters and shrimp. Use with care when applying to turf areas adjacent to any body of water. Drift and runoff from treated turf may adversely affect aquatic organisms in adjacent aquatic sites. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

GUARANTEED ANALYSIS:

SPREADER SETTINGS (Based on the South REGION rate recommendations)

	PROGRAM 1	PROGRAM 2	PROGRAM 3
Rates per 1,000 sq.ft.			
SPREADERS			

GENERAL INFORMATION FOR TURF USE

This product is a herbicide that provides control of crabgrass and other annual grasses and broadleaf weeds in established lawns and ornamental turfs including golf course fairways, roughs and tee boxes. This product also provides post-emergent crabgrass control up to the 4-leaf stage. This product will not harm nearby established ornamentals when used according to label directions.

USE PRECAUTIONS

Turfgrass Safety: This product may be used on seeded, sodded or sprigged lawns and ornamental turfs that are well established. The grass must have developed a good root system and a uniform stand, and have received at least two mowings following its seeding, sodding or sprigging before it can receive its first application of this product. Use of this product on lawns and ornamental turfs that are not well established, or on those that have been weakened by weather-, pest-, disease-, chemical-, or mechanical-related stress, may increase the chances for turf injury.

This product should only be applied to lawns and ornamental turfs that are composed of the following turfgrass species that have been determined to be tolerant to applications of this product. When applied as directed under the conditions described, the following established turfgrasses are tolerant to this product:

COOL-SEASON GRASSES: Bentgrass, creeping* (*Agrostis palustris*), Bluegrass, Kentucky (*Poa pratensis*), Fescue, fine** (*Festuca rubra*), Fescue, tall (*Festuca arundinacea*), Ryegrass, perennial (*Lolium perenne*).

WARM-SEASON GRASSES: Bahiagrass (*Paspalum notatum*), Bermudagrass (*Cynodon dactylon*), Buffalograss*** (*Buchloe dactyloides*), Carpetgrass (*Axonopus affinis*), Centipedegrass (*Eremochloa ophiuroides*), Kikuyugrass (*Pennisetum clandestinum*), St. Augustinegrass (*Stenotaphrum secundatum*), Zoysiagrass (*Zoysia japonica*). DO NOT apply this product to Colonial bentgrass (*Agrostis tenuis*) varieties.

*Use of this product on certain varieties of creeping bentgrass, such as "Carmen", "Cohansey", or "Seaside" and "Washington" may result in undesirable turf injury. Not all varieties of creeping bentgrass have been tested. **Use of this product on certain varieties of fine fescue may result in undesirable turf injury. The following fine fescue varieties have been found to be sensitive to this product: "Atlanta", "Banner", "Beauty", "Bilgort", "CF-2", "Enjoy", "HF-93", "Highlight", "Ivalo", "Jamestown", "Koket", "Majenta", "Mary", "Penlawn", "Tamara", "Tajana", "Waldorf", and "Waldina". Not all varieties of fine fescues have been tested. ***DO NOT use this product on seedling buffalograss in the spring or the first year of establishment until the turfgrass is fully green and has established new roots.

NOTE: It is not possible to evaluate all turfgrass varieties for tolerance to BEST EverGreen 18-5-0 with 0.20% Dimension. It is suggested that the user evaluate the responses of turfgrass varieties to this product on a limited area, at recommended use rates, prior to initiating large-scale use.

Reseeding, Overseeding, or Sprigging: Reseeding, overseeding, or sprigging of treated areas with this product should be delayed until 12 weeks from the time of application. Reseeding, overseeding or sprigging before 12 weeks after application may prevent establishment of desirable turfgrasses. When reseeding or overseeding, proper cultural practices such as soil cultivation, irrigation and fertilization should be followed. If overseeding with perennial ryegrass in the fall, overseed 8 weeks or later after a single application. For best results, use mechanical or power seeding equipment (slit seeders) designed to give good seed-to-soil contact.

Other Use Precautions: The early postemergence crabgrass control described below is limited. Postemergence applications are only effective on crabgrass if applied before crabgrass has displayed its fifth leaf or first tiller. For best results, cultural practices that disturb the soil such as core-, spike-, or hydro-aerification and verticutting, should be done before application of this product. DO NOT apply this product until grass has recovered from these cultural practices. DO NOT use clippings from treated turf for mulching around vegetables or fruit trees. Keep children and pets out of treated areas until dust has settled.

6008

APPLICATION DIRECTIONS – Control of Crabgrass

Preemergence and Early Postemergence Control

- This product provides "preemergence" control of crabgrass (including the large, smooth and southern species) when applied prior to the emergence of crabgrass from the ground in established lawns and ornamental turfs.
- It can also provide "early postemergence" control of crabgrass. Postemergence crabgrass control will be obtained only when this product is applied prior to the tillering of crabgrass, which generally corresponds to the time when you can first see the crabgrass plants in the lawn or turf. The practical benefit of this product's early postemergence activity is that it can give the user a 2-4 week longer period of time to make applications to control crabgrass.

Application Frequency and Timing:

- This product may be applied as a single application or as a split application for crabgrass control in the spring, summer or fall.
- DO NOT apply more than 5.74 lb. of this product/1,000 sq. ft. per application, and no more than 13.7 lbs. of this product per 1,000 sq. ft. per year. In New York State, this product may only be used by Professional Applicators at no more than 250 lbs. per acre (5.74 lbs./1,000 sq. ft.) (0.5 lb. active ingredient per acre) per year; and is prohibited from use in Nassau and Suffolk Counties.

Spring Applications:

- For single applications made in the spring or early summer, this product should be applied at the appropriate rate corresponding to one of the control programs listed in Table 1 below, depending on the user's location, the turfgrass mowing height, and whether the use is preemergence or early postemergence at the time of application. The duration of residual weed control provided by this product is directly related to the total rate applied, but will vary somewhat depending on weather, weed pressure, turfgrass competitiveness, and the user's location within a region.
- Use Program 1 for preemergence control at sites where the turfgrass is cut high (e.g., homeowner lawns). This program provides 3-5 months of preemergence crabgrass control. This program should not be used for early postemergence crabgrass control.
- Use Program 2 for preemergence control at sites where (a) turfgrass is cut low (e.g., golf fairways), and (b) turfgrass maintenance or weed control has not been conducted during the previous year. This program provides 4-6 months of preemergence crabgrass control.
- Use Program 3 for preemergence control at sites where (a) turfgrass is cut low (e.g., golf fairways) and (b) turf maintenance or weed control has not been conducted during the previous year. This program provides 4-6 months of preemergence crabgrass control. This program may also be used for early postemergence control at sites where turfgrass is cut low (e.g., golf fairways).
- Subsequent, sequential pre- and/or postemergence applications should be made where longer periods of control are desired.
- Where split fertilizer applications are recommended, the rates in Table 1 may be applied in two applications made 6-10 weeks apart and prior to crabgrass emergence.

TABLE 1: APPLICATION USE RATES*
(Lbs. of This Product / 1,000 sq.ft.)

PROGRAM	1	2	3
Use	Preemergence and Postemergence on Crabgrass	Preemergence and Postemergence on Crabgrass	Preemergence and Postemergence on Crabgrass
REGION			
North	1.43 lbs./1,000 sq.ft.	2.07 lbs./1,000 sq.ft.	2.87 lbs./1,000 sq.ft.
Transition	2.07 lbs./1,000 sq.ft.	2.87 lbs./1,000 sq.ft.	4.26 lbs./1,000 sq.ft.
South	2.87 lbs./1,000 sq.ft.	4.26 lbs./1,000 sq.ft.	5.74 lbs./1,000 sq.ft.
Coastal South	4.26 lbs./1,000 sq.ft.	5.74 lbs./1,000 sq.ft.	5.74 lbs./1,000 sq.ft.

*DO NOT apply more the 5.74 lbs. of this product/1,000 sq. ft. per application, and no more than 17.22 lb. of this product/1,000 sq. ft., per year. In New York State, this product may only be used by Professional Applicators at no more than 250 lbs. per acre (5.74 lbs./1,000 sq. ft.) (0.5 lbs. active ingredient per acre) per year; and is prohibited from use in Nassau and Suffolk Counties. Postemergence control is limited; see "Crabgrass Control" and "Precautions" sections above. Regions include areas listed below.

North: All other areas.

Transition: DE, KS, KY, MD, MO, NJ, VA, southeastern PA, southern areas of IL, IN, OH & coastal areas of CT, NY, & RI.

South: AL, AR, AZ, CA, GA, LA, MS, NC, NM, NV, OK, SC, TN, & TX.

Coastal South: HI, FL, & southern coastal areas of AL, GA, LA, MS, NC, SC, & TX.

Fall Applications

- Use as directed in the fall at the Program 3 rates listed in Table 1 to control winter weed species and for crabgrass control through the early part of next spring.
- The fall application should be followed by an appropriate spring application to provide season-long control.

Tips for Improved Control

- For best results, apply this product within a few days after mowing and delay mowing again for a few days after the application.
- When treated lawn or ornamental turf areas are watered or receive significant rainfall within a few days after application of this product, improved weed control may result.
- Use of split (half-rate) applications spaced 6-8 weeks apart may provide improved weed control.

Control of other Grass and Broadleaf Weeds

Spring Applications

Used as directed for crabgrass control in the spring, this product will also control (at the Program 3 rates) or suppress (at the Program 2 rates) the following weeds when applied prior to their emergence: barnyardgrass, bluegrass (annual, roughstalk), crowfootgrass, foxtail (green & yellow)*, goosegrass, kikuyugrass, lespedeza (common), oxalis (buttercup, creeping & yellow woodsorrel), purslane (common), ryegrass (annual & perennial), smutgrass, speedwell (corn), spurge (prostrate & spotted).

*Also controlled at the Program 2 rates.

Fall Applications

Used as directed, this product will also control (at the Program 3 rates) or suppress (at the Program 2 rates) the following weeds when applied prior to their emergence:

bluegrass (annual, roughstalk), bittercress, chickweed, henbit, geranium (Carolina), parsley-piert, pineappleweed, shepherd's purse.

APPLICATION EQUIPMENT AND INSTRUCTIONS

Apply this product with drop or rotary-type spreaders designed to apply granular herbicides. Avoid the use of spreaders that would apply this product in narrow rows or concentrated bands. Before each application, calibrate the spreader according to the equipment manufacturer's directions for adjusting the spreader settings such that the spreader delivers the appropriate application rate recommended above. Apply this product uniformly over the lawn or ornamental turf area. A more uniform application can be made by spreading half of the required amount over the area and then applying the remaining half at a right angle to the previous direction. Avoid streaking, skips, or overlaps during application. Check equipment frequently to ensure equipment is functioning properly and applying uniform distribution of granules.

7048

GENERAL INFORMATION – LANDSCAPE ORNAMENTALS

EverGreen 18-5-0 Minis + 5% Iron with 0.20% Dimension provides preemergence control of listed annual grass and broadleaf weeds in landscape ornamentals. It may be applied to soil surfaces or mulch where ornamental plants are grown for aesthetic purposes in ornamental gardens, parks, golf courses and residential properties. Ornamental plants on this label have shown tolerance to EverGreen 18-5-0 Minis + 5% Iron with 0.20% Dimension when applied as directed. It is impossible, however, to evaluate tolerance for all ornamental plant species or varieties or under all possible growing conditions. Under your growing conditions, a limited area involving only a few plants should be treated for tolerance evaluation prior to large-scale applications.

USE RATES

Apply EverGreen 18-5-0 Minis + 5% Iron with 0.20% Dimension with a properly calibrated spreader that will assure accurate, even particle distribution. This product may be applied as a single application or as split application. Refer to the following table for product rates when treating a small area.

DO NOT apply more than 5.74 lbs. of this product/1,000 sq. ft. per application, and no more than 13.7 lbs. of this product per 1,000 sq. ft. per year. In New York State, this product may only be used by Professional Applicators at no more than 250 lbs. per acre (5.74 lbs./1,000 sq. ft.), (0.5 lbs. active ingredient per acre) per year, and is prohibited from use in Nassau and Suffolk Counties.

Pounds Product/Acre	Pounds Product/1,000 Square Feet	Ounces Product/100 Square Feet
250	5.74	9.2

APPLICATION TIMING

EverGreen 18-5-0 Minis + 5% Iron with 0.20% Dimension is a preemergence herbicide that will not control established weeds. Make applications prior to weed seed germination or mulch or to bare ground. The best weed control is obtained when applied to soil that is free of clods, weeds and debris such as leaves. Prior to making an application, control existing vegetation by hand weeding, cultivation or with a postemergence herbicide. Once an application of this product has been made, do not disturb the soil surface as the herbicide barrier will be broken.

PRECAUTIONS

- Only apply this product to established ornamentals.
- DO NOT apply this product directly to bare root of ornamental plants as injury may result.
- DO NOT incorporate this product into soil. Dilution of active ingredient and possible injury to plant roots may occur.
- DO NOT apply to soil around ornamental plants that are under stress due to drought, flooding, excessive fertilizer or soil salts, wind injury, hail, frost damage, winter injury, injury from previously applied pesticides or injury due to insects, nematodes or disease.
- To avoid foliar damage to ornamentals, irrigate immediately after application to remove herbicide particles from the foliage.
- DO NOT apply under conditions that would affect product distribution. Wind could affect distribution and move product onto sensitive species. Uneven product distribution will result in uneven weed control.
- DO NOT apply this product on grasses grown for seed.
- DO NOT graze livestock or feed forage cut from areas treated with this product.

TOLERANT ORNAMENTALS

When applied as directed under the conditions described on this label, the ornamentals listed below have shown tolerance in field trials. However, this product has not been tested on all cultivars of each species or under all possible growing conditions. Follow directions given above to determine plant tolerance under your growing conditions prior to large-scale use.

Abelia, Dwarf (*Abelia X grandiflora*) Nana; **Ajuga** (*Ajuga reptans*) Bronze, (*Ajuga genevensis*) Bronze Beauty; **Almond, Flowering** (*Prunus gladiolosa*); **Apple*** (*Malus pumila*); **Arbovitae** (*Thuja occidentalis*) Nigra, Pyramidalis, Smarag, Techry, Woodwardii; **Arbovitae, Golden** (*Thuja orientalis*); **Aster, Chinese** (*Callistephus chinensis*) Dwarf Queen; **Ash, Green** (*Fraxinus pennsylvanica*); **Ash, Mountain** (*Sorbus aucuparia*); **Ash, Purple** (*Fraxinus Americana*); **Azalea** (*Rhododendron* spp.) Brilliant, Buccaneer, Carror, Chimes (Belgian), Elsie Lee, Exbury, Fashion, Hardizer Beauty, Hershey Red, Higasa, Hincorimson, Holland (Hybrid), Marion Lee, Northern Lights, Orange Cup, Orchid Lights, Snow, Southern Charm; **Azalea, Flame** (*Rhododendron calendulaceum*); **Azalea, Kirishima**; **Bamboo, Heavenly** (*Nandina domestica*); **Barberry** (*Berberis thunbergii*) Aurea, Dwarf Pygmy, Green, Kobold, Pygmy Red, Rose Glow; **Barberry, Purple** (*Berberis vulgaris*) Atropurpurea; **Basket Flower** (*Gaillardia grandiflora*); **Bearberry** (common) (*Arctostaphylos uva-ursi*) Massachusetts; **Bee Balm** (*Monarda didyma*); **Begonia** (*Begonia* spp.); **Birch, River** (*Betula nigra*); **Blackeyed Susan** (*Rudbeckia hirta*) Goldstrum; **Blanket Flower** (*Gaillardia* spp.) Bluecrop; **Blueberry*** (*Vaccinium* spp.) Blue Jay, Jersey, North Blue, Northland; **Bottlebrush** (*Callistemon citrinus*); **Boxwood, Japanese**, (*Buxus microphylla*) Japonica; **Boxwood, Weller** (*Buxus sempervirens*); **Broom** (*Cytisus* spp.) Moonlight; **Broom** (*Genista pilosa*) Vancouver Gold; **Bugle Carpet**; **Camellia, Bugle Carpet** (*Camellia japonica*) Debutante, Maholiana Supreme, Chansonette, (*Camellia sasangua*); **Candy Tuft** (*Iberis* spp.) Snow White; **Carex, Variegated** (*Carex*); **Cedar, Red** (*Juniperus virginiana*); **Celosia** (*Celosia* spp.); **Centaurea** (*Centaurea montana*) Scarlet Plumosa; **Cockscomb, Plumosa** (*Celosia cristata*) Red Kewpie; **Coleus** (*Coleus blumei*); **Columbine** (*Aquilegia* spp.); **Copper leaf** (*Acalypha wilkesiana*); **Coreopsis** (*Coreopsis* spp.) Moonbloom; **Cornflower** (*Centaurea* spp.); **Cotoneaster** (*Cotoneaster apiculatus*); **Coyotebrush** (*Baccharis pelularis*); **Cycads** (*Cycads revoluta*); **Cypress, Bald** (*Taxodium distichum*); **Cypress, Italian** (*Cupressus sempervirens*) Glauca; **Cypress, Japanese False** (*Chamaecyparis obtusa*) Gracilis; **Cypress, Leyland** (*Cupressocyparis leylandii*); **Daffodil** (*Narcissus* spp.) King Alfred; **Daylily** (*Hemerocallis* spp.) Aztec Gold, Bright Yellow (Hybrid), Single Gold (Evergreen), Wilson's Yellow; **Delphinium** (*Delphinium* spp.) Magic Fountain; **Dianthus** (*Sweet William*) (*Dianthus* spp.); **Dogwood** (*Cornus florida*); **Dogwood, American** (*Cornus sericea*) Flavarinaeae; **Douglas Fir** (*Pseudotsuga menziesii*); **Dusty Miller** (*Senecio cineraria*); **Elm, Drake** (*Ulmus parvifolia*); **Euonymus** (*Euonymus fortunei*) Argenteo-variegata, Aureo-marginata, Colorata, Emerald Gaiety, Emerald 'N Gold, Gold Edge, Gold Princess, Silver King, Tricolor, Vegetus; **Fan Palm, European** (*Chamaerops humilis*); **Fan Palm, Mexican** (*Washingtonia robusta*); **Fern** (various) (*Asparagus* spp.); **Fescue** (*Festuca glauca*); **Fetterbush** (*Leucothoe fontana*) Rainbow; **Ficus** (*Ficus retusa*) Nitida; **Fir Fraser** (*Abies fraseri*); **Forsythia** (*Forsythia X intermedia*) Arnold Dwarf, Lynwood Gold, Meadowlark, (*Spring Glory*) Weeping; **Fuchsia** (*Fuchsia* spp.); **Galium** (*Galium odoratum*); **Gardenia** (*Gardenia jasminoides*) Mystery, Radicans; **Geranium** (*Pelargonium X hortorum*); **Gum** (*Eucalyptus citriodora*); **Hawthorn** (*Crataegus* spp.) Cockspear White, Crimson Cloud, Enchantress, Jack Evans, Washington White; **Heather, Twisted** (*Erica cinerea*) Mediterranean Pink; **Hemlock, Canada** (*Tsuga canadensis*); **Hibiscus** (*Hibiscus* spp.) Blue Bird, Brilliant, Hula Girl, Holly (*Ilex* spp.) Blue Boy, Blue Girl, (*Ilex X meserveae*) Burfordii, China Girl, Compacia, Forsteri, Hellerie, Japanese Northern Beauty, Needlepoint, Nellie R. Stevens, (*Ilex X attenuata*) Savannah; **Holly, Chinese** (*Ilex cornuta*); **Holly, Japanese** (*Ilex crenata*); **Holly, Yaupon** (*Ilex vomitoria*); **Honeysuckle** (*Lonicera japonica*) Clavey's Dwarf, Hilliana, Tatarian Canadian White, Zebelli Red Hosta; **Hosta** (*Hosta* spp.) Albo Marginata (*Hosta lancifolia*); **Ice Plant** (*Carpobrotus edulis*); **Impatiens** (*Impatiens* spp.); **Iris** (*Iris* spp.) Wedgewood, Dwarf Blue; **Ivy, English** (*Hedera helix*) Bulgaria, Thorndale; **Jasmine, Asian** (*Trachelopermum asiaticum*); **Juniper** (*Juniperus* spp.) Arcadia, Armstrong, Bar Harbor (*Juniperus horizontalis*) Blue Chip, Blue Pacific, Blue Rug, Blue Star, Broadmoor, Buffalo, Calgary Carpet, Emerald Sea, Emerald Spreader, Endora Compacta, Fruitlandi, Green, Gold Tip, Hetzi, Hughes, Manhattan Blue, Parsoni, Pfitzeriana, Plumosa, Prince of Wales, Procumbens Dwarf, San Jose, Sargent Blue, Sargent Green, Scandia, Scopulorum Moonglow, Scopulorum Skyrocket, Spartan (*Juniperus chinensis*) Tamariscifolia (*Juniperus sabina*) Weberi, Youngstown, Yukon Belle; **King Palm** (*Archontophoenix cunninghamiana*); **Laurel, Australian** (*Pittosporum tobira*); **Laurel, Mountain** (*Kalmia latifolia*); **Leucothoe** (*Leucothoe fontanesiana*); **Ligustrum, Japanese** (*Ligustrum japonicum*); **Lily, African** (*Agapanthus africanus*) Albus, Peter Pan; **Lily, African Blue**; **Lily of the Valley** (*Piety japonica*) Mt. Fire; **Lilyturf** (*Liriope muscaris*) Evergreen Giant, Lilac Beauty, Majestic, Monroe White, Variegata; **Liriope, Green** (*Liriope spicata*); **Maple, Japanese** (*Acer japonicum*); **Maple, Norway** (*Acer platanoides*); **Maple, Red** (*Acer rubrum*); **Maple, Silver** (*Acer saccharinum*); **Maple Sugar** (*Acer saccharum*); **Marigold** (*Tagetes patula*) Honeycomb, Wheeler's Dwarf, Variegata; **Mock Orange*** (*Philadelphus* spp.) Golden, Snowflake Double White; **Mondo Grass** (*Ophiopogon japonicus*); **Moss Rose** (*Portulaca grandiflora*) Sunnyside; **Myrtle, Crape** (*Lagerstroemia indica*) Faurei, Langer, Muskogee, Standard Pink; **Myrtle, Wax** (*Myrica cerifer*); **Nadina** (*Nadina domestica*) Compacta, Nana; **Narcissus** (*Narcissus* spp.) Oak, Laurel (*Quercus laurifolia*); **Oak, Pin** (*Quercus palustris*); **Oak, Red** (*Quercus rubra*); **Oak, Southern** (*Quercus virginiana*); **Oak, Willow** (*Quercus phellos*); **Oleander** (*Nerium oleander*) Hardy Red, Petite Pink, Sister Agnes; **Oleosperrum** (*Osteospermum fruticosum*) Wirligig; **Pachysandra** (*Pachysandra terminalis*); **Palm, Bangalow**; **Pampas Grass** (*Cortadaria selloana*); **Pansy** (*Viola X wittrockiana*); **Paper Flower** (*Bougainvillea glabra*) Barbara Karst; **Peach*** (*Prunus persica*); **Periwinkle, Dwarf** (*Vinca minor*); **Petunia** (*Petunia X hybrida*) Picotti; **Photinia, Red Tip** (*Photinia X fraseri*); **Pieris** (*Pieris japonica*); **Pine, Australian** (*Pinus nigra*); **Pine, Japanese Black** (*Pinus thunbergiana*); **Pine, Loblolly** (*Pinus taeda*); **Pine, Longleaf** (*Pinus palustris*); **Pine, Mugho** (*Pinus mugho*); **Pine, Scotch** (*Pinus sylvestris*); **Pine, Slash** (*Pinus elliotii*); **Pine, Swiss Mt.** (*Pinus mugo*); **Pine, Virginia** (*Pinus virginiana*); **Pine, White** (*Pinus strobus*); **Pineapple, Guava** (*Fajoaellowaiiana*); **Pittosporum**, Japan; **Potentilla** (*Potentilla nepalensis*) Abbotswood (*Potentilla fruticosa*); **Privet** (*Ligustrum japonicum*) Golden Vicary, Regal, Texanum, Wax, Yellow Tipped; **Privet, Glossy** (*Ligustrum lucidum*); **Pyraanthia** (*Pyraanthia koidzumii*) Gnome, Lalandei, Victory; **Queen Palm** (*Areacstrum rammannzoffianum*); **Quince, Japanese***; **Rhododendron** (*Rhododendron* spp.) Album, Cunningham White, Fashion, Hardy, PJM, Purple Gem, Silvery Pink; **Rhododendron, Carolina** (*Rhododendron carolinianum*); **Rhododendron, Catawba** (*Rhododendron catawbiense*); **Rockcress** (*Arabis* spp.) Snowcap; **Rhode Max** (*Rosebay*) (*Rhododendron maximum*); **Rose** (*Rosa banksiae*) Luta; **Rosemary*** (*Rosmarinus officinalis*); **Rosemary, Bog** (*Andromeda polifolia*) Nana; **Salvia** (*Salvia farinacea*) Rhea; **Sedum** (*Sedum spurium*) Dragon Blood Red,

8088

Red Carpet, Yellow; Snapdragon (*Antirrhinum* spp.); Sourwood (*Oxydendrum arboreum*); Spiraea (*Astilbe X arendsii*) Fanali; Spiraea (*Spiraea* spp.) Anthony Waterer Red, Dolchica, Froebel Pink, Goldenflame Red, Snowmound White, Van Houtte White; Spiraea, Garland (*Spiraea* spp.); Spruce, Black Hills (*Picea glauca*); Spruce, Colorado Blue (*Picea pungens*); Spruce, Norway (*Picea abies*); Spruce, White (*Picea glauca*) Conica; Sweetflag, Grassyy-Le; Sweetgum (*Liquidambar styraciflua*); Sycamore (*Platanus occidentalis*); Tree Fern (Tiki Fern) (*Asparagus virgatus*); Trumpet Flower, Evening (*Gelsemium sempervirens*); Tulp (*Tulipa*, spp.) Apeldoorn; Verbena, Shrub (*Lantana sellowiana*); Vervain (*Verbena* spp.) St. Paul; Viburnum (*Viburnum* spp.) American Cranberry Bush, Arrowwood, Common Snowball, European Cranberry Bush, Linden, Mohican, Wright; Vinca (*Periwinkle*) (*Vinca minor*); Windmill Palm (*Trachycarpus fortunei*); Xylosma (*Xylosma congestum*); Yarrow (*Achillea* spp.); Yaupon (*Ilex vomitoria*) Dwarf; Yew (*Taxus cuspidata*) Denisformis (*Taxus X media*).

*Ornamental species only. DO NOT USE ON FOOD PRODUCING TREES AND ORNAMENTALS.

WEEDS CONTROLLED: Used as directed, BEST EverGreen 18-5-0 with 0.20% DIMENSION® will control or suppress the following annual grass and broadleaf weeds when applied prior to their emergence. This product will not control established grasses (except for crabgrass in early stages of development), and established broadleaf weeds, therefore, the area to be treated should be free of weeds prior to application.

Grasses: barley (*Hordeum* spp.); barmyard grass (*Enchinochloa crus-galli*); bluegrass, annual (*Poa annua*); brome (*Bromus* spp.); crabgrass, large (*Digitaria sanguinalis*); crabgrass, smooth (*Digitaria ischaemum*); crabgrass, Southern (*Digitaria ciliaris*) crowfootgrass (*Dactyloctenium aegyptium*); dallisgrass (seedling) (*Paspalum dilatatum*); goosegrass (*Eleusine indica*); foxtail, green (*Setaria verti*); foxtail, yellow (*Setaria faberii*); kikuyugrass (*Pennisetum clandestinum*) oats, wild (*Avena fatua*); ryegrass (annual & perennial) (*Lolium* spp.); smutgrass (*Sporobolus indicus*).

Broadleaf Weeds: bittercress (*Cardamine* spp.); carpetweed (*Mollugo verticillata*); chickweed (*Stellaria* spp.); geranium, Carolina (*Geranium carolinianum*); henbit (*Lamium* spp.); knotweed, prostrate (*Polygonum aviculare*); lespedeza, common (*Lepedeza striata*); maretail (*Conyza canadensis*); medic, black (*Medicago lupulina*); mustard (*Brassica* spp.); oxalis, buttercup (*Oxalis pes-caprae*); pineappleweed (*Matricaria matricarioides*); pigweed, redroot (*Amaranthus retroflexus*); parsley-piert (*Alochemilla arvensis*); purslane, common (*Portulaca oleracea*); rocket, London (*Sisymbrium irio*); shepherd's purse (*Capsella bursa-pastoris*); speedwell, com (*Veronica arvensis*); spurge, garden (*Euphorbia hirta*); spurge, prostrate (*Euphorbia humistrata*); spurge, spotted (*Euphorbia maculata*); woodsorrel, creeping (*Oxalis corniculata*); woodsorrel, yellow (*Oxalis stricta*).

APPLICATION EQUIPMENT AND INSTRUCTIONS

Apply this product with drop or rotary-type spreaders designed to apply granular herbicides. Avoid the use of spreaders that would apply this product in narrow rows or concentrated bands. Before each application, calibrate the spreader according to the equipment manufacturer's directions for adjusting the spreader settings such that the spreader delivers the appropriate application rate recommended above. Apply this product uniformly over the area to be treated. A more uniform application can be made by spreading half the required amount over the area and applying the remaining half at a right angle to the previous direction. Avoid streaking, skips, or overlaps during application. Check equipment frequently to ensure equipment is functioning properly and delivering uniform distribution of granules.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store this product only in its original container in a dry, cool, secured storage area.

SPILL: In case of spill, sweep up product and dispose of material according to "product disposal" directions listed below.

PRODUCT DISPOSAL: Wastes resulting from the use of this product that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticide disposal or in accordance with applicable Federal, State or local procedures.

EMPTY CONTAINER DISPOSAL: 50 LB. BAGS - Do not reuse bag. Completely empty bag into application equipment. Dispose of empty bag(s) in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke. BULK CONTAINERS (Woven polyethylene - 1,000 and 2,000 lbs.) - Empty thoroughly and return to specified destination for cleaning and reuse.

CONDITIONS OF SALE AND WARRANTY

J.R. Simplot Company warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label only when used in accordance with label directions under normal conditions of use. J.R. SIMPLOT COMPANY MAKES NO OTHER EXPRESS OR IMPLIED WARRANTIES EITHER OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR USE. Handling, storage and use of the product by Buyer or User are beyond the control of J.R. Simplot Company and Seller. Risks such as crop injury, ineffectiveness or other unintended consequences resulting from, but not limited to, weather or soil conditions, presence of other materials, disease, pest, drift to other crops or property or failure to follow label directions will be assumed by the Buyer or User. In no case will J.R. Simplot Company or Seller be held liable for consequential, special or indirect damages resulting from the handling, storage or use of this product.

DIMENSION® is a registered trademark of Dow AgroSciences LLC.

0303(01)

This product contains a chemical known to the State of California to cause cancer, birth defects or other reproductive harm. Proposition 65, the Safe Drinking Water and Toxic Enforcement Act of 1986, requires notification of potential exposure to substances identified by the State of California as causing cancer, birth defects or other reproductive harm.

Information regarding the contents and levels of metals in this product is available on the Internet at <http://www.regulatory-info-jr.com>

ITEM#79647050 EverGreen 18-5-0 w/DIM (LABELS/BEST/DIM R11-09-09)

Simplot

MANUFACTURED IN U.S.A.
J. R. SIMPLOT COMPANY
P.O. Box 198 • Lathrop, CA 95330



Tech sheets, MSDS and other
information on BEST products
available at www.bestfertilizer.com

NET WEIGHT 50 LB. BAG (22.68 Kg.)