

7001-377

1/24/2003

Page 176



Please read instructions on reverse before completing form.

Form Approved, OMB No. 2070-0060, Approval expires 2-28-95



United States
Environmental Protection Agency
Washington, DC 20460

☐ Registration
☐ Amendment
☒ Other

OPP Identifier Number

Application for Pesticide - Section I

1. Company/Product Number 7001-377	2. EPA Product Manager Joanne Miller	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) Turf Fertilizer with .107% Dimension	PM# 23	
5. Name and Address of Applicant (Include ZIP Code) J.R. Simplot Company P.O. Box 198 Lathrop, CA 95330 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	

Section - II

<input type="checkbox"/> Amendment - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below.

NOTIFICATION

JAN 24 2003

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

Notification of Adding, Revising or Deleting Advisory Statements per PR Notice 98-10.

This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.

Section - III

1. Material This Product Will Be Packaged In:			
Child-Resistant Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	2. Type of Container <input type="checkbox"/> Metal <input type="checkbox"/> Plastic <input checked="" type="checkbox"/> Glass <input type="checkbox"/> Paper <input type="checkbox"/> Other (Specify) _____
* Certification must be submitted		If "Yes" Unit Packaging wgt. No. per container	If "Yes" Package wgt No. per container
3. Location of Net Contents Information <input checked="" type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container 50 and 2000 lb.	
5. Location of Label Directions <input type="checkbox"/>		6. Manner in Which Label is Affixed to Product <input checked="" type="checkbox"/> Lithograph <input checked="" type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled <input type="checkbox"/> Other _____	

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)			
Name Lisa J. Strong		Title Regulatory Consultant	Telephone No. (Include Area Code) (209) 858-2511
Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment both under applicable law.			6. Date Application Received (Stamped)
2. Signature 		3. Title Regulatory Consultant	
4. Typed Name Lisa J. Strong		5. Date 12/9/02	

2 7 6



AgriBusiness

J.R. SIMPLOT COMPANY 16777 HOWLAND ROAD P.O. BOX 198
LATHROP, CALIFORNIA 95330-0198 (209) 858-2511 FAX (209) 858-2519

December 9, 2002

Document Processing Desk (NOTIF)
Office of Pesticide Program (7504C)
U.S. Environmental Protection Agency
Room 266A, Crystal Mall 2
1921 Jefferson Davis Hwy.
Arlington, VA 22202

ATTN: Ms. Joanne Miller
Product Manager (23)
And
Mr. Eugene Wilson

RE: EPA Reg. No. 7001-377
Turf Fertilizer with .107% Dimension


This serves as notification that, under the Directions For Use, J.R. Simplot Company is removing the following statement "In the State of New York do not use in Nassau County or Suffolk County" and replacing it with "In New York State this product may only be used by Professional Turf Applicators at no more than 465 lbs. (0.5 lbs. active ingredient) per acre per year and is prohibited from use in Nassau and Suffolk Counties."


As directed in PR Notice 98-10 we followed the guidelines established in PR Notice 95-2 for adding, revising or deleting advisory statements.

After your review and acceptance we request a pin-punched copy of the proposed label be returned to us in order to satisfy state registrations. We have enclosed a self-addressed stamped envelope in order to facilitate this request.

Five copies of our label are enclosed for your review. One copy highlights the changes.

Sincerely,
J.R. SIMPLOT COMPANY


Lisa J. Strong
Regulatory Consultant





TURF FERTILIZER WITH .107% DIMENSION®

PREEMERGENCE WEED CONTROL with PLANT FOOD

Turf Fertilizer with .107% Dimension® is a selective herbicide that provides control of listed annual grasses and broadleaf weeds in established lawns and ornamental turfs, including golf course fairways, roughs, and tee boxes. Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

ACTIVE INGREDIENT:

Dithiopyr, 3,5-pyridinedicarbothioic acid, 2-(difluoromethyl)-4-(2-methylpropyl)-6-(trifluoromethyl)-S,S-dimethyl ester

0.107%

INERT INGREDIENTS:

99.893%

Total 100.000%

Product protected by U.S. Patent No. 4,692,184. Other patents pending.

EPA Reg. No. 7001-377

EPA Est. No. 7001-MO-1

KEEP OUT OF REACH OF CHILDREN

CAUTION

NOTIFICATION
JAN 24 2003

NOTIFICATION
JAN 24 2003
326

READ THE ENTIRE LABEL BEFORE USING THIS PRODUCT.

Use only according to label instructions.

NOTICE: Before using this product, read the entire Precautionary Statements, Conditions of Sale and Warranty, Directions for Use, Use Restrictions and Storage and Disposal Instructions. If the Conditions of Sale and Warranty are not acceptable, return the product unopened within thirty days of purchase to the place of purchase.

FIRST AID

IF IN EYES-Immediately flush with plenty of water. Get medical attention if irritation persists.

IF INHALED-Remove individual to fresh air. Seek medical attention if breathing difficulty develops.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION! CAUSES EYE IRRITATION. HARMFUL IF INHALED. Avoid contact with eyes or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Prolonged or frequently repeated skin contact while handling this material may cause allergic reaction in some individuals.

X-X-X

GUARANTEED ANALYSIS

**REFER TO THE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS
AND DIRECTIONS FOR USE.**

NET WEIGHT: 50 lbs. (22.68 Kg)



486

Turf Fertilizer With .107% DIMENSION[®]

ENVIRONMENTAL HAZARDS

This product is toxic to fish and highly toxic to other aquatic organisms including oysters and shrimp. Use with care when applying to turf areas adjacent to any body of water. Drift and runoff from treated turf may adversely affect aquatic organisms in adjacent aquatic sites. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling in New York State. This product may only be used by Professional Turf Applicators at no more than 465 lbs. (0.5 lbs. active ingredient) per acre per year and is prohibited from use in Nassau and Suffolk Counties.

GENERAL INFORMATION

This product is a selective herbicide that provides control of crabgrass and other annual grasses and broadleaf weeds in established lawns and ornamental turfs, including golf course fairways, roughs, and tee boxes. This product will not harm nearby established ornamentals when used according to label directions.

USE PRECAUTIONS

Turfgrass Safety

This product may be used on seeded, sodded, or sprigged lawns and ornamental turfs that are well established. The grass must have developed a good root system and a uniform stand, and have received at least two mowings following its seeding, sodding, or sprigging before it can receive its first application of this product. Use of this product on lawns and ornamental turfs that are not well established, or on those that have been weakened by weather, pest, disease, chemical, or mechanical-related stress, may increase the chances for turf injury.

This product should only be applied to lawns and ornamental turfs that are composed of the following turfgrass species that have been determined to be tolerant to applications of this product. When applied as directed under the conditions described, the following established turfgrasses are tolerant to this product.

Cool Season Grasses		Warm Season Grasses	
Common Name	Scientific Name	Common Name	Scientific Name
Bentgrass, Creeping	<i>Agrostis palustris</i>	Bermudagrass	<i>Cynodon dactylon</i>
Bluegrass, Kentucky	<i>Poa pratensis</i>	Bulldozergrass	<i>Echinochloa polystachya</i>
Fescue, fine-leaved	<i>Festuca rubra</i>	Carpetgrass	<i>Axonopus affinis</i>
Fescue, tall	<i>Festuca arundinacea</i>	Centipedegrass	<i>Eremochloa ophiuroides</i>
Ryegrass, perennial	<i>Lolium perenne</i>	Kikuyugrass	<i>Pennisetum clandestinum</i>
		St. Augustinegrass	<i>Stenotaphrum secundatum</i>
		Zoysiagrass	<i>Zoysia japonica</i>

TURFGRASS REGIONS

- Northern States
- Transition States
- Southern States
- Coastal Southern



Subsequent, sequential pre- and/or postemergence applications should be made where longer periods of control are desired.

Where split fertilizer applications are recommended the rates in Table 1 may be split across two applications made 6-10 weeks apart and prior to crabgrass emergence.

Table 1. Recommended Single Application Use Rates ^a			
(Pounds of this product/1,000 sq. ft.)			
PROGRAM	#1 Preemergence (high-cut turf)	#2 Preemergence (low-cut turf)	#3 Postemergence (low-cut turf)
USE (Turfgrass cut)			
REGION			
North	2.68 lbs./1,000 sq. ft.	3.86 lbs./1,000 sq. ft.	5.36 lbs./1,000 sq. ft.
Transition	3.86 lbs./1,000 sq. ft.	5.36 lbs./1,000 sq. ft.	7.93 lbs./1,000 sq. ft.
South	5.36 lbs./1,000 sq. ft.	7.93 lbs./1,000 sq. ft.	10.72 lbs./1,000 sq. ft.
Coastal South	7.93 lbs./1,000 sq. ft.	10.72 lbs./1,000 sq. ft.	10.72 lbs./1,000 sq. ft.

^aDO NOT apply more than 10.72 lbs. of this product/1,000 sq. ft. per application, and no more than 32.16 pounds of this product/1,000 sq. ft. per year. Postemergence control is limited; see "Control of Crabgrass" and "Use Precautions" sections. Regions include areas listed below. See Map of United States.

North:

all other areas
DE, KS, KY, MD, MO, MI, VA, southeastern PA, southern areas of IL, IN, OH, & coastal areas of CT, NY, & RI

South:

AL, AR, AZ, CA, GA, LA, MS, NC, NM, NV, OK, SC, TN, & TX
FL, & southern coastal areas of AL, GA, LA, MS, NC, SC, & TX

Coastal South:

FL, & southern coastal areas of AL, GA, LA, MS, NC, SC, & TX

^bMay require split or sequential applications for full control.

FALL APPLICATIONS

This product can also be applied in the late summer or early fall (late August through November) at the "Program 3" use rates listed in Table 1 to provide control of crabgrass through the early part of the next spring. The fall application should be followed by an appropriate spring application to provide season-long control.

TIPS FOR IMPROVED CONTROL

For best results, apply this product within a few days after mowing and delay mowing again for a few days after the application. When treated lawn or ornamental turf areas are watered or receive significant rainfall within a few days after application of this product, improved weed control may result.

Use of split (half-rate) applications spaced 6-8 weeks apart may provide improved weed control.

Control of Other Grass and Broadleaf Weeds

Use of split (half-rate) applications spaced 6-8 weeks apart may provide improved weed control. Spring Applications. Used as directed for crabgrass control in the spring, this product will also control (at the "Program 3" rates) or suppress (at the "Program 2" rates) the following weeds when applied prior to their emergence:

- barnyardgrass
- bluegrass (annual)
- crowfootgrass
- foxtail (yellow & green)
- goosegrass
- kikuyugrass
- ryegrass (annual & perennial)
- smutgrass
- lespedeza (common)
- oxalis (buttercup, creeping & yellow wood-sorrel)
- purplestem (common)
- speckleweed (corn)
- spurge (prostrate & spotted)

^aAlso controlled at the Program 2 rates.

Fall Applications. Used as directed for late summer or early fall use, for crabgrass control through the early part of the next spring, this product will also control (at the "Program 3" rates) or suppress (at the "Program 2" rates) the following weeds when applied prior to their emergence:

- bluegrass (annual)
- chickweed
- geranium (Carolina)

shepherdspurse

DO NOT apply this product to Colonial bentgrass (*Agrostis tenuis*) varieties.

Use of this product on certain varieties of creeping bentgrass such as 'Cohansey', 'Carmen', 'Seaside', and 'Washington' may result in undesirable turfgrass injury. Not all varieties of creeping bentgrass have been tested. **Use** of this product on certain varieties of fine fescue may result in undesirable turf injury. The following fine fescue varieties have been found to be sensitive to this product: 'Atlanta', 'Banner', 'Beauty', 'Bigart', 'CF-2', 'Enjoy', 'HF-93', 'highlight', 'Ivalo', 'Jamestown', 'Koket', 'Mejenta', 'Mary', 'Pennlawn', 'Tamara', 'Tatjana', 'Waldorf', and 'Waldina'. Not all varieties of fine fescue have been tested. **DO NOT USE** this product on seedling buffalograss in the spring of the first year of establishment until the turfgrass is fully green and has established new roots.

RESEEDING, OVERSEEDING, OR SPRIGGING

Reseeding, overseeding, or sprigging of treated areas within 4 months after application of this product may inhibit the establishment of desirable turfgrasses. When reseeding or overseeding, proper cultural practices such as soil cultivation, irrigation and fertilization should be followed. For best results, use mechanical or power seeding equipment (slit seeders) designed to give good seed to soil contact.

GOLF COURSE PUTTING GREENS

DO NOT use this product on golf course putting greens.

OTHER USE PRECAUTIONS

The early postemergence crabgrass control described below is limited. Postemergence applications are only effective on crabgrass if applied before crabgrass has displayed its fifth leaf or first tiller. For best results, cultural practices that disturb the soil, such as core-, spike-, or hydro-aerification, and verticutting, should be done before application of this product. DO NOT apply this product until the grass has recovered from these cultural practices.

Apply this product directly to established lawns or ornamental turfs only. DO NOT apply to flowers, vegetables, shrubs or trees. DO NOT use clippings from treated turf for mulching around vegetables or fruit trees. DO NOT apply this product to pastures. Keep people and pets off treated areas until dust has settled.

APPLICATION DIRECTIONS

Control of Crabgrass - Preemergence and Early Postemergence Control.

This product provides "preemergence" control of crabgrass (including the large, smooth, and southern species) when applied prior to the emergence of crabgrass from the ground in established lawns and ornamental turfs.

It can also provide "early postemergence" control of crabgrass during the early stages of crabgrass growth after the crabgrass has emerged from the ground. However, it is often difficult to see the very small, early stages of crabgrass in well-established lawns and ornamental turfs. Postemergence crabgrass control will be obtained only when this product is applied prior to the tillering of crabgrass, which generally corresponds to the time when you can first easily see the crabgrass plants in the lawn or turf. So the practical benefit of this product's additional, early postemergence activity is that (compared to strictly preemergence crabgrass products) it can give the user a 2-4 week longer period of time to make applications and still control crabgrass.

APPLICATION AND FREQUENCY TIMING

This product may be applied as a single application, as a split application, or as a sequential application for crabgrass control in the spring, summer or fall. DO NOT apply more than 10.72 pounds of this product per 1,000 sq. ft. per application, and no more than 32.16 pounds of this product per 1,000 sq. ft. per year.

SPRING APPLICATIONS

For single applications made in the spring or early summer, this product should be applied at the appropriate rate corresponding to one of the three control programs listed in Table 1 below, depending on the user's location, the turfgrass mowing height, and whether the use is considered to be preemergence or early postemergence at the time of the application. The duration of residual weed control provided by this product is directly related to the total rate applied, but will vary somewhat depending on weather, weed pressure, turfgrass competitiveness, and the user's location within a region.

Use Program #1 for preemergence control at sites where the turfgrass is cut high (e.g., homeowner lawns). This program provides 3-5 months of preemergence crabgrass control. This program should not be used for early postemergence crabgrass control.

Use Program #2 for preemergence control at sites where (a) turfgrass is cut low (e.g., golf fairways), and (b) turfgrass maintenance or weed control has not been conducted during the previous year. This program provides 4-6 months of preemergence crabgrass control. This program may also be used for early postemergence control at sites where turfgrass is cut high (e.g., homeowner lawns).

Use Program #3 for preemergence control at sites where (a) turfgrass is cut low (e.g., golf fairways) and (b) turf maintenance or weed control has not been conducted during the previous year. This program provides 4-6 months of preemergence crabgrass control. This program may also be used for early postemergence control at sites where turfgrass is cut low (e.g., golf fairways).

APPLICATION EQUIPMENT AND INSTRUCTIONS

Apply this product with drop or rotary-type spreaders designed to apply granular herbicides. Avoid the use of spreaders that would apply this product in narrow rows or concentrated bands. Before each application, calibrate the spreader according to the equipment manufacturer's directions for adjusting the spreader settings such that the spreader delivers the appropriate application rate recommended above. Apply this product uniformly over the lawn or ornamental turf area. A more uniform application can be made by spreading half of the required amount over the area and then applying the remaining half at a right angle to the previous direction. Avoid streaking, skips or overlaps during application. Check equipment frequently to ensure equipment is functioning properly and applying uniform distribution of granules.

STORAGE AND DISPOSAL

STORAGE: Store this product only in its original container in a dry, cool, secured storage area. Do not contaminate water, foodstuffs, feed or seed by storage or disposal.

PRODUCT DISPOSAL: Wastes resulting from the use of this product that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticide disposal or in accordance with applicable Federal, State or local procedures.

EMPTY CONTAINER DISPOSAL: Do not reuse bag. Completely empty bag into application equipment. Dispose of emptied bag(s) in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

SPILL: In case of spill, sweep up material and dispose of material according to "product disposal" directions listed above.

**Dimension® is a registered trademark of Rohm and Haas Company.*

CONDITIONS OF SALE AND WARRANTY

J.R. SIMPLOT COMPANY warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label only when used in accordance with label directions under normal conditions of use. J.R. SIMPLOT COMPANY MAKES NO OTHER EXPRESS OR IMPLIED WARRANTIES EITHER OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR USE. Handling, storage and use of the product by Buyer or User are beyond the control of J.R. SIMPLOT COMPANY and Seller. Risks such as crop injury, ineffectiveness or other unintended consequences resulting from, but not limited to, weather or soil conditions, presence of other materials, disease, pest, drift to other crops or property or failure to follow label directions will be assumed by the Buyer or User. IN NO CASE WILL J.R. SIMPLOT COMPANY OR SELLER BE HELD LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE HANDLING, STORAGE OR USE OF THIS PRODUCT.

DIM 107 (LABELS/ St Louis/Rev. A 11-02)

NET WEIGHT 50lbs. (22.68kg)

Distributed By:

Simplot

J. R. SIMPLOT COMPANY

101 Ferry St., St. Louis, MO 63147

P.O. Box 198, Lathrop, CA 95330



6
9
2
9