

U.S. ENVIRONMENTAL PROTECTION AGENC Office of Pesticide Programs Registration Division (7505C) 1200 Pennsylvania Ave., N.W. Washington, D.C. 20460

EPA Reg Number

Date of Issuance: JUL 3 1 2002

Term of Issuance:

7001-382

Conditional

NOTICE OF PESTICIDE:

<u>X</u> Registration

__ Reregistration

Name of Pesticide Product:

PCNB 12.5%

Name and Address of Registrant (include ZIP Code):

J.R. Simplot Company

P.O. Box 198

Lathrop, CA 95330

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA Sec. 3(c)(7)(A) provided that you:

- 1. Submit and/or cite all data required for registration/ reregistration of your product under FIFRA Sec. 3(c)(5) when the Agency requires all registrants of similar products to submit such data; and submit acceptable responses required for reregistration of your product under FIFRA section 4.
- 2. Make the following labeling changes before you release the product for shipment:

Add the phrase "EPA Registration Number 7001-382".

3. Submit one copy of your final printed label prior to release of the product for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA Sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records.

Many J. Walles Waller, Product Manager (21)

7/31/02

Fungicide Branch, Registration Division (7505C)

PCNB 12.5%

With X-X-X Fertilizer

Aids in the Control and Prevention of Stripe Smut, Leaf Spot, Brown Patch, Dollar Spot and Snow Mold on a variety of turfgrasses.

ACTIVE INGREDIENT:

EPA Reg. No. 7001-GIE EPA Est. No. 7001-MO-1

KEEP OUT OF REACH OF CHILDREN CAUTION

FIRST AID

IF ON SKIN OR CLOTHING Take off contaminated clothing.

Rinse skin immediately with plenty of water for 15-20 minutes Call a poison control center or doctor for treatment advice.

IF IN EYES

Hold eye open and rinse slowly and gently with water for 15-20 minutes.

Remove contact lenses, if present, after the first 5 minutes, then continue rinsing

Call a poison control center or doctor for treatment advice.

IF INHALED

Move person to fresh air.

If person is not breathing, call 911 or an ambulance, then give artificial

respiration, preferably by mouth-to-mouth, if possible.

Call a poison control center or doctor for further treatment advice.

IF SWALLOWED

Call a poison control center or doctor immediately for treatment advice.

Have person sip a glass of water if able to swallow.

Do not induce vomiting unless told to do so by the poison control center

or doctor.

Do not give anything by mouth to an unconscious person.

HAVE THE PRODUCT CONTAINER OR LABEL WITH YOU WHEN CALLING A POISON CONTROL CENTER OR DOCTOR, OR GOING FOR TREATMENT.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Harmful if absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking chewing gum, or using tobacco. Wear long sleeved shirt and long pants, socks, shoes and gloves. Remove and wash contaminated clothing before reuse.

NET WEIGHT: 50 lbs. (22.68 Kg.)

ACCEPTED

JUL 3 1 2002

Under the Federal Issections KFA Reg. No.

NET WEIGHT: 1000 lbs. (453.6 Kg) NET WEIGHT: 2000 lbs. (907.20 Kg)

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Do not apply directly to water, to areas where water is present or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water when cleaning equipment or disposal of waste. Do not apply directly adjacent to potable water supplies.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

Do not apply this product through any type of irrigation system.

Do not graze treated areas or feed clippings to livestock.

USE PRECAUTIONS

Use on lawn and turfgrass areas only. Do not adjust application rates to meet fertilizer requirements for your soil. USE ONLY when both a fertilizer and fungicide are required. Apply when foliage is dry. If temperature is 80°F or above, water lightly to wash granules off foliage. Very slight but temporary yellowing of the grass may occur following treatment. This causes no harm and will disappear in a short time.

Cohansey Bentgrass is more sensitive to PCNB than other Bentgrass varieties. Use on Bentgrass only from mid-fall through early winter, except in California, Oregon and Washington where application can be made after early winter through the early fall. Not for use on dichondra or stands of predominately fine fescue. Not for use on tall fescue during the warm summer months in the Southern U.S.A. Not for use on Bermudagrass in Florida, Southern Alabama, Southern Mississippi, Louisiana, Southern Texas and California.

Not for use on winter overseeded Bermudagrass putting greens until seedlings are established. Not for use on Poa Annua where this turf is desired in the Southern part of the U.S.A.

Treat Bluegrass only in the fall through early spring and Bentgrass only in mid-fall through early winter In California, apply on Bluegrass and Bentgrass turf.

Not for use on Velvet Bentgrass.

FERTILIZER GUARANTEE X-X-X

SPREADER SETTINGS

Rotary spreaders should be used with caution. Do not apply on a windy day or under conditions where material comes in contact with the operator.

COOL SEASON GRASSES

BROWN PATCH, DOLLAR SPOT Northern or Cool Season lawns such as bluegrass, bluegrass-fine fescue mixtures, Perennial Ryegrass and similar varieties (CA apply to Bluegrass varieties only.) Apply 1.6 - 2 lbs. per 1,000 sq. ft. to dry foliage at the first sign of disease. Repeat every 7 to 10 days during warm, damp weather.

SNOW MOLD (Except CA, OR, WA) on Bluegrass, Bluegrass blends with Fine Fescue or Perennial Ryegrass, Bentgrass or Bentgrass mixtures and similar varieties. Apply 4 to 6 lbs. prior to the first heavy snow. Use 6 lbs. per 1,000 sq. ft. in areas where snow cover is present the entire winter. On Bentgrass and on putting greens and tees containing Bentgrass, make one application in the mid-fall through early

winter.

SNOW MOLD (CA, OR, WA) on Bluegrass, Bluegrass blends with Fine Fescue or Perennial Ryegrass, Bentgrass or Bentgrass mixtures and similar varieties: Apply 4 to 6 lbs. per 1,000 sq. ft. to dry foliage through the winter months. Wait a minimum of 14 days between applications. Make one application in areas where snow cover is present the entire winter. On Bentgrass and on putting greens and tees containing Bentgrass, make one application in the mid-fall through early winter.

STRIP SMUT (Except CA) on Bluegrass and Perennial Ryegrass Apply 6.1bs. per 1,000 sq. ft. at first sign of disease. If the disease is still present, make a second application in 3-4 weeks. As a preventative treatment apply 6 lbs. in the fall and spring.

STRIP SMUT (CA only) on Bluegrass: Apply 6 lbs. per 1,000 sq. ft. at first sign of disease. If the disease is still present in 3-4 weeks, make a second application. Do not apply more than 3 applications per year. As a Preventative treatment apply 6 lbs. per 1,000 sq. ft. in the fall through early spring. Do not apply more than 3 applications per year.

RUST on Bluegrass and Perennial Ryegrass. Apply 6 lbs. per 1,000 sq. ft. at first sign of disease. If the disease is still present in 3-4 weeks, make a second application.

WARM SEASON GRASSES

BROWN PATCH, DOLLAR SPOT: St. Augustine, Bermuda grass, Zoysia and similar varieties. Apply 6 lbs. per 1,000 sq. ft. to dry foliage in the fall as a preventative, or in the spring at the first sign of disease. Water the area to wash the granules to soil level. An additional application in 3 to 4 weeks may be needed if the disease is still present or if heavy rainfall or flooding occurs after the treatment.

DICHONDRA: Apply 6 lbs. per 1,000 sq. ft. to dry foliage and water lightly to insure movement to the soil level. Apply at first sign of disease. Repeat every 7 to 10 days during warm damp weather.

STORAGE AND DISPOSAL

PESTICIDE STORAGE: Do not contaminate water, food or feed by storage and disposal.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Paper and Plastic Bags. Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

BULK CONTAINERS: Woven Polyethylene (1000 and 2000 lbs.) Empty thoroughly and return to your local supplier for cleaning and reuse.

J.R. SIMPLOT COMPANY P.O. BOX 198 LATHROP, CA 95330