

**CONCENTRATED
FABRIC
SANITIZER**

Bio-Rinse G

**PROVIDES SELF-SANITIZING
ACTIVITY ON
TREATED FABRICS**

EPA REG. NO. 7048-6

DIRECTIONS

Use 1½ oz. per 100 lbs. of fabric in the final rinse after sour. For best results dilute dosage in 1-2 gallons of water and then add to final rinse before softener.

Bio-Rinse G is a special treatment to be used as an adjunct to standard laundering routine. It is not intended as a terminal disinfectant for fabric or hard surfaces at the concentrations recommended.

Do not reuse empty drum. Return to drum reconditioner or rinse well with soap solution and discard. Do not contaminate water by disposal of waste.

WHEN USED AS DIRECTED

Bio-Rinse G imparts a self-sanitizing residual to the fabric that reduces the number of *Staph* and *E. Coli* in the presence of moisture or high relative humidity.

Bio-Rinse G imparts a bacteriostatic residual to the fabric that will inhibit the growth of antibiotic resistant *staph* and enteric coliform organisms in the presence of moisture or high relative humidity.

Bio-Rinse G helps reduce the danger of cross infection due to contaminated linen.

Bio-Rinse G inhibits the growth of bacteria which produce ammonia in diapers.

ACCEPTED

11-15-71

UNDER THE FEDERAL INSECTICIDE
ACT AND REGULATIONS THEREOF
FOR RECORD NO. 7048-6

EDMAR CHEMICAL COMPANY

3939 VALLEY ROAD • CLEVELAND, OHIO 44109

DANGER

Keep out of reach of children

Corrosive. Causes severe eye and skin damage. Do not get in eyes, on skin, or on clothing. Wear goggles or face shield and rubber gloves when handling. Harmful or fatal if swallowed. Avoid contamination of food.

FIRST AID

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse.

If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution; or if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

NOTE TO PHYSICIAN

Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression, and convulsion may be needed.

ACTIVE INGREDIENTS:

n-Alkyl (60% C₁₄, 30% C₁₆, 5% C₁₂, 5% C₁₈)

dimethyl benzyl ammonium chlorides 12.5%

n-Alkyl (68% C₁₂, 32% C₁₄)

dimethyl ethylbenzyl ammonium chloride 12.5%

INERT INGREDIENTS: 75.0%

100.0%

NET CONTENTS—55 GALS.