DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Por use in homes, bakeries, food plants, restaurants, schools, hotels, motels, office buildings, hospitals, and food handling plants.

Remove protective cap, Hold container upright and point valve in desired direction and press button to sprey.

FOR EC. TITES, MOSQUITOES, AND GRATE: Close all doors and windows. Voint horsis upward, direct the spray to all parts of the room especially windows and other light sources which attract these insects. Fill the room with mist, then leave the treated area. Keep the room closed for at least 15 minutes. Ventilate the room when treatment is completed. Sweep up and destroy fallen insects. Repeat treatment as necessary.

WASPS: Application should be made in late evening when insects are at rest. Spray liberally into hiding and breeding places, contecting as many insects as possible. Repeat as mecessary.

FOR CRANLING INSECTS: Boaches, earwigs, grain mites, now bugs, spiders, and centipedes: spray thoroughly into hiding places such as cracks, crevices, moist areas, openiess around pipes and sinks, under refrigerators, baseboards, and storage areas. Spray directly on insects when possible. Repeat as necessary.

AMTS: Spray trails, sest and point of entry. Spray on ants when possible. Repeat as accusacry.

SPIDERS, CENTIFEDES, SON BUGS, CRICKETS: Spray directly on wahe and places where these peace crewl. Spray directly on insects when possible.

CARPET RESTLES, MOTH LAWVAR: Closm and air conteminated materials before treating. Spray directly on peats, repeating treatment as needed. Articles not stored in mothtight container should be retreated every 30 days.

SEESDOS: Sprey cracks and surfaces in basehoords, floors, moulding, and other contaminated areas. Take bed apart and treat mettress, springs and frame.

FLEAS: Spray floors, floor coverings, cracks, bedding, sleeping Tearters, and other erest frequented by pets. Repeat as necessar Do not spray pets.

CHEM SPRAY FORMULA NG. 1

RILLS RESISTANT PLIES AND ROACHES

Effective against flies, Roaches, Anta, Moths and Certain Other Insects

ACTIVE INGREDIENTS

Ayrethrins	C.3	1
Technical Piperenyl Butexide *	0.6	1
N-octyl Bicyclohoptone Dicarboximide	1.0	1
IMER: INGREDIENTS	98.1	•

*Consists of 0.48% (butylearloity)) (6-propylpiperonyl) ether and 0.12% related compounds.

KEEP OUT OF REACH OF CHILDREN

CAUTION See back panel for additional cautions.

Met Weight: 14 oz.

HALF-ICTURED BY

CSA LIMITED, INC. HOUSTON, TEXAS 77289

EPA Res. No. 7056-21

EPA Est. No. 7056-TX-1

STORAGE: Do not use & Store near Heat or oper flame

DISPOSAL: Wrap container and dispose of in trash. Do not inclmerate.

> PRECAUTICMARY STATEMENTS MASARDS TO HUMANS AND DOMESTIC ANIMALS

CADTION

Avoid contamination of food and food stuffs. Food should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before using. When using the product in these areas apply only when the facility is not in operation. In hospitals, patients should be removed from room prior to treatment. Room should be ventilated for two hours after spraying. Do not return patients to room until after ventilation. Cover or remove fish bowls. Remove all pets from area to be treated.

STATEMENT OF PRACTICAL TREATMENT

If on Skin: In case of contact, remove contaminated clothing and immediately wash skin with soap and water.

If in Eyes: Flush eyes with plenty of water for 15 minutes.

Call a physician. Remove Victim to fresh air. Apply respiration If Inhaled:

if indicated.

CHEMICAL BASABOS

Contents under pressure. Do not use or store mear heat or open flame. Do not puncture or incinerate container. Exposure to temperatures above 130°F may cause bursting.

Not Reviewed. Registrant claims to be in accordance with Fleregistration of Peaticide Product Guidance for Phase 2 Response, Page 2.12, Active to Inert Change in Status.

