

1(11-1)

7056

ON BEEF CATTLE, HORSES: To control Horn flies, Cattle Lice, Ear Tick, Lone Star Tick, Gulf Coast Tick, Winter Tick and Cattle Fever Tick, mix 1 gallon of CHEM-TOX with 100 gallons of water and stir well to form a creamy emulsion. Spray animals with a strong spray to penetrate to the skin, or dip thoroughly. Approximately 2 quarts of spray are required for a full grown cow, proportionately less for smaller animals. Be sure not to overdose young animals. For season long Horn fly control repeat treatment every 21 days. For season long tick control repeat treatments every 30 days.

For maximum long-lasting protection, mix 1 gallon of CHEM-TOX with 100 gallons of water as described above. However, for a lower cost spray for temporary control of cattle lice and horn flies, 1 gallon of CHEM-TOX may be mixed with 150 or 200 gallons of water and applied as above.

ON SHEEP AND GOATS: For control of Sheep Lice, Goat Lice and Keds, mix 1 gallon of CHEM-TOX with 200 to 400 gallons of water, mix well and spray or dip animals thoroughly. Repeat treatment in 14 days if control is not complete.

HOGS: To control sarcoptic mange and hog lice, dilute one gallon of CHEM-TOX to 100 gallons of water, mix well, and spray or wash animals thoroughly. Be sure not to overdose young animals. Repeat treatment in 14 days if control is not complete.

MANGE AND SCAB: For excellent control and prevention of sarcoptic, psoroptic and chorioptic mange (scab), of beef cattle, sheep and hogs, mix 1 gallon of CHEM-TOX with 100 gallons of water and dip animals thoroughly. Repeat treatment in 14 days.

CHEM-TOX

TOXAPHENE-MALATHION

LIVESTOCK SPRAY & DIP

BACKRUBBER CONCENTRATE

For Spraying Livestock, Barns, Sheds, Animal Shelters, Fences and Garbage Cans, for Control of Many Insect Pests.

ACTIVE INGREDIENTS:

Toxaphene*	43.00%
*Technical Chlorinated Camphene (Chlorine Content 67-69%)	
Malathion (0,0-Dimethyl Dithiophosphate of Diethyl Mercaptosuccinate)	4.30%
Xylene	4.79%

INERT INGREDIENTS:

47.91%

WARNING:

KEEP OUT OF REACH OF CHILDREN
See back panel for additional cautions.

NET CONTENTS: 1 GALLON

Mfg by
CHEM SPRAY AEROSOLS, INC.
Houston, Texas 77098

9(142)

WARNING:

May be fatal if swallowed. Do not breathe spray mist. Do not get in eyes, on skin or clothing. In case of spillage on person or clothing, immediately remove clothing and flush skin or eyes with plenty of water; for eyes, get medical attention. Do not apply in dairy barns or to dairy animals. Avoid contamination of feed and foodstuffs. Do not use around or on poultry, dogs, or fish.

This product is toxic to fish, birds, and other wildlife. Keep out of streams, lakes, and ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

Do not reuse container. Break or crush empty container and bury in isolated area away from water supplies. Do not apply in dry areas not specified on the label. Do not use on animals within 3 days of slaughter.

FOR USE ON BACKRUBBERS: As an aid in the control of hornflies and lice on beef cattle, dilute 1 gallon of CHEM TOX with 10 to 20 gallons of fuel oil or other suitable carrier and fill insecticide compartment of backrubber device or soak sack or cloth as occasion requires. Recharge as necessary. Place backrubber device where animals feed, water, or loaf.

AS A PREMISE SPRAY FOR BARNYARD AND BUILDING: To kill Fleas, Mosquitoes, Roaches, Stable Flies and many other insects around barns, barnyards and outbuildings, dilute 1 gallon of CHEM-TOX with 20 gallons of water. Thoroughly wet all surfaces where insects alight or crawl, at the rate not in excess of 4 gallons per 1,000 square feet. For Fleas, thoroughly treat places where animals sleep. Repeat applications as necessary for effective season-long control. Do not treat pets or animals with this concentration. Do not use in dwellings.

CHEM-TOX DILUTION TABLE

AMOUNT OF CHEM TOX	VOLUME OF WATER				
	BEEF CATTLE HORSES HOGS	SHEEPS AND GOATS LICE AND KEYS SPRAY	DIP	SHEEP SCAPES	PREMISE SPRAY
Pint	12 Gal.	25 Gal.	60 Gal.	12 Gal.	2 Gal.
Quart	25 Gal.	50 Gal.	125 Gal.	25 Gal.	5 Gal.
Gallon	100 Gal.	200 Gal.	500 Gal.	100 Gal.	20 Gal.

EPA Reg No. 7056-72
EPA Est. 7056-TX-1

ACCEPTED
1-31-74
7056-72

CAUTION:

As a general precaution in spraying animals, avoid treating during cold, stormy weather; avoid treating overheated or sick animals; water animals before dipping to prevent them from drinking the dip. If calves must be treated, spray with a minimum amount of material. Do not use on very young animals or calves under 3 months of age. Do not use in poultry houses. Malathion is a cholinesterase inhibitor. Do not use this product simultaneously or within a few days before or after exposure to cholinesterase inhibiting drugs, pesticide or chemicals. Atropine is antidotal. DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME.

1 (of 2)

ON BEEF CATTLE, HORSES: To control Horn flies, Cattle Lice, Ear Tick, Lone Star Tick, Gulf Coast Tick, Winter Tick and Cattle Fever Tick, mix 1 gallon of CHEM-TOX with 100 gallons of water and stir well to form a creamy emulsion. Spray animals with a strong spray to penetrate to the skin, or dip thoroughly. Approximately 2 quarts of spray are required for a full grown cow, pro-portionately less for smaller animals. Be sure not to overdose young animals. For season long Horn fly control repeat treatment every 21 days. For season long tick control repeat treatments every 30 days.

For maximum long-lasting protection, mix 1 gallon of CHEM-TOX with 100 gallons of water as described above. However, for a lower cost spray for temporary control of cattle lice and horn flies, 1 gallon of CHEM-TOX may be mixed with 150 or 200 gallons of water and applied as above.

ON SHEEP AND GOATS: For control of Sheep Lice, Goat Lice and Keds, mix 1 gallon of CHEM-TOX with 200 to 400 gallons of water, mix well and spray or dip animals thoroughly. Repeat treatment in 14 days if control is not complete.

HOGS: To control sarcoptic mange and hog lice, dilute one gallon of CHEM-TOX to 100 gallons of water, mix well, and spray or wash animals thoroughly. Be sure not to overdose young animals. Repeat treatment in 14 days if control is not complete.

MANGE AND SCAB: For excellent control and prevention of sarcoptic, psoroptic and chorioptic mange (scab), of beef cattle, sheep and hogs, mix 1 gallon of CHEM-TOX with 100 gallons of water and dip animals thoroughly. Repeat treatment in 14 days.

3(0f2)

WARNING:

May be fatal if swallowed. Do not breathe spray mist. Do not get in eyes, on skin or clothing. In case of spillage on person or clothing, immediately remove clothing and flush skin or eyes with plenty of water; for eyes, get medical attention. Do not apply in dairy barns or to dairy animals. Avoid contamination of feed and foodstuffs. Do not use around or on poultry, dogs, or fish.

This product is toxic to fish, birds, and other wildlife. Keep out of streams, lakes, and ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.


Do not reuse container. Break or crush empty container and bury in isolated area away from water supplies. Do not apply in any manner not specified on the label. Do not use on animals within 28 days of slaughter.

CHEM-TOX DILUTION TABLE

AMOUNT OF CHEM TOX	VOLUME OF WATER				
	BEEF CATTLE, HORSES, HOGS	SHEEPS AND GOATS LICE AND KEDS		SHEEP SCABIES	PREMISE SPRAY
		SPRAY	DIP		
Pint	12 Gal.	25 Gal.	60 Gal.	12 Gal.	2 Gal.
Quart	25 Gal.	50 Gal.	125 Gal.	25 Gal.	5 Gal.
Gallon	100 Gal.	200 Gal.	500 Gal.	100 Gal.	20 Gal.

E.P.A. Reg No. 7056-72

E.P.A. Est. 7056-TX-1



FOR USE ON BACKRUBBERS: As an aid in the control of hornflies and lice on beef cattle, dilute 1 gallon of CHEM TOX with 10 to 20 gallons of fuel oil or other suitable carrier and fill insecticide compartment of backrubber device or soak sack or cloth as occasion requires. Recharge as necessary. Place backrubber device where animals feed, water, or loaf.

AS A PREMISE SPRAY FOR BARNYARD AND BUILDING: To kill Fleas, Mosquitoes, Roaches, Stable Flies and many other insects around barns, barnyards and outbuildings, dilute 1 gallon of CHEM-TOX with 20 gallons of water. Thoroughly wet all surfaces where insects alight or crawl, at the rate not in excess of 4 gallons per 1,000 square feet. For Fleas, thoroughly treat places where animals sleep. Repeat applications as necessary for effective season-long control. Do not treat pets or animals with this concentration. Do not use in dwellings.

CAUTION:

As a general precaution in spraying animals, avoid treating during cold, stormy weather; avoid treating overheated or sick animals; water animals before dipping to prevent them from drinking the dip. If calves must be treated, spray with a minimum amount of material. Do not use on very young animals or calves under 3 months of age. Do not use in poultry houses. Malathion is a cholinesterase inhibitor. Do not use this product simultaneously or within a few days before or after exposure to cholinesterase inhibiting drugs, pesticide or chemicals. Atropine is antidotal. DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME.