

Reg no: 7056-158

PM-15

124

US ENVIRONMENTAL PROTECTION AGENCY OFFICE OF PESTICIDES PROGRAMS REGISTRATION DIVISION (75-767) WASHINGTON, DC 20460 NOTICE OF PESTICIDE: <input type="checkbox"/> REGISTRATION <input type="checkbox"/> REREGISTRATION (Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended)	EPA REGISTRATION NO. 7056-158	DATE OF ISSUANCE JUN 11 1990
	TERM OF ISSUANCE Until Reregistration	
	NAME OF PESTICIDE PRODUCT Inside/Outside Insecticide Spray	

NAME AND ADDRESS OF REGISTRANT (Include ZIP code)

[Faint registration details]

161/265160
18/2

BEST AVAILABLE COPY

NOTE: Changes in labeling formula differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above U.S. EPA registration number.


On the basis of information furnished by the registrant, the above named pesticide is hereby Registered/Reregistered under the Federal Insecticide, Fungicide, and Rodenticide Act.

A copy of the labeling accepted in connection with this Registration/Reregistration is returned herewith.

Registration is in no way to be construed as an indorsement or approval of this product by this Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

[Faint, mostly illegible text, likely containing registration details and correspondence]

ATTACHMENT IS APPLICABLE

SIGNATURE OF APPROVING OFFICIAL 

DATE
June 11, 1990

Handwritten: 6-11-90

- d. Delete the statement, "Nonfood areas are areas where food is not stored, served, prepared or handled" since it conflicts with the definition of non-food areas on the front panel.
- e. Under the instructions for use on dogs/cats add the following precautions:

Not for use on puppies/kittens less than six weeks old. Do not use on old, sick or convalescing animals.

3. Submit five (5) copies of your final printed labeling before you release the product for shipment. Refer to the A-79 Enclosure for a further description of final printed labeling.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

George T. LaRocca
Product Manager 15
Insecticide-Rodenticide Branch
Registration Division (H7505C)

3 9 4 1

INSIDE OUTSIDE
INSECTICIDE SPRAY
NUMBER TWO

A COMBINATION OF INSECTICIDES
OFFERING QUICK KNOCKDOWN
AND
RESIDUAL KILL
OF CRAWLING INSECTS
FOR UP TO FOUR WEEKS

ACTIVE INGREDIENTS:

*†-Permethrin (3-phenoxyphenyl)methyl(+)-cis, trans-3-(2,2-dichloroethyl)-2,2-dimethylcyclopropanecarboxylate 0.15%/w/w
d-Trans Allethrin (allyl homolog of Cyanozin I) 0.25%

INERT INGREDIENTS: 99.60%
100.00%/w/w

*Cis/trans isomers ratio: min. 35%(z)cis and max. 65%(z)trans.
†U.S. Patent No. 4,024,183.
Permet-Bio UCLAF's Prameze brand of permethrin insecticide.
Permet-Bio UCLAF Corp., 400 Sylvan Avenue, Englewood Cliffs, N.J. 07632

PRECAUCION AL CONSUMIDOR: Si usted no lee Ingles, no use este producto hasta que la etiqueta le haya sido explicada amplemente.
(TO THE USER: If you cannot read English, do not use this product until the label has been fully explained to you.)

KEEP OUT OF REACH OF CHILDREN
CAUTION
SEE SIDE PANEL FOR ADDITIONAL
PRECAUTIONARY STATEMENTS AND
STATEMENT OF PRACTICAL TREATMENT

Can be used in non-food areas of food handling establishments such as garbage rooms, lavatories, floor drains (to sewers), entries, vestibules and hallways, offices, locker rooms, machine rooms, basements, boiler rooms, garages, utility rooms, mop closets and storage rooms (after canning or bottling).

Net Weight 8oz

EPA REG. NO. 7056-XX

EPA EST. 7056-TX-1

MANUFACTURED BY

CSA LIMITED, INC
HOUSTON, TX 77269-0347

PRECAUTIONARY STATEMENTS
HAZARDOUS TO HUMANS AND DOMESTIC ANIMALS
CAUTION

Harmful if swallowed. May be absorbed through skin. Wash thoroughly after handling and before smoking or eating foodstuffs. Remove pets and cover fish tanks before use or humans. Do not allow children or pets to contact.

STATEMENT OF PRACTICAL TREATMENT

- IF SWALLOWED: Drink one to two glasses of water and gargle the back of the throat with fingers. If necessary, a physician immediately. Do not induce vomiting.
- IF INHALED: Remove affected person to fresh air.
- IF ON SKIN: Remove contaminated clothing and wash with soap and water.
- IF IN EYES: Flush eyes with plenty of water. Contact a physician immediately.

ENVIRONMENTAL HAZARDS

This product is toxic to fish and other aquatic organisms. Do not contaminate water by cleaning of equipment or disposal.

PHYSICAL AND CHEMICAL HAZARDS

EXTREMELY FLAMMABLE. Do not use or store near fire, pilot lights in gas appliances and unplug or turn off pilot lights. After airing out phase is completed, relight pilot lights. Do not smoke in the use area. Contents under pressure. Exposure to temperatures above 130°F may cause explosion. Do not apply this product in switch boxes or other electrical equipment because it may cause electrical failure or shorting of electrical surfaces that can be harmed or stained by water.

ACCEPTED
with COMMENT
in EPA Letter D-100

JUN 11 1999

Under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA) as amended.

70

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE: Do not store near heat or open flame. Keep in locked storage area. Do not store this water-based formulation at temperatures below 32°F (0°C).

CONTAINER DISPOSAL: Replace cap and wrap container in several layers of newspaper. Discard in trash. Do not puncture or incinerate.

Shake well before each use. Remove protective cap. Aim spray opening away from person. Push button to spray.

Do not use in food areas of food handling establishments, restaurants or other areas where food is commercially prepared or processed. Do not use in serving areas while food is exposed or facility is in operation. Serving areas are areas where prepared foods are served such as dining rooms but excluding areas where food may be prepared or held. In the home, all food processing surfaces and utensils should be covered during treatment or thoroughly washed before use. Exposed food should be covered or removed. Nonfood areas are areas where food is not stored, served, prepared, or handled. This product is for use in nonfood areas indoors and outside premises of hotels, motels, homes, preschools, nurseries, day camps, greenhouses, lathhouses, seed houses, food handling establishments, commercial, industrial, office, institutional and storage buildings, dairies, milk storage rooms, egg processing plants, supermarkets, restaurants, bottling plants, canneries, flour and feed mills, bakeries and hospitals. In hospitals, patients should be removed prior to treatment. Also for use in campers, mobile homes buses, automobiles, trucks, and boxcars. Also for use in dog kennels and horse stables. Not for use in horse stables if horses will be used for food.

PESTS Indoors

For residual and contact control of cockroaches (waterbugs), crickets, silverfish (firebrats), clover mites, millipedes, centipedes, earwigs, sowbugs (pillbugs), spiders, ants, clothes moths, carpet beetles, boxelder bugs, scorpions, spider beetles, grasshoppers, house flies, mosquitoes, gnats, fruit flies, cluster flies, flying moths, confused flour beetles, saw-toothed grain beetles, shiny spider beetles, grain mites, red flour beetles, angoumois grain moths, grain weevils, Indian meal moths, Mediterranean flour moths, lesser grain borers, cigarette beetles, drug store beetles, chocolate moths. Spray thoroughly into hiding places such as cracks, crevices, moist areas, openings around pipes and sinks, under refrigerators, baseboards, and storage areas. Spray directly on insects where possible.

BRAIN INSECTS: For residual use, to control confused flour beetles, saw-toothed grain beetles, spider beetles, shiny spider beetles, grain mites, red flour beetles, angoumois grain moths, grain weevils, Indian meal moths, Mediterranean flour moths, lesser grain borers, cigarette beetles, drug store beetles, chocolate moths, carpet beetles. Avoid contamination of other food and foodstuffs in the area by removing before spraying. Spray directly on exposed stages and into cracks and crevices where grain insects may be hiding. Dispose of all infested products. The shelves and drawers should be cleaned and thoroughly sprayed so as to contact all pests, and covered with shelf paper after allowing shelves to dry. Any utensils that are contacted with spray should be washed before using. Repeat as necessary.

SPIDERS: Spray directly on these pests and use repeated applications in the webbing and all cracks and crevices which harbor spiders.

ANTS: To kill ants, spray trails, nests and points of entry within buildings and homes. Spray directly on ants where possible. Repeat as necessary.

WASPS, HORNETS, YELLOW JACKETS, BEES: Application should be made in the late evening when insects are at rest. Spray liberally including and breeding places, especially under eaves rafters, contacting as many insects as possible. Repeat as necessary.

HOUSE FLIES, MOSQUITOES, CLUSTER FLIES, GNATS, FRUIT FLIES, FLYING MOTHS: Close off all doors and windows. Point nozzle upward. Direct spray mist to all parts of the room, especially windows and other light sources which attract these insects. Fill the room with mist, then leave the treated area. Keep the room closed for at least 15 minutes. Ventilate the room when treatment is completed.

CARPET BEETLES AND CLOTHES MOTHS: To Treat Carpets-Hold container 14-18 inches (36-46 cm) from carpet. Apply spray evenly, covering the entire surface until slightly moist. Spray underside of carpet when possible. Avoid wetting or soaking. To Treat Garments, Blankets, Carpets and Other Fabrics Before Storage-Dry clean or wash before treatment and after storage. Apply spray evenly, covering the material on both sides until slightly moist. For best results, place treated materials in plastic bags after drying, and store in a dark place. Also spray floor and baseboards, closets, cracks, crevices, infested areas of shelving and/or wherever these insects are seen or suspected.

CONTACT APPLICATION ONLY: (Not For Residual Control Of Fleas, Ticks and Bedbugs)

FLEAS, BROWN DOG AND LONE STAR TICKS ON DOGS, CATS AND PREMISES: Spray animals along back, under forelegs and on other areas, except acrotum and anus. Part hair to contact skin

and insure good coverage. Hold nozzle about one foot (30 cm) away while spraying. Apply until hair is thoroughly moist. Do not spray directly in eyes, mouth or nose of animal. To prevent reinfestation, thoroughly spray the animals' bedding and other resting areas until the surface is slightly moist contacting as many insects as possible. Treat the floor areas, cracks and crevices in the floor and along baseboards at the rate of 2 seconds per linear foot (30 cm) at a distance of 12 to 18 inches (30-45 cm). Also spray the entire inside surface of dog houses.

FOR BEDBUGS: Apply as a spot treatment to cracks and crevices, around baseboards, floorboards, bedboards and walls. First thoroughly clean and air mattresses and springs. Then spot treat only areas that may harbor bedbugs, concentrating on tufts, folds and seams. Repeat as necessary. For persistent infestations, consult professional applicator.

PESTS OUT-OF-DOORS

PESTS ON OUTSIDE SURFACES OF BUILDINGS: Use this product to control ants, clover mites, crickets, earwigs, millipedes and sowbugs (pillbugs) on the outside surfaces of buildings. To help prevent infestation of buildings treat the building foundation to a height of 2 to 3 feet (0.5 to 1 m), where pests are active and may find entrance. Also, apply as a residual spray to outside surfaces of buildings, including porches, screens, window frames, eaves, patios, garages, refuse dumps, and other areas where these pests congregate or have been seen. Repeat treatment as needed to maintain effectiveness.

HOUSE FLIES, MOSQUITOES, CLUSTER FLIES, GNATS, FLYING MOTHS: For outdoor use only as aid in reducing annoyance from these insects. Spray outside surfaces of screens, doors, window frames or wherever these insects may enter the room. Also treat surfaces around light fixtures on porches, in garages, and other places where these insects alight or congregate. Repeat as necessary.

ANTS, LONE STAR TICKS, CRICKETS, FLEAS, AND EARWIGS: For treatment of localized infestations of these insects in outdoor areas where there are weeds or bushy non-crop areas, spray infested areas thoroughly. For ants, thoroughly wet hills and runways. Repeat application as infestations warrant and as reinfestations occur.

WASPS, HORNETS, YELLOW JACKETS, BEES: Application should be made in late evening when insects are at rest. Aim spray at nest openings in ground, bushes, and in cracks and crevices which may harbor nests, saturating nest openings and contacting as many insects as possible.

FOR USE ON PLANTS

CONTACT AND RESIDUAL SPRAY- Apply spray to plants covering all foliage. Avoid heavy wetting.

INDOORS: To protect plants in greenhouses and lathhouses, and for indoor landscaping-Spray directly on insects to kill leafrollers, armyworms, cutworms, cabbage worms on chrysanthemums; fungus gnats on velvet plants. Spray as directed above. Spray as directed above to control aphids, armyworms, corn earworms fall cankerworms, inchworms, Japanese beetles, leafminers, leafrollers, and whiteflies on the following plants.

Ageratum	English Ivy	Piggy-back plant
Ardisia	Fuchsia	Poinsettia
Aster	Gladious	Portulaca
Azalea	Grape Ivy	Poinos

Baby's breath	Hypoestes	Prayer Plant
Begonia	Ivy	Purple Passion
Birds-nest fern	Juniper	Rabbit's foot fern
Bleeding heart	Lily	Rose
Carnation	Marigold	Schefflera
Coleus	Orchid	Snagdragon
Crown of thorns	Palm	Snake Plant
Cockscomb	Pansy	Statice
Cyclamen	Piperomia	Velvet Plant
Dracaena	Petunia	Verbena
Dumbcane	Philodendron	Zinnia

If application is to be made indoors, remove infested plants from living areas prior to application. Spray plants in an area not likely to be occupied by children or pets (particularly fish). When leaves dry, plants can be returned to original location.

Use with CAUTION on ferns, palms and luchsia. As with all plants listed above, especially carnations, chrysanthemums and roses, there exists a wide variety of species and cultivars, and therefore, a wide variety of susceptibility to injury. To ascertain possible phytotoxicity, spray a few plants as described above, wait 24 hours, and check for possible injury.

Repeat applications may be necessary to control citrus black flies on non-bearing citrus; adult Japanese beetles on roses.

OUTDOORS: Spray as described above in the late afternoon or evening, when the temperature ranges from 50°F to 75°F and when there is little or no wind. Spray at the first indication of insect pests. On those plants that will flower during the season, the spray schedule should be completed before blossom formation. Repeat as necessary; use intervals of 4 to 8 days. PRAMEX application can be made up to the day of collection.

*Spray as directed above to control aphids, cicadas, corn earworms, fall cankerworms, inchworms, leafminers, leafrollers, mealybugs, spider mites, exposed thrips and whiteflies on the following plants:

Ageratum	Gold bells	Pea shrub
Aster	Hypoestes	Petunia
Azalea	Ivy	Philodendron
Begonia	Juniper	Poinsettia
Coleus	Lilac	Rhododendron
Common ninebark	Marigold	Rose
Common snowberry	Mock Orange	Snagdragon
Excucum	Nannyberry	Taxus
Euonymus	Orchid	Wax plants
English Ivy	Palm	Weeping fig
Gladious	Pansy	Zinnia

TO KILL TENT CATERPILLARS: Application should be made when caterpillars are young and tents are first noticed (usually late spring). For best results, apply in the late afternoon or evening when caterpillars have returned to their tents. Aim spray directly at tents, and saturate on all sides. Repeat as necessary.